## Correction to "Characterization of human $\alpha 4\beta 2$ -nicotinic acetylcholine receptors stably and heterologously expressed in native nicotinic receptor-null SH-EP1 human epithelial cells"

In the above article [Eaton JB, Peng J-H, Schroeder KM, George AA, Fryer JD, Krishnan C, Buhlman L, Kuo Y-P, Steinlein O, and Lukas RJ (2003) Mol Pharmacol 64:1283-1294], Table 3 was incorrect due to printing errors. The corrected table appears below.

We regret this error and apologize for any confusion or inconvenience it may have caused.

## TABLE 3 Parameters for interactions of indicated drugs with $h\alpha 4\beta 2$ -nAChR in transfected SH-EP1 cells

ts presented in Figs. 6 to 9 and in Tables 1 onal $\rm IC_{50}$ values given in parentheses for $\rm r$ BDN binding competition $K_{\rm i}$ values (column e indicated ligands (drug; column 1). Also in alue but corrected for the concentration of E totations following the drug name for selectitive inhibitor (C), or a mixed competitive	nicotine and suberyldicholin 4), all in micromolar, are ex ndicated (F/B; column 5) is t I-EBDN used in those assay: ted ligands indicate, from /noncompetitive inhibitor (N	e from fits to the self-inhibit pressed to allow comparisons. the ratio between the function s and for the H-EBDN binding results shown in Fig. 9, function.	ory phase of agonist dose-response probetween functional and radioligand bin at $\mathrm{EC_{50}/IC_{50}}$ and the binding competing $K_{\mathrm{D}}$ of 10 pM using the Cheng-Prusoff	offiles for those drugs), and ading competition affinities tion $K_i$ value (based on the correction) for each ligand.
Drug	Functional		$\begin{array}{c} \text{Binding Competition} \\ K_{\text{i}} \end{array}$	F/B
	$\mathrm{EC}_{50}$	$IC_{50}$	11	
		$\mu M$	$\mu M$	
Epibatidine	0.0085		0.000012	710
Nicotine	0.85	(5800)	0.0020	430
Cytisine	1.3		0.00024	5400
Suberyldicholine	1.3	360 (200)	0.013	100
Acetylcholine	1.7		0.018	94
Dimethyl-phenyl-piperazinium	1.9		0.020	95
Carbamylcholine	17		0.15	110
Acetylthiocholine	100		0.17	590
Choline	$\sim$ 3800		7.0	>540
Lobeline		21	0.0019	11,000
Succinylcholine		≫1000	2.9	>340
Decamethonium C		300	1.3	230
D-tubocurarine N/C		62	2.0	31
Dihydro- $\beta$ -erythroidine C		1.5	0.085	18
Methyllycaconitine C		6.6	2.3	2.8
Eserine N		130	>24	< 5.4
Vecuronium		28	24	1.2
Alcuronium		26	19	1.4
Pancuronium N/C		91	57	1.6
Hexamethonium N/C		11	45	0.24
Mecamylamine N		0.47	$\gg$ 24	< 0.02