

Correction to “Preferential formation of MT1/MT2 melatonin receptor heterodimers with distinct ligand interaction properties compared with MT2 homodimers”

In the above article [Ayoub MA, Levoe A, Delagrangre P, and Jockers R (2004) *Mol Pharmacol* **66**:312–321], information regarding the ligands in Fig. 5 was removed in error during copyediting. A corrected version of Fig. 5 appears below. The online version has been corrected in departure from the print version.

We regret this error and apologize for any confusion or inconvenience it may have caused.




Fig. 5. Correlations between EC_{50} values of MT₁/MT₂ heterodimers and K_i values of MT₁ and MT₂ receptors. EC_{50} values of the MT₁/MT₂ heterodimer shown in Table 4 were plotted against K_i values determined in cells coexpressing MT₁ and MT₂ receptors (A) or expressing MT₁-Rluc (B) or MT₂-YFP (C) receptors individually. D, correlation between EC_{50} values of the MT₂ homodimer and the MT₁/MT₂ heterodimer. Ligands: 1, melatonin; 2, S20098; 3, S22153; 4, S20928; 5, 4P-PDOT; 6, luzindole; 7, S24773; 8, 2-iodomelatonin. B and C, nonselective (●) and selective (○) compounds. Ligand 9 shown in B corresponds to the MT₁-selective S26284 compound. Data were fitted using a linear regression equation (GraphPad Prism software).