

Blockade of Vascular Endothelial Growth Factor Receptor Signal Pathway and Antitumor Activity of ON-III (2',4'-Dihydroxy-6'-methoxy-3',5'-dimethylchalcone), a Component from Chinese Herbal Medicine

Xiao-Feng Zhu, Bin-Fen Xie, Jun-Min Zhou, Gong-Kan Feng, Zong-Chao Liu, Xiao-Yi Wei, Feng-Xian Zhang, Mei-Fang Liu, and Yi-Xin Zeng

Department of Experimental Research, Cancer Center, Sun Yat-sen University, Guangzhou, China (X.-F.Z., B.-F.X., J.-M.Z., G.-K.F., Z.-C.L., Y.-X.Z.); and Southern China Institute of Botany, the Chinese Academy of Sciences, Guangzhou, China (X.-Y.W., F.-X.Z., M.-F.L.)

Received November 30, 2004; accepted February 8, 2005

ABSTRACT

Antiangiogenesis is a promising strategy of cancer treatment. Vascular endothelial growth factor receptor [fetal liver kinase/kinase-inserting domain-containing receptor (KDR)] is a tyrosine kinase receptor and has been strongly implicated in tumor angiogenesis. In this study, we report that 2',4'-dihydroxy-6'-methoxy-3',5'-dimethylchalcone (ON-III), extracted from the dried flower *Cleistocalyx operculatus*, used in traditional Chinese medicine, reversibly inhibited KDR tyrosine kinase phosphorylation, but epidermal growth factor receptor tyrosine kinase phosphorylation was unaffected under the same concentrations of ON-III. ON-III also inhibited mitogen-activated protein kinase (MAPK) and AKT activation of KDR signal transduction in downstream molecules without reduced

total MAPK and AKT. The results in vitro showed that ON-III inhibited growth of human vascular endothelial HDMEC cells in the presence of VEGF preferentially, compared with epidermal growth factor. Systemic administration of ON-III at nontoxic doses in nude mice resulted in inhibition of subcutaneous tumor growth of human hepatocarcinoma Bel7402 and lung cancer GLC-82 xenografts. The tumor vessel density decreased, as determined by immunohistochemical staining, for CD31 after ON-III treatment. These results indicated that ON-III inhibited KDR tyrosine kinase, shut down KDR-mediated signal transduction, and inhibited tumor growth of human xenografts in vivo.

Angiogenesis, the formation of new blood vessels by sprouting from pre-existing endothelium, is a significant component of a wide variety of biological processes, including embryonic vascular development and differentiation, wound healing, organ regeneration, and pathological processes including tumorigenesis (Folkman, 1990; Eberhard et al., 2000). Many growth factors involve this process. Of the nu-

merous growth factors and cytokines that have been shown to have angiogenic effects, VEGF seems to be a pivotal factor in pathological situations that involve neovascularization (Smith et al., 2000; Foekens et al., 2001). VEGF is produced by normal and transformed cells and plays a key role in the physiology of normal vasculature and in tumor-induced angiogenesis, which makes it important to understand the mechanisms through which this mitogen promotes cell proliferation (Nakopoulou et al., 2002).

VEGF first binds to either of two tyrosine kinase receptors, Flk1/KDR or Flt1. Signaling by such receptors facilitates activation of the intrinsic tyrosine kinase followed by autophosphorylation of tyrosine residues in the cytoplasmic sig-

This study was supported by grants 39900183 and 30171092 from the National Natural Science Foundation of China, and grants 013126, 2003C30101, and 2004Z3-E4021 from the Natural Science Foundation of Guangdong Province.

Article, publication date, and citation information can be found at <http://molpharm.aspetjournals.org>.
doi:10.1124/mol.104.009894.

ABBREVIATIONS: VEGF, vascular endothelial growth factor; KDR, kinase inserting domain-containing receptor; MAPK, mitogen-activated protein kinase; VEGFR, vascular endothelial growth factor receptor; EGFR, epidermal growth factor receptor; HDMEC, human dermal microvessel endothelial cells; ON-III, 2',4'-dihydroxy-6'-methoxy-3',5'-dimethylchalcone; DMSO, dimethyl sulfoxide; PBS, phosphate-buffered saline; TBST, Tris-buffered saline/Tween 20; MTT, 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide; PAGE, polyacrylamide gel electrophoresis; PTK, protein tyrosine kinase; SU5416, 3-[(2,4-dimethylpyrrol-5-yl)methylidene]-indolin-2-one; SU6668, (Z)-3-[2,4-dimethyl-5-(2-oxo-1,2-dihydro-3-ylidenemethyl)-1H-pyrrol-3-yl]-propionic acid; ZD6474, N-(4-bromo-2-fluorophenyl)-6-methoxy-7-[(1-methylpiperidin-4-yl)methoxy]quinazolin-4-amine; PTK787/ZK222584, 1-[4-chloroanilino]-4-[4-pyridylmethyl]phthalazine succinate; PTY, phosphotyrosine.

naling molecules that connect the activated receptor to transduction cascades and promote cellular responses (Geng et al., 2001; Colavitti et al., 2002; Dias et al., 2002). KDR receptor possesses intrinsic tyrosine kinase activity that is stimulated after ligand binding and receptor dimerization and is mandatory for transmission of a cytoplasmic signaling response (Binetruy-Tournaire et al., 2000; Zhu et al., 2001). The activation of KDR has also been shown to correlate with tumor growth, lymph node metastasis, and resistance to chemotherapy in many kinds of tumors (Thakker et al., 1999; Baek et al., 2000). Binding of VEGF to KDR receptor on the surface of endothelial cells facilitates its autophosphorylation of the protein tyrosine kinase domain (Takahashi et al., 1999; Ichikura et al., 2001; Zhang et al., 2002). Activation of KDR by VEGF results in activation of phosphatidylinositol 3-kinase/AKT, MAPK, and protein kinase C; the ultimate cellular response is DNA synthesis and cell proliferation. Vascular endothelial growth factor receptors (VEGFRs) are up-regulated on endothelium at sites of active angiogenesis, providing an opportunity for selective therapeutic intervention (Hanahan, 1997; Zhang et al., 2002).

To exploit targeting at KDR tyrosine kinase for cancer treatment, we have been actively pursuing small molecule therapeutic strategies targeted at KDR receptor-mediated signal transduction pathway (Spiekermann et al., 2002; Wedge et al., 2002). In this report, we describe that ON-III, which is a chalcone derivative from the dried flower *Cleisto-calyx operculatus*, used in traditional Chinese medicine, could inhibit tyrosine phosphorylation of KDR and growth of human cancer xenografts. Although some synthetic molecules have been reported to inhibit the VEGFR kinase, this is a low molecular weight inhibitor of the VEGF/VEGF receptor system reported to be active at inhibiting VEGF-mediated processes from traditional Chinese medicine.

Materials and Methods

Cell Culture and Reagents. The human dermal microvessel endothelial cells (HDMEC) were prepared as described previously (Qian et al., 1999). HDMEC cells, human lung adenocarcinoma GLC-82 cells, and hepatocarcinoma cell line Bel7402 cells were cultivated in RMPI 1640 medium (Invitrogen, Carlsbad, CA) (Zhu et al., 2000). Cells were grown in an incubator at 37°C under 5% CO₂ in air. KDR, anti-EGFR, and anti-tyrosine antibodies (PY99) were obtained from Santa Cruz Biotechnology (Santa Cruz, CA). Anti-Akt, phospho-Akt, MAPK, and phospho-MAPK antibody were purchased from Cell Signaling (Beverly, MA).

Extraction and Isolation. ON-III was extracted and isolated from the flower *C. operculatus* used in traditional Chinese medicine. Its structure is shown in Fig. 1. The following is the steps of extraction and isolation of ON-III. The fresh buds (5.0 kg) of *C. operculatus*,

collected at South China Botanical Garden, Guangzhou, China, were extracted with 95% EtOH three times at room temperature. The EtOH extract, after concentration in vacuo, was suspended in water and the aqueous suspension was sequentially extracted three times each with ether, EtOAc, and *n*-BuOH. The combined EtOAc solution, upon evaporation, yielded a brown syrup (13.0 g). This syrup was subjected to silica gel column chromatography, eluted with CHCl₃-MeOH mixtures of increasing polarities (99.5:0.5 to 97:3). The fractions obtained on elution with CHCl₃-MeOH (99.5:0.5) were combined and concentrated to afford a brown residue. This residue, upon recrystallization in benzene, yielded compound (690 mg): orange needles, melting point 124 to 126°C. The compound was identified by co-thin-layer chromatography with an authentic sample and by direct comparison of its spectral data with those reported previously (Zhang et al., 1990).

Immunoblot Analysis. After the different treatments, cells were placed on ice and washed twice with ice-cold PBS. Cell lysates were harvested in lysis buffer (50 mM HEPES, 150 mM NaCl, 10% glycerol, 1% Triton X-100, 1.5 mM MgCl₂, and 1 mM EGTA) with protein inhibitors (10 µg/ml trypsin, 10 µg/ml aprotinin, 10 µg/ml leupeptin, 1 mM phenylmethylsulfonyl fluoride, 200 µM Na₂VO₄, and 100 mM NaF). After 10 min of centrifugation at 14,000g, the supernatant was transferred to clean microcentrifuge tubes, protein was determined by bicinchoninic acid protein assay (Pierce, Rockford, IL). Equal amounts of protein were then boiled in 5× sample buffer for 5 min at 90°C and resolved on 10% Tris-glycine gels (Novex, San Diego, CA). After electrophoresis, protein was electrotransferred to nitrocellulose membrane (Amersham Biosciences, Piscataway, NJ). Membrane was blocked with 5% milk in TBST (10 mM Tris, pH 7.4, 150 mM NaCl, and 0.05% Tween 20) for 40 min at room temperature and then incubated with a 1:2000 dilution of primary antibody in 1% milk in TBST overnight at 4°C. After washing with TBST three times, the membrane was probed with a 1:2000 dilution of anti-mouse or anti-rabbit antibody conjugated with horseradish peroxidase (Santa Cruz Biotechnology, San Francisco, CA) for 30 min at room temperature. Washing with TBST three times again, the membrane was developed by enhanced chemiluminescence (ECL; Amersham Biosciences).

Immunoprecipitation. Cells were washed twice with ice-cold PBS and then lysed with lysis buffer for 20 min at 4°C, and the protein content of the supernatants was determined by the bicinchoninic acid method (Pierce). Five hundred micrograms of protein was immunoprecipitated at 4°C overnight using the KDR or EGFR antibody (10 µg) and 40 µl of agarose-protein G (Roche Molecular Biochemicals, Mannheim, Germany). After being washed twice with lysis buffer, the KDR or EGFR immune complex was resolved by SDS-PAGE, and immunoblot assay was performed as described above.

Phosphorylation Assays. Cells were seeded in 24-well plate in 10% fetal bovine serum/RMPI 1640 medium. On the next day, the cells were washed twice with serum-free RMPI 1640 medium and starved overnight (14 h). On the third day, cells were treated by different concentrations of compounds for 30 min at 37°C before the cells were stimulated by VEGF for 5 min. Then, cell lysates were collected. Immunoprecipitation was conducted using anti-KDR or anti-EGFR antibodies; KDR, EGFR, phosphorylated KDR, or phosphorylated EGFR was examined using anti-PTY or anti-KDR or anti-EGFR antibodies through immunoblot analysis as shown under Immunoblot Analysis.

Reversibility Test Protocol. We have used the protocol described by Smaill et al. (1999). Treated cells were washed with warm serum-free media, incubated for 30 min, washed again, and incubated for another 8 h. This set of cells was tested for phosphorylation.

Cell Viability Assay. In brief, cells were seeded in 96-well plates (Falcon, Lincoln Park, NJ) at various densities per well. On the next day, compound was prepared at 2-fold dilutions in medium and incubated with the cells until the control cells reached 100% confluence. Cell viability was determined by MTT assay. In brief, MTT (Sigma-Aldrich, St. Louis, MO) was dissolved and sterilized in PBS




Fig. 1. The structure of ON-III.

at 5 mg/ml, and 10 μ l was added into each well. The plate was incubated in incubator (37°C, 5% CO₂) for 4 h, and all the medium was removed. DMSO (100 μ l) was added to each well to dissolve the dark blue crystal, and the plate was shaken gently for 5 min. The optical density values were read with a test wavelength of 570 nm and a reference wavelength of 650 nm (Zhu et al., 2003).

In Vivo Antitumor Activity. The human liver and lung cancer models were established in nude mice. BALB/c nude mice were obtained from the Experimental Animal Center at Sun Yat-sen University. Female mice were 4 to 6 weeks old. All manipulations were performed under sterile conditions. The procedures involving mice and their care were in accordance with the National Institutes of Health *Guide for the Care and Use of Laboratory Animals* and with the UKCCCR (1998). Tumor xenografts were established by 1×10^6 Bel7402 cells injected s.c. into nude mice. Tumors were measured with a caliper in larger diameter and smaller diameter. The tumor volume was calculated by the formula $1/2 \times \text{larger diameter} \times (\text{smaller diameter})^2$. Treatments were initiated on day 7 after inoculation, by which time the the volume had reached $\sim 20 \text{ mm}^3$. ON-III was tested alone in mice bearing transplanted Bel7402 cells and GLC-82 cells for determining the efficacious dosage for these tumors. ON-III was given at 5.0, 10.0, and 20.0 mg/kg i.p. five times a week for 4 weeks. Approximate tumor sizes and body weight were measured twice each week. Average volumes and standard deviations were calculated for each group and plotted. Finally, the mice were sacrificed, and the tumor tissues were excised and weighed. The growth inhibitory effect was calculated with the formula: % inhibitory rate = (T control - T drug)/T control $\times 100\%$.

Quantification of Tumor Vessel Counts. To determine whether tumor growth inhibition by ON-III was associated with inhibition of tumor vessel formation, groups of mice bearing the Bel7402 hepatocarcinoma were used for histological examination. The tumor-bearing mice were treated with either 20 mg/kg/day of ON-III or vehicle. Two weeks after starting treatment, the tumor tissues was quickly frozen and stored at -70°C . Frozen sections were

fixed and stained with primary antibodies to CD31. Five random 0.159-mm² fields at 200 \times magnification were captured for each tumor by using a Sony three-chip camera mounted on a Zeiss universal microscope (Dreves et al., 2000; Laird et al., 2002).

Results

The Effect of ON-III on Tyrosine Phosphorylation of KDR. Autophosphorylation of KDR increases the velocity of the kinase reaction, and only phosphorylated KDR can provide binding sites for its substrates (such as phospholipase C γ , Grb-2, Shc, etc.). Thus, we first detected that ON-III inhibited tyrosine phosphorylation of KDR. To examine the relationship of dose-effect of inhibition, HDMEC cells were treated for 30 min in serum-free medium with the concentrations of ON-III indicated in Fig. 2, then stimulated with 10 ng/ml VEGF for 5 min. Cells were lysed and immunoprecipitated using KDR antibody, then resolved in SDS-PAGE. Western blot analysis was conducted with anti-phosphotyrosine antibody and KDR antibody. The results suggested that ON-III significantly inhibited phosphorylation of KDR in a dose-dependent manner at concentrations of 0, 2.5, 5.0, 10.0, and 20.0 μ g/ml; the IC₅₀ value was 6.20 μ g/ml in HDMEC cells. But levels of KDR protein were unaffected after ON-III treatment (Fig. 2).

The Selectivity of Inhibition of KDR Phosphorylation by ON-III. ON-III inhibited tyrosine phosphorylation of KDR. It is of interest to investigate the relative selectivity of ON-III toward KDR. To elucidate this issue, human epidermal EGFR-overexpressing A431 cells were employed. A431 cells were treated with 10 and 40 μ g/ml ON-III for 30 min and then lysed and immunoprecipitated using EGFR antibody and resolved on SDS-PAGE. Western blot analysis was conducted with anti-phosphotyrosine antibody and EGFR antibody. The results showed that ON-III had no inhibitory effect on tyrosine phosphorylation of EGFR after more than 40 μ g/ml of ON-III treatment for 30 min in A431 cells (Fig. 3).

Reversible Inhibition of KDR Tyrosine Kinase Phosphorylation by ON-III. The results shown above indicate




Fig. 2. Inhibition of KDR phosphorylation by ON-III in HDMEC cells. HDMEC cells were treated with different concentrations of ON-III for 30 min. Cells were collected, lysed, and immunoprecipitated using KDR antibody, then resolved in SDS-PAGE. Western blot analysis was conducted and probed with anti-KDR or anti-PTY antibody. Bottom shows the ratios of protein quantitation of phospho-KDR to KDR. This experiment is representative of three experiments.




Fig. 3. Inhibition of EGFR phosphorylation by ON-III. A431 cells were treated with 0 (lane 1), 0.2% DMSO (lane 2), 10 μ g/ml ON-III (lane 3), and 40 μ g/ml ON-III (lane 4) for 30 min. Cells were collected, lysed, and immunoprecipitated using EGFR antibody, then resolved in SDS-PAGE. Western blot analysis was conducted and probed with anti-EGFR or anti-PTY antibody. This experiment is representative of two experiments.




Fig. 4. Reversible inhibition of KDR phosphorylation by ON-III. HDMEC cells were treated with 0 (lane 1), vehicle (lane 2), or 10 μ g/ml ON-III for 30 min (lane 3), then were washed and incubated for 30 min (lane 4) or 8 h (5) with fresh medium. Immunoprecipitation was performed with KDR antibody. Western blot analysis was performed and probed with anti-PTY or anti-KDR antibody. This experiment is repeated three times.

that ON-III selectively inhibits tyrosine phosphorylation of KDR receptor. To investigate whether inhibition of tyrosine phosphorylation of KDR by ON-III is reversible, HDMEC cells were treated with 10 $\mu\text{g/ml}$ ON-III in serum-free medium for 30 min, and the cells were washed with serum-free medium twice, then incubated separately for 30 min and 8 h. The tyrosine phosphorylation of KDR was detected. The results showed that the levels of phosphorylation of KDR completely recovered 8 h after ON-III was washed out (Fig. 4).

Effect of ON-III on the Activation of MAPK and AKT, Downstream Targets of the KDR Signal Pathway. Activation of KDR leads to endothelial cell proliferation, presumably by inducing activation of extracellular signal-regulated kinase 1/2 and phosphatidylinositol 3-kinase/AKT pathway. To examine whether ON-III can inhibit activation of MAPK and AKT, HDMEC cells were treated with varying concentrations of ON-III for 30 min, and phospho-MAPK and MAPK were detected with anti-phospho-MAPK and anti-MAPK antibody in HDMEC cells. Phospho-Akt and Akt were detected with anti-phospho-AKT and anti-AKT antibody. Phospho-MAPK and phospho-Akt were the active types of MAPK and Akt, respectively. The results showed that phospho-MAPK and phospho-Akt decreased significantly after treatment of HDMEC cells with ON-III (Fig. 5), but total




Fig. 5. Inhibition of MAPK and AKT activation by ON-III. HDMEC cells were treated with different concentrations of ON-III for 30 min in the presence of VEGF. Cells were collected and lysed. Western blot analysis was conducted and probed with anti-phospho-MAPK, anti-MAPK, anti-phospho-AKT, or anti-AKT antibody. This experiment is representative of three experiments.




Fig. 6. Inhibitory effect of ON-III on the growth of HDMEC cells in the presence of VEGF or EGF. HDMEC cells were treated with different concentrations of ON-III in the presence of VEGF or EGF for 72 h. Cell viability was determined with MTT assay and shown as cell viability rate \pm S.E. ($n = 4$).

MAPK and Akt were unaffected. However, phospho-MAPK and phospho-AKT in A431 were unaffected by ON-III (data not shown). This suggested that ON-III has potent inhibitory effect on activation of MAPK and Akt in a dose-dependent manner in VEGF-stimulating HDMEC cells after treatment with ON-III for 30 min.

Effect of ON-III on the Growth of Human Endothelial Cells in the Presence of VEGF. The results described above suggested that ON-III inhibited tyrosine phosphorylation of KDR in HDMEC cells. Can ON-III inhibit cell growth




Fig. 7. The inhibitory effect of ON-III on the growth of human hepatocarcinoma cell Bel7402 (A) and lung adenocarcinoma GLC-82 (B) xenograft tumors. Bel7402 and GLC-82 cells were implanted s.c. in athymic mice. Treatment was started when palpable primary tumors reached a size of $\sim 0.16 \text{ cm}^3$. Once-daily i.p. administration of ON-III or vehicle was then started and continued for the duration of the experiment. Data points represent a mean tumor volume \pm S.E. from 10 mice. A, ON-III has significant inhibitory effect on hepatocarcinoma cell growth with inhibition rates of 38%, 45%, and 65% at dosages of 5, 10, and 20 mg/kg, respectively ($P < 0.05$). B, ON-III inhibited lung cancer cell growth with inhibitory rates of 21.8% ($P > 0.05$) or 40% ($P < 0.05$) at dosages of 5 or 20 mg/kg, respectively.

more potently in the presence of VEGF? To elucidate this issue, HDMEC cells and MTT analysis were employed. The results shown in Fig. 6 indicate that ON-III had more potential inhibitory effect on HDMEC cells in the presence of VEGF. It indicated that ON-III inhibited VEGF-mediated cell growth in HDMEC cells.

Antitumor Activity of ON-III in Vivo. ON-III inhibited KDR tyrosine kinase phosphorylation and KDR-mediated signal transduction. This suggested that ON-III might inhibit the tumor growth in vivo. Therefore, we established human liver cancer and lung adenocarcinoma xenografts to detect antitumor activity of ON-III in vivo. ON-III administration was initiated on day 7 after implantation, when the tumor size was approximately $\sim 20 \text{ mm}^3$. ON-III was alone given by i.p. injection every day at doses of 5, 10, or 20 mg/kg, and tumor growth was inhibited with 38%, 45%, and 65%, respectively, after 4 weeks of treatment. There were significant differences ($P < 0.05$) compared with control (Fig. 7a). No obvious toxicity was observed in mice receiving the dosage treatment. ON-III had no effect on body weight of nude mice

TABLE 1

Effect of ON-III on body weight of nude mice bearing hepatocarcinoma cell BEL7402

The treatment of ON-III has no significant effect on body weight of nude mice compared with DMSO group, $P > 0.05$.

Groups	Before Treatment		After Treatment	
	No. of Mice	Body Weight	No. of Mice	Body Weight
		<i>g</i>		<i>g</i>
NS	10	23.1 ± 1.84	10	28.1 ± 2.46
DMSO	10	23.4 ± 1.93	10	26.3 ± 1.67
ON-III				
5 mg/kg	10	22.8 ± 1.35	10	26.2 ± 1.39
10 mg/kg	10	23.1 ± 1.69	10	26.9 ± 2.52
20 mg/kg	10	23.6 ± 2.61	10	24.4 ± 2.33

NS, 0.9% NaCl solution.




Fig. 8. Analysis of microvessel density of ON-III-treated Bel7402 tumor xenografts ($n = 5$). A, histological sections of control (vehicle-treated) and ON-III (20 mg/kg/day)-treated Bel7402 tumors stained for CD31 and presented at $200\times$ magnification. A significant reduction in CD31 (endothelial cells) staining was evident after ON-III treatment. Mean CD31-positive area/ $5000\text{-}\mu\text{m}^2$ viable tumor (B), as determined by morphometric image analysis after ON-III treatment, are shown. $P < 0.01$ by one-tailed t test.

bearing hepatocarcinoma cell Bel7402 (Table 1). In addition, ON-III inhibited human lung cancer GLC-82 xenografts in nude mice. The inhibitory rates were 21.8% ($P > 0.05$) and 40% ($P < 0.01$) after 4 weeks treatment at dose of 5 and 20 mg/kg ON-III, respectively (Fig. 7b).

Effect of ON-III on Tumor Vascularization. ON-III can inhibit VEGFR tyrosine kinase phosphorylation. Thus, inhibition of tumor growth in vivo by ON-III may occur by inhibiting tumor angiogenesis. Therefore, we employed immunohistochemical staining for CD31 to detect vessels in tumor tissues. The results revealed a significant decrease in tumor vessel counts in the ON-III groups compared with those in the control group (Fig. 8).

Discussion

Tumor growth depends on angiogenesis. This angiogenic switch is characterized by oncogene-driven tumor expression of pro-angiogenic proteins, such as VEGF, basic fibroblast growth factor, interleukin-8, and others. The significance of VEGF/VEGFR signaling in tumor initiation, progression, and metastasis make them important targets for development of specific inhibitors (Blagosklonny, 2004). Activation of KDR by VEGF results in its receptor tyrosine kinase phosphorylation. Phosphorylated KDR receptor provides binding sites for SH-2 domain-containing protein, including the adaptor proteins Grb-2 and Shc. In addition to SH-2 domain, Grb-2 binds to the small nucleotide exchange proteins, such as Sos through its SH-3 domain, then activates ras by exchanging GTP for GDP on ras (Jeong et al., 2002). Activated ras binds to and facilitates raf activation. Activated raf stimulates extracellular signal-regulated kinase activity, which, in turn, activates the MAPK pathway. Inhibiting VEGF production and blocking VEGF/VEGFR signaling can inhibit tumor angiogenesis and tumor growth in vivo (Teicher et al., 2001; Keyes et al., 2004). Although some small

molecules targeting KDR tyrosine kinase (e.g., SU5416, SU6668, ZD6474, and PTK787) have been reported, it is very interesting to identify VEGFR tyrosine kinase inhibitors from traditional Chinese medicine because it has usually no or low toxicity to host (Carlomagno et al., 2002; Mendel et al., 2003; O'Farrell et al., 2003).

KDR tyrosine kinase is a potent target for cancer treatment (Huang et al., 2002; Itokawa et al., 2002). In this study, we found that ON-III, which is one of the chalcone derivatives from the dried flower *C. operculatus*, inhibited tyrosine phosphorylation of KDR. But the expression of KDR protein was unaffected. This indicates that inhibition of phosphorylation of KDR was not via reduced KDR expression. The data demonstrated that inhibition of KDR phosphorylation by ON-III was reversible and relatively selective compared with EGFR tyrosine kinase phosphorylation. Our results demonstrated that ON-III inhibited activation of MAPK and Akt and shut down KDR downstream signaling. ON-III inhibited VEGF-stimulating HDMEC cell proliferation significantly. In comparison, growth inhibition of HDMEC cells by ON-III in the presence of EGF, but without VEGF, was obviously weaker. Moreover, ON-III produced a dose-dependent inhibition of tumor growth in human tumor xenograft models (hepatocarcinoma Bel7402 and lung adenocarcinoma GLC-82), despite their varied histological origin (liver and lung, respectively) and different growth inhibitory rates. In both cases, the percentage of inhibition on tumor growth kept increasing as the duration of drug application was extended, which is indicative of a sustained antitumor effect of ON-III. Furthermore, histological examination showed that tumor vessel counts decreased in tumor tissues in ON-III-treated mice. This may be the result of VEGF signaling blockage. In all of the in vivo experiments described in this report, ON-III was extremely well tolerated, with no obvious effects on body weight or animal behavior, and no target organ toxicity was observed.

From this experiment, we found that efficacy of ON-III after daily i.p. dosing was dependent on the growth rate of the tumors and was more optimal against slowly growing tumors and more variable against quickly growing tumors. Our findings that ON-III is less effective against lung carcinoma GLC-82 than hepatocarcinoma cancer Bel7402 is consistent with studies reported using the tyrosine kinase inhibitors SU5416 and PTK787/ZK222584 and neutralizing antibodies against VEGF (Laird et al., 2000; Mendel et al., 2000). These inhibitors are also less effective against some tumors than others, perhaps because of differences in the requirement of new blood vessels for the growth of particular tumors and the angiogenic factors produced by a particular tumor cell population. Although others have shown that almost all tumor cell lines in vitro and tumors grown in vivo produce VEGF, cytokines or growth factors other than VEGF may also contribute to endothelial cell survival and tumor angiogenesis and even be up-regulated when the effects of VEGF are inhibited (Fong et al., 1999; Buchdunger et al., 2000). We have observed in vitro that ON-III has a low inhibitory effect on endothelial cell growth if EGF is present in the medium. This suggests that several different factors support endothelial survival and tumor vascularization. Anti-VEGF therapy may be more effective against some types of tumors than others, and future therapy may necessitate a combination of antiangiogenic agents with different

mechanisms of action, as well as conventional therapies targeting the tumor cells. This may give VEGF inhibitors additional therapeutic applications over antiangiogenic agents with other mechanisms of action. Soluble VEGF receptors and antibodies against VEGF or its receptors have also been proposed as agents to block VEGF; compared with ON-III, however, the antibodies have the disadvantages of large proteins (Klement et al., 2000; Dias et al., 2001).

Taken together, we identified ON-III, a potent inhibitor of VEGF signaling with antitumor effect in vivo. This compound is from the dried flower *C. operculatus* used in traditional Chinese medicine for treatment of inflammation for many years. No adverse events have been observed clinically. Therefore, ON-III is a promising anticancer agent targeting VEGFR tyrosine kinase.

References

- Baek JH, Jang JE, Kang CM, Chung HY, Kim ND, and Kim KW (2000) Hypoxia-induced VEGF enhances tumor survivability via suppression of serum deprivation-induced apoptosis. *Oncogene* **19**:4621–4631.
- Binetruy-Tournaire R, Demangel C, Malavaud B, Vassy R, Rouyre S, Kraemer M, Plouet J, Derbin C, Perret G, and Mazie JC (2000) Identification of a peptide blocking vascular endothelial growth factor (VEGF)-mediated angiogenesis. *EMBO (Eur Mol Biol Organ) J* **19**:1525–1533.
- Blagosklonny MV (2004) Antiangiogenic therapy and tumor progression. *Cancer Cell* **5**:13–17.
- Buchdunger E, Cioffi CL, Law N, Stover D, Ohno-Jones S, Druker BJ, and Lydon NB (2000) Abl protein-tyrosine kinase inhibitor STI571 inhibits in vitro signal transduction mediated by c-kit and platelet-derived growth factor receptors. *J Pharmacol Exp Ther* **295**:139–145.
- Carlomagno F, Vitagliano D, Guida T, Ciardiello F, Tortora G, Vecchio G, Ryan AJ, Fontanini G, Fusco A, and Santoro M (2002) ZD6474, an orally available inhibitor of KDR tyrosine kinase activity, efficiently blocks oncogenic RET kinases. *Cancer Res* **62**:7284–7290.
- Colavitti R, Pani G, Bedogni B, Anzevino R, Borrello S, Waltenberger J, and Galeotti T (2002) Reactive oxygen species as downstream mediators of angiogenic signaling by vascular endothelial growth factor receptor-2/KDR. *J Biol Chem* **277**:3101–3108.
- Dias S, Hattori K, Heissig B, Zhu Z, Wu Y, Witte L, Hicklin DJ, Tateno M, Bohlen P, Moore MA, et al. (2001) Inhibition of both paracrine and autocrine VEGF/VEGFR-2 signaling pathways is essential to induce long-term remission of xenotransplanted human leukemias. *Proc Natl Acad Sci USA* **98**:10857–10862.
- Dias S, Shmelkov SV, Lam G, and Rafii S (2002) VEGF (165) promotes survival of leukemic cells by Hsp90-mediated induction of Bcl-2 expression and apoptosis inhibition. *Blood* **99**:2532–2540.
- Dreys J, Hofmann I, Hugenschmidt H, Wittig C, Madjar H, Muller M, Wood J, Martiny-Baron G, Unger C, and Marme D (2000) Effects of PTK787/ZK 222584, a specific inhibitor of vascular endothelial growth factor receptor tyrosine kinases, on primary tumor, metastasis, vessel density and blood flow in a murine renal cell carcinoma model. *Cancer Res* **60**:4819–4824.
- Eberhard A, Kahler T, Goede V, Hemmerlein B, Plate KH, and Augustin HG (2000) Heterogeneity of angiogenesis and blood vessel maturation in human tumors: implications for antiangiogenic tumor therapies. *Cancer Res* **60**:1388–1393.
- Foekens JA, Peters HA, Grebenchtchikov N, Look MP, Meijer-van Gelder ME, Geurts-Moespot A, van der Kwast TH, Sweep CG, and Klijn JG (2001) High tumor levels of vascular endothelial growth factor predict poor response to systemic therapy in advanced breast cancer. *Cancer Res* **61**:5407–5414.
- Folkman J (1990) What is the evidence that tumors are angiogenesis dependent? *J Natl Cancer Inst* **82**:4–6.
- Fong TA, Shawver LK, Sun L, Tang C, App H, Powell TJ, Kim YH, Schreck R, Wang X, Risau W, et al. (1999) SU5416 is a potent and selective inhibitor of the vascular endothelial growth factor receptor (Flk-1/KDR) that inhibits tyrosine kinase catalysis, tumor vascularization and growth of multiple tumor types. *Cancer Res* **59**:99–106.
- Geng L, Donnelly E, McMahon G, Lin PC, Sierra-Rivera E, Oshinka H, and Hallahan DE (2001) Inhibition of vascular endothelial growth factor receptor signaling leads to reversal of tumor resistance to radiotherapy. *Cancer Res* **61**:2413–2419.
- Hanahan D (1997) Signaling vascular morphogenesis and maintenance. *Science (Wash DC)* **277**:48–50.
- Huang X, Wong MK, Yi H, Watkins S, Laird AD, Wolf SF, and Gorelik E (2002) Combined therapy of local and metastatic 4T1 breast tumor in mice using SU6668, an inhibitor of angiogenic receptor tyrosine kinases and the immunostimulator B7.2-IgG fusion protein. *Cancer Res* **62**:5727–5735.
- Ichikura T, Tomimatsu S, Ohkura E, and Mochizuki H (2001) Prognostic significance of the expression of vascular endothelial growth factor (VEGF) and VEGF-C in gastric carcinoma. *J Surg Oncol* **78**:132–137.
- Itokawa T, Nohikawa H, Nishioka Y, Sone S, Iwamoto Y, Yamada Y, Cherrington J, McMahon G, Shibuya M, Kuwano M, et al. (2002) Antiangiogenic effect by SU5416 is partly attributable to inhibition of Flt-1 receptor signaling. *Mol Cancer Ther* **1**:295–302.
- Jeong SJ, Itokawa T, Shibuya M, Kuwano M, Ono M, Higuchi R, and Miyamoto T (2002) Costunolide, a sesquiterpene lactone from *Saussurea lappa*, inhibits the VEGFR KDR/Flk-1 signaling pathway. *Cancer Lett* **187**:129–133.

- Keyes KA, Mann L, Sherman M, Galbreath E, Schirtzinger L, Ballard D, Chen YF, Iversen P, and Teicher BA (2004) LY317615 decreases plasma VEGF levels in human tumor xenograft-bearing mice. *Cancer Chemother Pharmacol* **53**:133–140.
- Klement G, Baruchel S, Rak J, Man S, Clark K, Hicklin DJ, Bohlen P, and Kerbel RS (2000) Continuous low-dose therapy with vinblastine and VEGF receptor-2 antibody induces sustained tumor regression without overt toxicity. *J Clin Invest* **105**:R15–R24.
- Laird AD, Christensen JG, Li G, Carver J, Smith K, Xin X, Moss KG, Louie SG, Mendel DB, and Cherrington JM (2002) SU6668 inhibits Flk-1/KDR and PDGFR-beta in vivo, resulting in rapid apoptosis of tumor vasculature and tumor regression in mice. *FASEB J* **16**:681–690.
- Laird AD, Vajkoczy P, Shawver LK, Thurnher A, Liang C, Mohammadi M, Schlessinger J, Ullrich A, Hubbard SR, Blake RA, et al. (2000) SU6668 is a potent antiangiogenic and antitumor agent that induces regression of established tumors. *Cancer Res* **60**:4152–4160.
- Mendel DB, Laird AD, Xin X, Louie SG, Christensen JG, Li G, Schreck RE, Abrams TJ, Ngai TJ, Lee LB, et al. (2003) In vivo antitumor activity of SU11248, a novel tyrosine kinase inhibitor targeting vascular endothelial growth factor and platelet-derived growth factor receptors: determination of a pharmacokinetic/pharmacodynamic relationship. *Clin Cancer Res* **9**:327–337.
- Mendel DB, Schreck RE, West DC, Li G, Strawn LM, Tancigco SS, Vasile S, Shawver LK, and Cherrington JM (2000) The angiogenesis inhibitor SU5416 has long-lasting effects on vascular endothelial growth factor receptor phosphorylation and function. *Clin Cancer Res* **6**:4848–4858.
- Nakopoulou L, Stefanaki K, Panayotopoulou E, Giannopoulou I, Athanassiadou P, Gakiopoulou-Givalou H, and Louvrou A (2002) Expression of the vascular endothelial growth factor receptor-2/Flk-1 in breast carcinomas: correlation with proliferation. *Hum Pathol* **33**:863–870.
- O'Farrell AM, Abrams TJ, Yuen HA, Ngai TJ, Louie SG, Yee KW, Wong LM, Hong W, Lee LB, Town A, et al. (2003) SU11248 is a novel FLT3 tyrosine kinase inhibitor with potent activity in vitro and in vivo. *Blood* **101**:3597–3605.
- Qian CN, Min HQ, Lin HL, Feng GK, Ye YL, Wang LG, and Kuang ZJ (1999) Anti-tumor effect of angiogenesis inhibitor TNP-470 on the human nasopharyngeal carcinoma cell line NPC/HK1. *Oncology* **57**:36–41.
- Smaill JB, Palmer BD, Rewcastle GW, Denny WA, McNamara DJ, Dobrusin EM, Bridges AJ, Zhou H, Showalter HD, et al. (1999) Tyrosine kinase inhibitors. 15. 4-(Phenylamino)quinazoline and 4-(phenylamino)pyrido[d]pyrimidine acrylamides as irreversible inhibitors of the ATP binding site of the epidermal growth factor receptor. *J Med Chem* **42**:1803–1815.
- Smith BD, Smith GL, Carter D, Sasaki CT, and Haffty BG (2000) Prognostic significance of vascular endothelial growth factor protein levels in oral and oropharyngeal squamous cell carcinoma. *J Clin Oncol* **18**:2046–2052.
- Spiekermann K, Faber F, Voswinckel R, and Hiddemann W (2002) The protein tyrosine kinase inhibitor SU5614 inhibits VEGF-induced endothelial cell sprouting and induces growth arrest and apoptosis by inhibition of c-kit in AML cells. *Exp Hematol* **30**:767–773.
- Takahashi T, Ueno H, and Shibuya M (1999) VEGF activates protein kinase C-dependent, but Ras-independent Raf-MEK-MAP kinase pathway for DNA synthesis in primary endothelial cells. *Oncogene* **18**:2221–2230.
- Teicher BA, Menon K, Alvarez E, Galbreath E, Shih C, and Faul MM (2001) Antiangiogenic and antitumor effects of a protein kinase Cbeta inhibitor in human HT-29 colon carcinoma and human CaKi1 renal cell carcinoma xenografts. *Anticancer Res* **21**:3175–3184.
- Thakker GD, Hajjar DP, Muller WA, and Rosengart TK (1999) The role of phosphatidylinositol 3-kinase in vascular endothelial growth factor signaling. *J Biol Chem* **274**:10002–10007.
- UKCCCR (1998) United Kingdom Co-ordinating Committee on Cancer Research (UKCCCR) Guidelines for the Welfare of Animals in Experimental Neoplasia (Second Edition). *Br J Cancer* **77**:1–10.
- Wedge SR, Ogilvie DJ, Dukes M, Kendrew J, Chester R, Jackson JA, Boffey SJ, Valentine PJ, Curwen JO, Musgrove HL, et al. (2002) ZD6474 inhibits vascular endothelial growth factor signaling, angiogenesis and tumor growth following oral administration. *Cancer Res* **62**:4645–4655.
- Zhang F, Liu M, and Lu R (1990) Studies on the chemical constituents from the bud of *Cleistocalyx operculatus*. *Acta Botanica Sinica* **32**:469–472.
- Zhang XS, Zhu XF, Gao JS, Ye YL, Feng QS, Liu ZC, and Zeng YX (2002) Variable sensitivity of endothelial cells to epirubicin in xenografts of human nasopharyngeal carcinoma CNE-2 cells. *Cancer Biol Ther* **1**:263–265.
- Zhu XF, Liu ZC, Xie BF, Feng GK, and Zeng YX (2003) Ceramide induces cell cycle arrest and upregulates p27kip in nasopharyngeal carcinoma cells. *Cancer Lett* **193**:149–154.
- Zhu XF, Liu ZC, Xie BF, Li ZM, Feng GK, Yang D, and Zeng YX (2001) EGFR tyrosine kinase inhibitor AG1478 inhibits cell proliferation and arrests cell cycle in nasopharyngeal carcinoma cells. *Cancer Lett* **169**:27–32.
- Zhu XF, Zhang XS, Li ZM, Yao YQ, Xie BF, Liu ZC, and Zeng YX (2000) Apoptosis induced by ceramide in hepatocellular carcinoma Bel7402 cells. *Acta Pharmacol Sin* **21**:225–228.

Address correspondence to: Dr. Yi-Xin Zeng, Cancer Center, Sun Yat-sen University, 651 DongFeng Road East, GuangZhou 510060, China. E-mail: yxzeng@gzsums.edu.cn