

MINIREVIEW

Insect GABA Receptors: Splicing, Editing, and Targeting by Antiparasitics and Insecticides

Steven David Buckingham, Philip Charles Biggin, Benedict Mark Sattelle, Laurence Andrew Brown, and David Barry Sattelle

Medical Research Council Functional Genetics Unit, Department of Human Anatomy and Genetics (S.D.B., B.M.S., L.A.B., D.B.S.) and Department of Biochemistry (P.C.B.), University of Oxford, Oxford, United Kingdom

Received June 2, 2005; accepted July 18, 2005

ABSTRACT

Ionotropic GABA receptors are abundant in both vertebrate and invertebrate nervous systems, where they mediate rapid, mostly inhibitory synaptic transmission. A GABA-gated chloride channel subunit from *Drosophila melanogaster* [Resistant to Dieldrin (RDL)] has been cloned, functionally expressed, and found to exhibit many aspects of the pharmacology of native, bicuculline-insensitive insect GABA receptors. RDL is the target of the commercially important insecticide fipronil. A point mutation in the channel-lining region of the RDL molecule is known to underlie most cases of resistance to insecticides acting on GABA receptors. RDL is widely distributed throughout the in-

sect nervous system, but the subunit composition of RDL-containing in native receptors is unknown. It is possible that in some instances, RDL coexpresses with glutamate-gated chloride channel subunits. Other ionotropic receptor subunits (LCCH3 and GRD) form GABA-gated cation channels when heterologously expressed. Interest in RDL as a model ligand-gated anion channel has been enhanced by the recent discovery of pre-mRNA A-to-I editing, which, together with alternative splicing, adds to the functional diversity of this GABA receptor subunit.

Ionotropic GABA receptor molecules (GABARs) are members of the dicysteine-loop ('Cys-loop') superfamily of neurotransmitter receptors, which also includes nicotinic acetylcholine receptors, type 3 5-hydroxytryptamine receptors, and glycine receptors (Karlin and Akabas, 1995; Karlin, 2002; Olsen et al., 2004). Ionotropic GABARs are pentameric proteins. Each polypeptide subunit possesses a long N-terminal extracellular domain containing residues that contribute to the neurotransmitter binding site and four transmembrane regions (M1-M4), the second of which (M2) provides many of the residues that line the integral chloride channel (Whiting, 2003; Kim et al., 2004; Darlison et al., 2005). Vertebrate ionotropic GABARs may be divided into two pharmacological categories: bicuculline-sensitive GABA_A receptors (composed of α_{1-6} , β_{1-3} , γ_{1-3} , δ , ϵ , θ , and π_{1-3} subunits and allosterically

modulated by benzodiazepines and barbiturates as well as pregnane steroids) and bicuculline-insensitive GABA_C receptors (composed of the three known isoforms of the ρ subunit and insensitive to the majority of modulators of GABA_A receptors) (Mustafa, 1995; Sieghart, 1995; McKernan and Whiting, 1996; Whiting, 2003; Connolly and Wafford, 2004; Rudolph and Mohler, 2004). This division into A and C subtypes is difficult to reconcile with recent findings on GABARs in brainstem neurons, which are composed of ρ_1 subunits coexpressed with $\alpha 1$ and $\gamma 2$ subunits (Milligan et al., 2004) to yield receptors with properties of both GABA_A and GABA_C subtypes. Furthermore, GABA_A and GABA_B receptors, which differ structurally and in their signaling mechanisms, may be closely functionally coupled. This is supported by the recent finding that the GABA_A $\gamma 2S$ subunit forms a complex with GABA_BR1 subunits (thereby enhancing GABA_B receptor trafficking to the cell surface, which otherwise requires coexpression with the GABA_BR2 subunit). The GABA_A $\gamma 2S$

Article, publication date, and citation information can be found at <http://molpharm.aspetjournals.org>.
doi:10.1124/mol.105.015313.

ABBREVIATIONS: GABAR, GABA receptor; RDL, resistant to dieldrin; GRD, GABA receptor of *Drosophila*; TM, transmembrane; CACA, *cis*-aminocrotonic acid; BIDN, 3,3-bis-trifluoromethyl-bicyclo-[2,2,1]heptane-2,2-dicarbonitrile; Ro5-4864, 4'-chlorodiazepam; LCCH3, Ligand-gated chloride channel homolog 3; nAChR, nicotinic acetylcholine receptor.

subunit also forms a complex with both GABA_BR1 and GABA_BR2 subunits to allow agonist-induced receptor internalization (Balasubramanian et al., 2004).

Insect ionotropic GABARs do not readily fit the vertebrate GABA_A/GABA_C receptor categories. The majority are distinguished from the GABA_A type of vertebrate receptors by their insensitivity to bicuculline and differ from GABA_C receptors in that they are subject to allosteric modulation, albeit weak, by benzodiazepines and barbiturates (Sattelle, 1990). An insect ionotropic *Drosophila melanogaster* GABAR subunit, RDL (Resistant to Dieldrin), can be heterologously expressed to form functional homo-oligomeric receptors, the pharmacology of which closely resembles that of the majority of native insect GABARs and so has proved to be of value in investigating GABAR physiology and pharmacology. Here, we show that RDL very closely mimics the pharmacology of most insect native neuronal GABARs reported so far and is therefore a useful homomeric model insect GABAR. As such, it facilitates the interpretation of site-directed mutagenesis experiments.

The Insect GABA-Gated Chloride Channel Family

The GABA-receptor subunit-encoding *Rdl* gene was isolated from a naturally occurring dieldrin-resistant strain of the dipteran *D. melanogaster* (Ffrench-Constant et al., 1991, 1993; Ffrench-Constant and Rocheleau, 1993). Highly conserved *Rdl*-like receptor genes have since been identified in several other insect orders. RDL subunits are distributed throughout the adult (Harrison et al., 1996) and embryonic (Aronstein and Ffrench-Constant, 1995) nervous system of *D. melanogaster* but not on muscle (Harrison et al., 1996). Their expression pattern suggests roles in fast GABAergic synaptic transmission (Buchner et al., 1988; DiAntonio et al., 1993; Harrison et al., 1996) and learning (Harrison et al., 1996; Strambi et al., 1998; Sattelle et al., 2000) as well as visual and olfactory processing (Harrison et al., 1996). Recordings of spontaneous GABA-mediated currents in *D. melanogaster* embryonic neurons have provided electrophysiological evidence that RDL contributes to synaptic GABARs (Lee et al., 2003).

Other subunits in the *D. melanogaster* genome are predicted to have anion channel characteristics based on signature sequences in their predicted transcripts, among which there are GABAR subunit candidates. These include CG6927, CG7589, CG11340, and CG12344. Until functional expression data are available, these candidates are difficult to distinguish from other insect ligand-gated anion channels, which include histamine-gated chloride channels (Gengs et al., 2002), a glutamate-gated chloride channel α subunit (Cully et al., 1996), candidate GABA-glycine-gated ion channels GRD (Harvey et al., 1994) and CG7589 (identified in genome annotations), as well as CG8916, a protein identified in genome annotations as a GABA_A receptor but so far not functionally expressed. Furthermore, although the GABAR-like sequences of LCCH3, another GABAR-like subunit, and GRD suggested possible anion channel activity, coexpression of these two subunits in *Xenopus laevis* oocytes results in a cationic channel (Gisselmann et al., 2004). This finding counsels caution when interpreting annotations based on sequence data alone. RDL and LCCH3 coexpress to form a

picrotoxinin-insensitive, bicuculline-sensitive channel with pharmacology quite unlike native insect GABARs (Zhang et al., 1995). So far, coexpression of RDL with GRD has not been reported. It is noteworthy that RDL is not expressed on cockroach muscle (Harrison et al., 1996) despite the existence of GABA-gated chloride currents in these cells (Rauh et al., 1997a; Schnee et al., 1997). This suggests the existence of muscle GABA-gated chloride channels that do not contain RDL.

Alternative Splicing of the *Rdl* Gene

By means of alternative splicing in two of its nine exons (exons 3 and 6), the RDL gene encodes four distinct polypeptides (Fig. 1A), all of which are expressed (Ffrench-Constant and Rocheleau, 1993). Three splice variants (RDLac, RDLbd, and RDLab) have been cloned and functionally expressed (Hosie and Sattelle, 1996a). Residues that vary in exon 3 lie upstream to loop D of the GABA binding site and near the equivalent positions of known determinants of agonist potency in GABA_A receptors of vertebrates. Alternative splicing occurs in other ligand-gated ion channels (Villarroel, 1999) and has been shown to affect agonist sensitivity in glycine receptors (Miller et al., 2004). It is noteworthy that alternative splicing of the *D. melanogaster* nicotinic acetylcholine receptor subunit $\alpha 6$ (Grauso et al., 2002) also occurs in exon 3, which aligns to exon 3 of RDL. The exon 6 alternatively spliced residues of RDL lie in loops F and C (Fig. 1B) and exon 6 splicing affects the expressed receptor's affinity for GABA (Hosie and Sattelle, 1996a) (Fig. 1C). The agonist profile for one splice variant (RDLbc) remains to be determined. Nevertheless, two of the variant amino acid residues are particularly interesting: the two splice variants containing the 'a' form of exon 3 (RDLac and RDLad) have Val53 and Leu57, respectively, whereas those containing the 'b' form (RDLbc and RDLbd) have Leu53 and Lys57. These two residues are near a valine (Val46 in the nicotinic α subunit of modified muscle cells in the electric organ of *Torpedo marmorata*), which is highly conserved among ligand-gated ion channels and in *T. marmorata* is proposed to form the closest contact between the ligand-binding and transmembrane domains (Miyazawa et al., 2003). This suggests that RDL exon 3 is involved in the coupling of ligand binding to pore opening. Structural perturbation due to alternative splicing in exon 6 has been shown to account for differences in the potency of three glycine receptor agonists (Miller et al., 2004).

The 10 alternatively-spliced residues affected by exon 6 splicing are located in loops F and C of the RDL binding site on the external face of the vestibule (Fig. 1B). Loop F is known to be involved in GABA binding and chloride channel gating in the vertebrate GABA_A $\alpha 1$ subunit (Newell and Czajkowski, 2003). Our comparative models of the RDL receptor suggest that exon 6 alternative splicing may influence the putative agonist binding site at the RDL subunit interface (S. D. Buckingham, P. C. Biggin, B. M. Sattelle, L. A. Brown, and D. B. Sattelle, unpublished observations).

The pharmacological diversity among vertebrate GABA_A receptor subtypes is largely determined by different subunit isoforms that make up the receptor. Because the *D. melanogaster* genome contains only a small number of genes iden-

tified as GABAR subunits, two of which can be coexpressed to form GABA-gated cation channels, the alternative splicing of the *Rdl* gene in *D. melanogaster* may serve to increase functional diversity in the absence of a large number of GABAR subunits. This possibility is strengthened by findings from immunocytochemical studies suggesting that RDL is very widely distributed and hence a likely component of many insect nervous system GABARs (Aronstein et al., 1996; Harrison et al., 1996).

Pre-mRNA Editing of RDL

Pre-mRNA A-to-I editing is widespread in animal tissues (Athanasiadis et al., 2004). It involves substitution of an inosine for an adenosine, the result being interpreted by the translation machinery as a guanosine. RDL is edited at four sites that may, based on comparison with equivalent residues in other receptors, affect the receptor's responses to GABA. They are R122G in the N-terminal domain,




Fig. 1. Structure and post-transcriptional modifications of RDL. **A**, exon structure of the *Rdl* gene; gray boxes represent transmembrane (TM) regions 1 to 3 coded by exon 7 and the TM 4 region encoded by exon 9. Alternative splicing occurs in exons 3 and 6. The two versions of exon 3 (a and b) differ by two amino acid residues. The two versions of exon 6 (c and d) differ by 10 amino acid residues. [Adapted from Hosie AM, Aronstein K, Sattelle DB, and French-Constant RH (1997) Molecular biology of insect neuronal GABA receptors. *Trends Neurosci* 20: 578–583. Copyright © 1997 Elsevier B.V. Used with permission.] **B**, model of the extracellular domain (ECD) of RDLac, based on the acetylcholine binding protein (Protein Data Bank code 1I9B). Red and yellow spheres denote alternatively spliced residues of exons 3 and 6, respectively, and orange spheres show the site of RNA-editing on the N terminus (R122G). **C**, alternative splicing of RDL exon 6 produces variations in the agonist dose-response curves for GABA and its analogs isoguvacine (ISOG), isonipecotic acid (ISON), and 3-aminopropanesulfonic acid (3-APS). All four agonists were more potent on RDL_{ac} homomeric receptors compared with homomeric RDL_{ad} receptors (points represent mean \pm S.E.M. from at least three oocytes). [Reprinted from Hosie AM, Buckingham SD, Presnail JK, and Sattelle DB (2001) Alternative splicing of a Drosophila GABA receptor subunit gene identifies determinants of agonist potency, *Neuroscience* 102:709–714. Copyright © 2001 Elsevier B.V. Used with permission.]

I283V in the first transmembrane domain, N294D in the TM1–TM2 loop, and M360V in the TM3–TM4 loop (Fig. 2A). So far, the effects of these substitutions on receptor function have not been tested, but comparison with equivalent residues in other subunits suggests that at least some of these edited residues are at positions likely to affect agonist actions. The residue Arg122 is in loop A of the agonist binding site, adjacent to the equivalent residue (Trp93) in nicotinic acetylcholine receptors known to be critical in agonist binding (Fig. 2B). The edited residue Met360 is located in the large intracellular loop between TM3 and TM4, which has been shown to influence GABAR desensitization kinetics and GABA EC_{50} (Fisher, 2004) and aligns with Leu277 of the human $\alpha 2$ GABA_A receptor subunit, which, when mutated to alanine, results in a 50-fold increase in the EC_{50} for GABA (O'Shea and Harrison, 2000).

RDL As a Model of Native Insect GABA Receptors

In addition to its potential as a model for anionic ligand-gated ion channels, RDL may also provide a convenient model for insect GABARs, which are the targets of commercially important insecticides, such as fipronil, used in animal health and crop protection applications. Although little is known of the pharmacology or physiology of native *D. melanogaster* GABA receptors, RDL shares a number of distinctive features with native GABA receptors of other insect species.

The Agonist Profile and Bicuculline Insensitivity of RDL Resembles the Most Abundant Native Insect GABARs. RDL homomeric receptors are distinguished from vertebrate GABA_A receptors by their insensitivity to bicuculline, their low sensitivity to 3-aminopropanesulfonic acid, and their high sensitivity to both isoguvacine (Hosie and Sattelle, 1996a) and muscimol (Buckingham et al., 1994a) (Fig. 1C). They are distinguished from vertebrate GABA_C receptors by the high efficacy of muscimol, *cis*-aminocrotonic acid (CACA), and isoguvacine. Such pharmacology is typical of many native insect GABARs (Lummis and Sattelle, 1985, 1986; Sattelle et al., 1988). Unlike many native insect GABARs (Brotz and Borst, 1996), however, RDL is sensitive to CACA, an agonist of vertebrate GABA_C, but not of GABA_A receptors (Hosie and Sattelle, 1996a). However, it should be noted that CACA does act on some insect GABARs, including those that mediate inhibition by identified filiform hair receptors of an identified projection interneuron (Gauglitz and Pfluger, 2001).

A feature of most insect GABARs is an insensitivity to bicuculline, a feature shared by extrasynaptic GABARs on the motor neuron D_f of *Periplaneta americana* (Sattelle et al., 1988; Buckingham et al., 1994b), synaptic GABARs on the giant interneuron 2 of *P. americana* (Buckingham et al., 1994b), GABARs that mediate inhibition by identified filiform hair receptors of an identified projection interneuron (Gauglitz and Pfluger, 2001), motion-sensitive visual interneurons of the blowfly *Calliphora erythrocephala* (Brotz and Borst, 1996), GABA-mediated inhibitory postsynaptic potentials recorded in an identified locust (*Schistocerca gregaria*)




Fig. 2. Pre-mRNA A-I editing of RDL. A, schematic representation of the RDL subunit, showing how potential A-I editing of mRNA for RDL results in changes to the amino acid sequence at residues 122, 283, 294, and 360. B, comparative model illustrating residues changed by alternative splicing and RNA-editing. Images on the left are side views; those on the right show pentamers viewed from above. The model of the TM domain of RDLac (B, left) is based on coordinates of *Torpedo* nAChR (Protein Data Bank code 1OED). Orange and red spheres denote sites of RNA editing in the TM1 region and the TM1–TM2 loop, respectively. Yellow spheres denote site of A→S/G mutation conferring resistance to insecticides.

interneuron (Watson and Burrows, 1987) and many, but not all, locust (*Locusta migratoria*) neuron ionotropic GABARs (Lees et al., 1987; Benson, 1988). However, it is clear that, in addition to bicuculline-insensitive receptors, a class of bicuculline-sensitive GABARs also exists in insects. For example, bicuculline blocks ionotropic GABARs of some adult (Wal-drop et al., 1987) and larval *Manduca sexta* abdominal ganglion neurons (Sattelle et al., 2003) as well as electrically

evoked or odor-evoked inhibitory postsynaptic potentials and GABA responses in projection neurons of antennal lobes of this species (Christensen et al., 1998a,b). It is clear, however, that RDL and its equivalent cloned from *Heliothis virescens* (Wolff and Wingate, 1998) resemble, in their insensitivity to bicuculline, the majority of insect ionotropic GABARs.

RDL Provides a Useful Model of the Native Convulsant Antagonist Site. The convulsant antagonist site is the




Fig. 3. Physiological characteristics of RDL receptors. A, chemical structures for convulsant compounds BIDN, dieldrin, and fipronil. B, actions of selected convulsant antagonists on the currents evoked by 10 μM GABA at wild-type RDL receptors expressed in *X. laevis* oocytes. Traces for GABA alone, followed by GABA and 10 μM picrotoxinin (PTX), fipronil, *t*-butylbicyclophosphorothionate (TBPS), and 4-*n*-propyl-4'-ethynylbicycloorthobenzoate (EBOB), respectively. [Adapted from Buckingham SD, Hosie AM, Roush RL, and Sattelle DB (1994) Actions of agonists and convulsant antagonists on a *Drosophila melanogaster* GABA receptor (Rdl) homo-oligomer expressed in *Xenopus* oocytes. *Neurosci Lett* 181:137–140. Copyright © 1994 Elsevier B.V. Used with permission.] C, effects of the convulsant BIDN on GABA-evoked currents at the RDL receptor. Top, compared with control, preincubation of oocytes with 100 μM BIDN shifts the dose-response curve for GABA to the right and reduces the maximal currents evoked, suggesting that the antagonist actions of BIDN are neither strictly competitive nor noncompetitive. Bottom, dose inhibition curves using 100 μM GABA show that compared with wild-type RDL (□), the A302S mutation (○) decreases the inhibitory effects of the BIDN at the RDL receptor (points represent mean ± S.E.M. from at least three oocytes). [Adapted from Hosie AM, Shirai Y, Buckingham SD, Rauh JJ, Roush RT, Baylis HA, and Sattelle DB (1995) Blocking actions of BIDN, a bicyclic dinitrile convulsant compound, on wild-type and dieldrin-resistant GABA receptor homo-oligomers of *Drosophila melanogaster* expressed in *Xenopus* oocytes. *Brain Res* 693:257–260. Copyright © 1995 Elsevier B.V. Used with permission.]

target of insecticides, including fipronil (Fig. 3), although fipronil, at micromolar concentrations, has also been shown to be a potent blocker of insect (Ikeda et al., 2003; Zhao et al., 2004) and nematode (Horoszok et al., 2001) L-glutamate-gated chloride channels. Pharmacological differences between insect and vertebrate GABARs at this site have been shown to underlie the selectivity of these compounds for insects. RDL shares much of the convulsant pharmacology that distinguishes insect from vertebrate GABARs (Buckingham et al., 1996). The pharmacology of the convulsant site of RDL has been delineated in detail in a structure/function analysis of the sensitivity of RDL homomers to a range of picrodendrin analogs (Hosie et al., 1996). As is the case for in situ insect GABARs, RDL is also blocked by 3,3-bis-trifluoromethyl-bicyclo-[2,2,1]heptane-2,2-dicarbonitrile (BIDN) (Hosie et al., 1995; Buckingham et al., 1996) (Fig. 4), an important new probe of the insect convulsant site (Rauh et al., 1997a,b). Although both BIDN and fipronil (Fig. 3) are potent blockers of RDL (Buckingham et al., 1994a, 1996; Rauh et al., 1997a; Grolleau and Sattelle, 2000) (Figs. 3 and 4) and in situ insect GABARs (Rauh et al., 1997a), single channel studies have revealed that they act at distinct albeit possibly overlapping sites (Grolleau and Sattelle, 2000) (Fig. 3, C and D). Indeed, as many as four convulsant binding sites have been identified in housefly head membranes (Deng et al., 1993).

RDL Mimics the Barbiturate and Steroid Insensitivity of Native Insect GABA Receptors. GABARs of insects are only weakly sensitive to pregnane steroids and to pentobarbital, which potentially enhance vertebrate GABA_A receptors (Lummis and Sattelle, 1986). Phenobarbital and pentobarbital (10 μ M–1 mM) enhanced the GABA responses of RDL homomers (Buckingham et al., 1996). The pregnane steroid 5 α -pregnan-3 α -ol-20-one (10 μ M) yielded a moderate enhancement of the GABA-response of RDL homomers (Miller et al., 1994), reflecting the low steroid sensitivity of native insect GABARs. The potency of pentobarbital and phenobarbital on RDL homomers (Hosie and Sattelle, 1996b) is similar to that observed on native locust GABARs (Lees et al., 1987)

and less than that seen on vertebrate GABA_A receptors. The steroid site on vertebrate GABARs remains elusive; once detected, however, it will be interesting to compare it with the equivalent regions of RDL, which is much less sensitive to these compounds.

Benzodiazepines Are Less Potent on RDL Homomers than on Native Insect GABA Receptors. Although RDL mimics well the agonist, competitive antagonist, and convulsant sites of insect GABARs, differences have emerged among RDL and known native insect GABARs in their sensitivity to benzodiazepines. For example, whereas the agonist responses of certain bicuculline-insensitive insect GABARs are enhanced by micromolar concentrations of flunitrazepam (Sattelle et al., 1988), the GABA responses of RDL homomers (Miller et al., 1994) or its *H. virescens* equivalent (Wolff and Wingate, 1998) are unaffected by flunitrazepam (100 μ M). More importantly, insect native GABARs and RDL homomers are both enhanced by lower concentrations of 4'-chlorodiazepam (Ro5-4864) (it inhibits native receptors at higher concentrations; R. Higashino and D. B. Sattelle, unpublished observations), although RDL receptors were approximately 100-fold less sensitive to this compound than native insect receptors (Sattelle et al., 1988; Buckingham et al., 1996; Hosie and Sattelle, 1996b).

Thus, RDL homomers do not match closely the benzodiazepine or barbiturate pharmacology of known insect GABARs. It is therefore probable that native GABARs of insects are composed of RDL coexpressed with other as-yet-unidentified subunits.

Coassembly of RDL with Glutamate-Gated Chloride Receptor Subunits?

One possible explanation for the difference between RDL homomers and native receptors at the benzodiazepine site is that native insect GABARs are not RDL homomers but include other subunit types or are a mix of RDL splice variants. Indeed, there is evidence that the GABARs present at the




Fig. 4. Analysis of the actions of BIDN and fipronil using single channel recordings from S2 cells expressing Rdl. A, examples of currents recorded from an outside-out patch from an S2-RDL cell exposed to 50 μ M GABA (c, closed; o, open). B, current amplitude histograms for GABA and GABA with the convulsant antagonist BIDN show that the presence of BIDN leads to a reduction in the size of channel conductance. C, the convulsant antagonist fipronil reduces the probability of channel openings (P_o) in the S2-RDL cells, with addition of BIDN further reducing the P_o . D, plotting the increasing concentration of BIDN, or 300 nM BIDN with increasing concentrations of fipronil, against the reduction in P_o reveals that the effects of these two antagonists do not seem to be via the same site on the RDL receptor. [Adapted from Grolleau F and Sattelle DB (2000) Single channel analysis of the blocking actions of BIDN and fipronil on a *Drosophila melanogaster* GABA receptor (RDL) stably expressed in a *Drosophila* cell line. *Br J Pharmacol* 130:1833–1842. Copyright © 2000 British Pharmacological Society. Used with permission.]

neuromuscular junction of *Caenorhabditis elegans* are heteromultimers composed of two splice variants encoded by the *unc-49* gene (Bamber et al., 2005). In heterologously expressed vertebrate GABA_A receptors, the efficacy of benzodiazepines is highly dependent upon receptor subtype composition (Buhr and Sigel, 1997; Smith, 2001). Indeed, the single channel properties of RDL-containing receptors on cultured *D. melanogaster* neurons differ from those of RDL homomers (Zhang et al., 1995). Thus, native RDL-containing receptors are likely to be hetero-oligomers of RDL and other, as-yet-unknown subunits. Heterologous expression studies provide evidence that coexpression of RDL with other insect putative GABAR subunits results in receptors with distinct pharmacologies (Table 1). In a recent study (Ludmerer et al., 2002), antibodies to RDL and to a *D. melanogaster* glutamate-gated chloride channel subunit both immunoprecipitated the entire fraction of receptors for the insecticide (nodulosporic acid), suggesting that RDL coassembles in vivo with a glutamate receptor subunit—the first evidence for coassembly of subunits from different subclasses of ligand-gated ion channels. Because RDL staining is not observed in muscle, even though GABA-gated chloride channels are present in insect muscle, such coexpression must be confined to certain regions of the nervous system.

GABA-Gated Cation Channels

It was recently reported (Gisselmann et al., 2004) that GRD and LCCH3 form GABA-gated cation channels when coexpressed in *X. laevis* oocytes. These channels are bicuculline-insensitive but are blocked by picrotoxin. Unlike RDL, they are insensitive to dieldrin and diazepam. It would be interesting to pursue further the expression patterns of RDL and these two subunits—RDL expression is not known to overlap with that of LCCH3 (Aronstein et al., 1996), but the expression patterns of GRD have not yet been reported, so the prediction that the expression patterns of LCCH3 and GRD overlap remains untested. A GABA-gated cation channel (EXP-1) has also been cloned from *C. elegans* (Beg and Jorgensen, 2003). No native GABA-gated cation channel has yet been recorded from any insect neuron, so if such receptors exist in insects, they may be rare.

Studies on other homomeric receptors from the ligand-gated ion channel superfamily offer some insight into the residues that define the charge-selectivity filter of the RDL receptor. In the case of both the $\alpha 7$ nicotinic acetylcholine receptor (Corringer et al., 1999a,b) and the 5HT₃ receptor (Gunthorpe and Lummis, 2001), selected residues in the TM1-TM2 loop and within TM2 dictate whether the receptors are cationic or anionic. An alignment of RDL with LCCH3, GRD, and the $\alpha 7$ nAChR suggests that GRD at least shares

similarities with the cationic $\alpha 7$ receptor at these key residues (Fig. 5A).

Molecular Target for Insecticides and a Point Mutation in RDL That Accounts for Resistance

As GABARs are prevalent in the nervous systems of insects, they are the targets of naturally occurring (e.g., picrotoxinin and a wide range of picrotoxin-like molecules of plant origin) as well as man-made (e.g., dieldrin, fipronil) insecticides. In the past, dieldrin, the structure of which can be related to picrotoxin, was a major insecticide, but it has been banned because it persists in the environment and because its intensive use in the years after World War II resulted in many examples of resistance to insecticide. Fipronil is the first of the phenylpyrazole group of chemicals to be introduced for pest control and is now a major pesticide in use with crops and as an antiparasitic (flea and mite control), with an estimated world market of US\$150 million. The usefulness of an insecticide is often limited by the development of resistance, which is common in field populations of many insect species (Georghiu, 1986). In the case of dieldrin resistance in *D. melanogaster*, insecticide resistance arises from the substitution of a single amino acid: Ala→Ser at the 2' position in the lumen of the GABA receptor's channel (Hosie et al., 1997; ffrench-Constant et al., 2000; Bass et al., 2004) (Fig. 5B). In the peach aphid, *Myzus persicae*, cyclo-diene resistance results from an Ala→Gly mutation (Anthony et al., 1998). Fipronil resistance in the RDL equivalent in *Drosophila simulans* has also been shown to arise from the two mutations, Ala→Gly similar to the equivalent mutation in *D. melanogaster*, and a T350M mutation in the third transmembrane domain (Le Goff et al., 2005). The mutation at Thr350 has not been described in any field populations of dieldrin-resistant *D. melanogaster* or in any mosquito populations. Resistance raised in the laboratory (Le Goff et al., 2005) may arise from a selection pressure different from that in the field, resulting in different mutations. In functional expression studies comparing the wild-type and resistant forms of *D. melanogaster* RDL, 2' Ala→Ser confers resistance to a variety of insecticides, such as fipronil, dieldrin, picrotoxinin, and picrodendrin-O, which act allosterically as noncompetitive antagonists of insect GABARs. This substitution also renders RDL homomers resistant to the antagonists *t*-butylbicyclophosphorothionate, lindane, and BIDN (Sattelle, 1990; Buckingham et al., 1994a, 1996; Hosie et al., 1995) (Fig. 2D), although these compounds interact noncompetitively in radioligand binding studies, suggesting that they have distinct binding sites. The ability of RDL to form functional homomers allows a more convenient study of the

TABLE 1

Heterologous expression of RDL alone or with other putative insect GABAR subunits results in receptors with distinct pharmacologies

Subunit combination	Heterologous Expression	Pharmacology	Reference
RDL homomer	Oocytes, S2 cells (anion channel)	Picrotoxinin-sensitive; Bicuculline-insensitive	Hosie and Sattelle, 1996a
RDL + LCCH3	Sf-9	Picrotoxinin-insensitive; Bicuculline-sensitive	Zhang et al., 1995
RDL + GRD	Not reported	Not reported	N/A
GRD + LCCH3	Oocytes (cation channel)	Picrotoxinin-sensitive; Bicuculline-insensitive	Gisselmann et al., 2004




Fig. 5. Alignments of RDL. **A**, alignments of the *D. melanogaster* RDL sequence (from above) with other *D. melanogaster* GABA-gated channels and the human α7 nicotinic acetylcholine receptor subunit (a cation channel) showing variations in residues of TM2 reported to define ion-selectivity of the channel. Sequences: *D. melanogaster* LCCH3 (GenBank accession number AAL28208), GRD (GenBank accession number CAA55144) and human α7 nAChR (GenBank accession number NP_000737). Alignments were created using ClustalX and GeneDoc programs. **B**, alignments of the TM1 and TM2 regions of the RDL sequence in *D. melanogaster* and other insect species reveals conserved sequence of amino acid residues. Sequences: *D. melanogaster* (fruit fly; GenBank accession number NP_729462), *Musca domestica* (housefly; GenBank accession number BAD16658), *Anopheles gambiae* (malaria mosquito; GenBank accession number XP_316072), *Apis mellifera* (honeybee; GenBank accession number AAC63381), and *Heliothis virescens* (tobacco budworm; GenBank accession number AAB62575). Region of alignment contains the alanine residue, where mutation to a serine or glycine has been reported to confer resistance to a number of compounds, including insecticides.

effects of the A302S/G mutation upon single channel properties of the channel, as well as simplifying the interpretations of such experiments. For example, the single channel properties of dieldrin-resistant RDL stably expressed in a *D. melanogaster* cell line provided the first evidence that the insecticide fipronil and the new convulsant site probe BIDN act at separate sites (Grolleau and Sattelle, 2000) (Summarized in Fig. 4).

Conclusions

The cloned RDL receptor is not only a convenient model of insect ionotropic GABARs but also a convenient homomeric anionic ligand-gated ion channel accessible to combined computational and site-directed mutagenesis approaches to understanding the structural basis of anion channel function and insecticide resistance. The experimental work so far indicates that RDL is widely distributed throughout the central nervous systems of insects of different orders (*Lepidoptera*, *Diptera*, *Coleoptera* and *Dictyoptera*), is located at synaptic sites and on neuronal cell bodies, and that RDL homomers mimic the characteristic agonist, barbiturate, and pregnane steroid pharmacology of native bicuculline-insensitive insect GABARs. In all these respects, RDL very closely resembles native insect GABARs. The close similarity between RDL homomers and native insect GABARs suggests that RDL-like subunits, which are widely distributed in the *D. melanogaster* nervous system, may underlie several aspects of the distinct pharmacology of bicuculline-insensitive insect GABARs.

The RDL subunit offers potential for generating insect-vertebrate chimeric GABARs, which will help identify the binding sites of modulators that differ in their actions on vertebrate and insect GABARs. Its ability to form homomeric

receptors that resemble native receptors also facilitates computational studies, just as the vertebrate α7 receptor enabled molecular modeling of cationic ligand-gated ion channels. Expression studies on the fourth as-yet-uncharacterised splice variant will complete the description of all splice variants for this receptor. Finally, expression studies of edited RDL receptors will provide the first functional characterization of the consequences of pre-mRNA A-to-I editing in GABARs.

Acknowledgments

We are indebted to Drs. Alistair Hosie and Andrew Jones for helpful discussions.

References

- Anthony N, Unruh T, Ganser D and Ffrench-Constant R (1998) Duplication of the Rdl GABA receptor subunit gene in an insecticide-resistant aphid, *Myzus persicae*. *Mol Genet* **260**:165–175.
- Aronstein K, Auld V, and Ffrench-Constant R (1996) Distribution of two GABA receptor-like subunits in the *Drosophila* CNS. *Invert Neurosci* **2**:115–120.
- Aronstein K and Ffrench-Constant R (1995) Immunocytochemistry of a novel GABA receptor subunit Rdl in *Drosophila melanogaster*. *Invert Neurosci* **1**:25–31.
- Athanasiadis A, Rich A, and Maas S (2004) Widespread A-to-I RNA editing of Alu-containing mRNAs in the human transcriptome. *PLoS Biol* **2**:e391.
- Balasubramanian S, Teissere JA, Raju DV, and Hall RA (2004) Heterooligomerization between GABAA and GABAB receptors regulates GABAB receptor trafficking. *J Biol Chem* **279**:18840–18850.
- Bamber BA, Richmond JE, Otto JF, and Jorgensen EM (2005) The composition of the GABA receptor at the *Caenorhabditis elegans* neuromuscular junction. *Br J Pharmacol* **144**:502–509.
- Bass C, Schroeder I, Turberg A, Field LM, and Williamson MS (2004) Identification of the Rdl mutation in laboratory and field strains of the cat flea, *Ctenocephalides felis* (Siphonaptera: Pulicidae). *Pest Manag Sci* **60**:1157–1162.
- Beg AA and Jorgensen EM (2003) EXP-1 is an excitatory GABA-gated cation channel. *Nat Neurosci* **6**:1145–1152.
- Benson JA (1988) Bicuculline blocks the response to acetylcholine and nicotine but not to muscarine or GABA in isolated insect neuronal somata. *Brain Res* **458**:65–71.
- Brotz TM and Borst A (1996) Cholinergic and GABAergic receptors on fly tangential cells and their role in visual motion detection. *J Neurophysiol* **76**:1786–1799.
- Buchner E, Bader R, Buchner S, Cox J, Emson PC, Flory E, Heizmann CW, Hemm S, Hofbauer A, and Oertel WH (1988) Cell-specific immuno-probes for the brain of

- normal and mutant *Drosophila melanogaster*. I. Wildtype visual system. *Cell Tissue Res* **253**:357–370.
- Buckingham SD, Hosie AM, Roush RL, and Sattelle DB (1994a) Actions of agonists and convulsant antagonists on a *Drosophila melanogaster* GABA receptor (Rdl) homo-oligomer expressed in *Xenopus* oocytes. *Neurosci Lett* **181**:137–140.
- Buckingham SD, Hue B, and Sattelle DB (1994b) Actions of bicuculline on cell body and neuropilar membranes of identified insect neurones. *J Exp Biol* **186**:235–244.
- Buckingham SD, Matsuda K, Hosie AM, Baylis HA, Squire MD, Lansdell SJ, Millar NS, and Sattelle B (1996) Wild-type and insecticide-resistant homo-oligomeric GABA receptors of *Drosophila melanogaster* stably expressed in a *Drosophila* cell line. *Neuropharmacology* **35**:1393–1401.
- Buhr A and Sigel E (1997) A point mutation in the gamma2 subunit of gamma-aminobutyric acid type A receptors results in altered benzodiazepine binding site specificity. *Proc Natl Acad Sci USA* **94**:8824–8829.
- Christensen TA, Waldrop BR, and Hildebrand JG (1998a) GABAergic mechanisms that shape the temporal response to odors in moth olfactory projection neurons. *Ann NY Acad Sci* **855**:475–481.
- Christensen TA, Waldrop BR, and Hildebrand JG (1998b) Multitasking in the olfactory system: context-dependent responses to odors reveal dual GABA-regulated coding mechanisms in single olfactory projection neurons. *J Neurosci* **18**:5999–6008.
- Connolly CN and Wafford KA (2004) The Cys-loop superfamily of ligand-gated ion channels: the impact of receptor structure on function. *Biochem Soc Trans* **32**:529–534.
- Corringer PJ, Bertrand S, Galzi JL, Devillers-Thiery A, Changeux JP, and Bertrand D (1999a) Molecular basis of the charge selectivity of nicotinic acetylcholine receptor and related ligand-gated ion channels. *Novartis Found Symp* **225**:215–224; discussion 224–30.
- Corringer PJ, Bertrand S, Galzi JL, Devillers-Thiery A, Changeux JP, and Bertrand D (1999b) Mutational analysis of the charge selectivity filter of the alpha7 nicotinic acetylcholine receptor. *Neuron* **22**:831–843.
- Cully DF, Pareiss PS, Liu KK, Schaeffer JM, and Arena JP (1996) Identification of a *Drosophila melanogaster* glutamate-gated chloride channel sensitive to the anti-parasitic agent avermectin. *J Biol Chem* **271**:20187–20191.
- Darlison MG, Pahal I, and Thode C (2005) Consequences of the evolution of the GABA_A receptor gene family. *Cell Mol Neurobiol* **25**:607–624.
- Deng Y, Palmer CJ, and Casida JE (1993) House fly head GABA-gated chloride channel: four putative insecticide binding sites differentiated by [³H]EBOB and [³⁵S]TBPS. *Pestic Biochem Physiol* **47**:98–112.
- DiAntonio A, Burgess RW, Chin AC, Deitcher DL, Scheller RH, and Schwarz TL (1993) Identification and characterization of *Drosophila* genes for synaptic vesicle proteins. *J Neurosci* **13**:4924–4935.
- ffrench-Constant RH, Anthony N, Aronstein K, Rocheleau T, and Stilwell G (2000) Cyclodiene insecticide resistance: from molecular to population genetics. *Annu Rev Entomol* **45**:449–466.
- ffrench-Constant RH, Mortlock DP, Shaffer CD, MacIntyre RJ, and Roush RT (1991) Molecular cloning and transformation of cyclodiene resistance in *Drosophila*: an invertebrate gamma-aminobutyric acid subtype A receptor locus. *Proc Natl Acad Sci USA* **88**:7209–7213.
- ffrench-Constant RH and Rocheleau TA (1993) *Drosophila* gamma-aminobutyric acid receptor gene Rdl shows extensive alternative splicing. *J Neurochem* **60**:2323–2326.
- ffrench-Constant RH, Steichen JC, Rocheleau TA, Aronstein K, and Roush RT (1993) A single-amino acid substitution in a gamma-aminobutyric acid subtype A receptor locus is associated with cyclodiene insecticide resistance in *Drosophila* populations. *Proc Natl Acad Sci USA* **90**:1957–1961.
- Fisher JL (2004) The alpha1 and alpha6 subunit subtypes of the mammalian GABA(A) receptor confer distinct channel gating kinetics. *J Physiol* **561**:433–448.
- Gauglitz S and Pflüger HJ (2001) Cholinergic transmission via central synapses in the locust nervous system. *J Comp Physiol A* **187**:825–836.
- Gengs C, Leung HT, Skingsley DR, Iovchev MI, Yin Z, Semenov EP, Burg MG, Hardie RC, and Pak WL (2002) The target of *Drosophila* photoreceptor synaptic transmission is a histamine-gated chloride channel encoded by ort (hclA). *J Biol Chem* **277**:42113–42120.
- Georghiou GP (1986) The magnitude of the resistance problem, in *Pesticide Resistance: Strategies and Tactics for Management*, p. 14–43, National Academy Press, Washington, DC.
- Gisselmann G, Plonka J, Pusch H, and Hatt H (2004) *Drosophila melanogaster* GRD and LCCH3 subunits form heteromultimeric GABA-gated cation channels. *Br J Pharmacol* **142**:409–413.
- Grauso M, Reenan RA, Culetto E, and Sattelle DB (2002) Novel putative nicotinic acetylcholine receptor subunit genes, Dalpha5, Dalpha6 and Dalpha7, in *Drosophila melanogaster* identify a new and highly conserved target of adenosine deaminase acting on RNA-mediated A-to-I pre-mRNA editing. *Genetics* **160**:1519–1533.
- Grolleau F and Sattelle DB (2000) Single channel analysis of the blocking actions of BIDN and fipronil on a *Drosophila melanogaster* GABA receptor (RDL) stably expressed in a *Drosophila* cell line. *Br J Pharmacol* **130**:1833–1842.
- Gunthorpe MJ and Lummis SC (2001) Conversion of the ion selectivity of the 5-HT_{3A} receptor from cationic to anionic reveals a conserved feature of the ligand-gated ion channel superfamily. *J Biol Chem* **276**:10977–10983.
- Harrison JB, Chen HH, Sattelle E, Barker PJ, Huskisson NS, Rauh JJ, Bai D, and Sattelle DB (1996) Immunocytochemical mapping of a C-terminus anti-peptide antibody to the GABA receptor subunit, RDL in the nervous system in *Drosophila melanogaster*. *Cell Tissue Res* **284**:269–278.
- Harvey RJ, Schmitt B, Hermans-Borgmeyer I, Gundelfinger ED, Betz H, and Darlison MG (1994) Sequence of a *Drosophila* ligand-gated ion-channel polypeptide with an unusual amino-terminal extracellular domain. *J Neurochem* **62**:2480–2483.
- Horoszok L, Raymond V, Sattelle DB, and Wolstenholme AJ (2001) GLC-3: a novel fipronil and BIDN-sensitive, but picrotoxinin-insensitive, L-glutamate-gated chloride channel subunit from *Caenorhabditis elegans*. *Br J Pharmacol* **132**:1247–1254.
- Hosie AM, Aronstein K, Sattelle DB, and ffrench-Constant RH (1997) Molecular biology of insect neuronal GABA receptors. *Trends Neurosci* **20**:578–583.
- Hosie AM, Ozoe Y, Koike K, Ohmoto T, Nikaido T, and Sattelle DB (1996) Actions of picrodendrin antagonists on dieldrin-sensitive and -resistant *Drosophila* GABA receptors. *Br J Pharmacol* **119**:1569–1576.
- Hosie AM and Sattelle DB (1996a) Agonist pharmacology of two *Drosophila* GABA receptor splice variants. *Br J Pharmacol* **119**:1577–1585.
- Hosie AM and Sattelle DB (1996b) Allosteric modulation of an expressed homo-oligomeric GABA-gated chloride channel of *Drosophila melanogaster*. *Br J Pharmacol* **117**:1229–1237.
- Hosie AM, Shirai Y, Buckingham SD, Rauh JJ, Roush RT, Baylis HA, and Sattelle DB (1995) Blocking actions of BIDN, a bicyclic dinitrile convulsant compound, on wild-type and dieldrin-resistant GABA receptor homo-oligomers of *Drosophila melanogaster* expressed in *Xenopus* oocytes. *Brain Res* **693**:257–260.
- Ikedo T, Zhao X, Kono Y, Yeh JZ, and Narahashi T (2003) Fipronil modulation of glutamate-induced chloride currents in cockroach thoracic ganglion neurons. *Neurotoxicology* **24**:807–815.
- Karlin A (2002) Emerging structure of the nicotinic acetylcholine receptors. *Nat Rev Neurosci* **3**:102–114.
- Karlin A and Akabas MH (1995) Toward a structural basis for the function of nicotinic acetylcholine receptors and their cousins. *Neuron* **15**:1231–1244.
- Kim S, Chamberlain AK, and Bowie JU (2004) A model of the closed form of the nicotinic acetylcholine receptor m2 channel pore. *Biophys J* **87**:792–799.
- Lee D, Su H, and O'Dowd DK (2003) GABA receptors containing Rdl subunits mediate fast inhibitory synaptic transmission in *Drosophila* neurons. *J Neurosci* **23**:4625–4634.
- Lees G, Beadle DJ, Neumann R, and Benson JA (1987) Responses to GABA by isolated insect neuronal somata: pharmacology and modulation by a benzodiazepine and a barbiturate. *Brain Res* **401**:267–278.
- Le Goff G, Hamon A, Berge JB, and Amichot M (2005) Resistance to fipronil in *Drosophila* simulans: influence of two point mutations in the RDL GABA receptor subunit. *J Neurochem* **92**:1295–1305.
- Ludmerer SW, Warren VA, Williams BS, Zheng Y, Hunt DC, Ayer MB, Wallace MA, Chaudhary AG, Egan MA, Meinke PT, et al. (2002) Ivermectin and nodulisporic acid receptors in *Drosophila melanogaster* contain both gamma-aminobutyric acid-gated Rdl and glutamate-gated GluCl alpha chloride channel subunits. *Biochemistry* **41**:6548–6560.
- Lummis SCR and Sattelle DB (1985) Insect CNS GABA_A receptors. *Neurosci Lett* **60**:13–18.
- Lummis SCR and Sattelle DB (1986) Binding sites for [³H]GABA, [³H]flunitrazepam and [³⁵S]TBPS in insect CNS. *Neurochem Int* **9**:287–293.
- McKernan RM and Whiting PJ (1996) Which GABA_A-receptor subtypes really occur in the brain? *Trends Neurosci* **19**:139–143.
- Millar NS, Buckingham SD, and Sattelle DB (1994) Stable expression of a functional homo-oligomeric *Drosophila* GABA receptor in a *Drosophila* cell line. *Proc R Soc Lond B Biol Sci* **258**:307–314.
- Miller PS, Harvey RJ, and Smart TG (2004) Differential agonist sensitivity of glycine receptor alpha2 subunit splice variants. *Br J Pharmacol* **143**:19–26.
- Milligan CJ, Buckley NJ, Garret M, Deuchars J, and Deuchars SA (2004) Evidence for inhibition mediated by coassembly of GABA_A and GABAC receptor subunits in native central neurons. *J Neurosci* **24**:7241–7250.
- Miyazawa A, Fujiyoshi Y, and Unwin N (2003) Structure and gating mechanism of the acetylcholine receptor pore. *Nature (Lond)* **423**:949–955.
- Mustafa BA (1995) Diversity of GABA receptors in the vertebrate outer retina. *Trends Neurosci* **18**:118–120.
- Newell JG and Czajkowski C (2003) The GABA_A receptor alpha 1 subunit Pro174-Asp191 segment is involved in GABA binding and channel gating. *J Biol Chem* **278**:13166–13172.
- Olsen RW, Chang CS, Li G, Hancher HJ, and Wallner M (2004) Fishing for allosteric sites on GABA(A) receptors. *Biochem Pharmacol* **68**:1675–1684.
- O'Shea SM and Harrison NL (2000) Arg-274 and Leu-277 of the gamma-aminobutyric acid type A receptor alpha2 subunit define agonist efficacy and potency. *J Biol Chem* **275**:22764–22768.
- Rauh JJ, Benner E, Schnee ME, Cordova D, Holyoke CW, Howard MH, Bai D, Buckingham SD, Hutton ML, Hamon A, et al. (1997a) Effects of [³H]BIDN, a novel bicyclic dinitrile radioligand for GABA-gated chloride channels of insects and vertebrates. *Br J Pharmacol* **121**:1496–1505.
- Rauh JJ, Holyoke CW, Kleier DA, Presnail JK, Benner EA, Cordova D, Howard MH, Hosie AM, Buckingham SD, et al. (1997b) Polycyclic dinitriles: a novel class of potent GABAergic insecticides provides a new radioligand, [³H]BIDN. *Invert Neurosci* **3**:261–268.
- Rudolph U and Mohler H (2004) Analysis of GABA_A receptor function and dissection of the pharmacology of benzodiazepines and general anesthetics through mouse genetics. *Annu Rev Pharmacol Toxicol* **44**:475–498.
- Sattelle D (1990) GABA receptors of insects. *Adv Insect Physiol* **22**:1–113.
- Sattelle DB, Bai D, Chen HH, Skeer JM, Buckingham SD, and Rauh JJ (2003) Bicuculline-insensitive GABA-gated Cl⁻ channels in the larval nervous system of the moth *Manduca sexta*. *Invert Neurosci* **5**:37–43.
- Sattelle DB, Harrison JB, Chen HH, Bai D, and Takeda M (2000) Immunocytochemical localization of putative gamma-aminobutyric acid receptor subunits in the head ganglia of *Periplaneta americana* using an anti-RDL C-terminal antibody. *Neurosci Lett* **289**:197–200.
- Sattelle DB, Pinnock RD, Wafford KA, and David JA (1988) GABA receptors on the cell-body membrane of an identified insect motor neuron. *Proc R Soc Lond B Biol Sci* **232**:443–456.
- Schnee M, Rauh J, Buckingham SD, and Sattelle DB (1997) Pharmacology of skeletal muscle GABA-gated chloride channels in the cockroach *Periplaneta americana*. *J Exp Biol* **200**:2947–2955.

- Sieghart W (1995) Structure and pharmacology of gamma-aminobutyric acid A receptor subtypes. *Pharmacol Rev* **47**:181–234.
- Smith TA (2001) Type A gamma-aminobutyric acid (GABAA) receptor subunits and benzodiazepine binding: significance to clinical syndromes and their treatment. *Br J Biomed Sci* **58**:111–121.
- Strambi C, Cayre M, Sattelle DB, Augier R, Charpin P, and Strambi A (1998) Immunocytochemical mapping of an RDL-like GABA receptor subunit and of GABA in brain structures related to learning and memory in the cricket *Acheta domestica*. *Learn Mem* **5**:78–89.
- Villarroel A (1999) Alternative splicing generates multiple mRNA forms of the acetylcholine receptor gamma-subunit in rat muscle. *FEBS Lett* **443**:381–384.
- Waldrop B, Christensen TA, and Hildebrand JG (1987) GABA-mediated synaptic inhibition of projection neurons in the antennal lobes of the sphinx moth, *Manduca sexta*. *J Comp Physiol A* **161**:23–32.
- Watson AH and Burrows M (1987) Immunocytochemical and pharmacological evidence for GABAergic spiking local interneurons in the locust. *J Neurosci* **7**:1741–1751.
- Whiting PJ (2003) The GABAA receptor gene family: new opportunities for drug development. *Curr Opin Drug Discov Dev* **6**:648–657.
- Wolff MA and Wingate VP (1998) Characterization and comparative pharmacological studies of a functional gamma-aminobutyric acid (GABA) receptor cloned from the tobacco budworm, *Heliothis virescens* (Noctuidae:Lepidoptera). *Invert Neurosci* **3**:305–315.
- Zhang HG, Lee HJ, Rocheleau T, French-Constant RH, and Jackson MB (1995) Subunit composition determines picrotoxin and bicuculline sensitivity of *Drosophila* γ -aminobutyric acid receptors. *Mol Pharmacol* **48**:835–840.
- Zhao X, Yeh JZ, Salgado VL, and Narahashi T (2004) Fipronil is a potent open channel blocker of glutamate-activated chloride channels in cockroach neurons. *J Pharmacol Exp Ther* **310**:192–201.

Address correspondence to: David B. Sattelle, MRC Functional Genetics Unit, Department of Human Anatomy and Genetics, University of Oxford, Oxford, UK. E-mail: david.sattelle@anat.ox.ac.uk
