

Substance P, Neurokinins A and B, and Synthetic Tachykinin Peptides Protect Mesencephalic Dopaminergic Neurons in Culture via an Activity-Dependent Mechanism

Bénédicte Salthun-Lassalle, Sabine Traver, Etienne C. Hirsch, and Patrick P. Michel

Institut National de la Santé et de la Recherche Médicale Unité 679, Experimental Neurology and Therapeutics, Hôpital de la Salpêtrière, Paris, France

Received June 3, 2005; accepted August 2, 2005

ABSTRACT

We evaluated the neuroprotective potential of tachykinin peptides using a model system in which mesencephalic dopaminergic (DA) neurons die spontaneously and selectively as they mature. The three native tachykinins, substance P (SP), neurokinin (NK) A, and NKB afforded substantial protection against DA cell demise. The selective NK₁ receptor antagonist [D-Pro⁹,[spiro- γ -lactam] Leu¹⁰,Trp¹¹]substance P (GR71251) was sufficient in itself to suppress the effect of SP, whereas a cotreatment with GR71251 and the NK₃ receptor antagonist (R)-N-[α -(methoxycarbonyl)benzyl]-2-phenylquinoline-4-carboxamide (SB218795) was required to prevent the effects of both NKA and NKB. Consistent with these results, D-Ala-[L-Pro⁹,Me-Leu⁸]substance P(7–11) (GR73632), a selective agonist of NK₁ receptors and [pro⁷]-NKB, a selective agonist of NK₃ receptors, conferred protection to DA neurons, whereas

(Lys³, Gly⁸-R- γ -lactam-Leu⁹)neurokinin A(3–10) (GR64349), which activates specifically NK₂ receptors, did not. DA neurons rescued by tachykinins accumulated [³H]DA efficiently, which suggests that they were also totally functional. Neuroprotection by tachykinins was highly selective for DA neurons, rapidly reversed upon treatment withdrawal, and reproduced by but independent of glial cell line-derived neurotrophic factor. Survival promotion by tachykinins was abolished by blocking voltage-gated Na⁺ channels with tetrodotoxin or N-type voltage-gated Ca²⁺ channels with ω -conotoxin-MVIIA, which indicates that an increase in neuronal excitability was crucially involved in this effect. Together, these data further support the notion that the survival of mesencephalic DA neurons during development depends largely on excitatory inputs, which may be provided in part by tachykinins.

Tachykinins belong to a family of neuropeptides that share a common C-terminal sequence, Phe-Xaa-Gly-Leu-Met-NH₂, that is crucial for interaction with their receptors (Severini et al., 2002; Pennefather et al., 2004). The mammalian tachykinins include substance P (SP), neurokinin (NK) A, and NKB, the effects of which are mediated by the receptors NK₁, NK₂, and NK₃. SP is the most potent tachykinin at the NK₁ receptor site, whereas NKA exhibits the highest affinity for the NK₂ receptor and NKB for the NK₃ receptor (Pennefather et al., 2004). The effects through NK receptors are generally

coupled to a G protein (G_q/G₁₁)-regulated phosphoinositide pathway (Khawaja and Rogers, 1996), but G protein-independent coupling has also been described previously (Yang et al., 2003).

NK receptors are implicated in various biological effects that include smooth muscle contraction, inflammatory processes, hypotensive effects, and stimulation of gland secretion (Severini et al., 2002). In the nervous system, tachykinins also operate as neuromodulators/neurotransmitters via mechanisms that are generally excitatory (Stacey et al., 2002). In addition, there is evidence that tachykinins also have intrinsic neuroprotective properties. They were found to reverse β -amyloid-induced toxicity (Kowall et al., 1991), to protect against excitotoxic cell death (Calvo et al., 1996), and to limit neurodegeneration caused by trophic factor deprivation (Lallemend et al., 2003).

This work was supported by Institut National de la Santé et de la Recherche Médicale. B.S.-L. was supported by a Doctoral Research Fellowship from Ministère de l'Éducation Nationale, de la Recherche et de la Technologie, and P.P.M. was supported by Pierre Fabre Research Center (Castres, France).

Article, publication date, and citation information can be found at <http://molpharm.aspetjournals.org>.
doi:10.1124/mol.105.015453.

ABBREVIATIONS: araC, cytosine arabinoside; Boc-D-FMK, *N*-tert-butoxycarbonyl-D-fluoromethylketone; DA, dopamine or dopaminergic; DIV, day in vitro; GDNF, glial cell line-derived neurotrophic factor; 5-HT, serotonin; MAP2, microtubule-associated protein-2; NK, neurokinin; PBS, phosphate-buffered saline; SP, substance P; TH, tyrosine hydroxylase; TTX, tetrodotoxin; RT-PCR, reverse-transcriptase polymerase chain reaction; AB, antibody; GR71251, [D-Pro⁹,[spiro- γ -lactam]Leu¹⁰,Trp¹¹]substance P; SB218795, (R)-N-[α -(methoxycarbonyl)benzyl]-2-phenylquinoline-4-carboxamide; GR73632, D-Ala-[L-Pro⁹,Me-Leu⁸]substance P(7–11); GR64349, (Lys³,Gly⁸-R- γ -lactam-Leu⁹)neurokinin A(3–10); R396, acetyl-Leu-Asp-Gln-Trp-Phe-GlyNH₂; GBR-12909, 1-(2(bis(4-fluorophenyl)methoxy)ethyl)-4-(3-phenylpropyl)piperazine di-HCl.

The loss of nigrostriatal dopaminergic (DA) neurons in Parkinson's disease and related disorders leads to profound motor impairment (Agid, 1991). The identification of signals or factors that control the survival and function of DA neurons is therefore of interest because it might not only provide a better understanding of the pathophysiological mechanisms of these diseases but also foster the development of new therapeutic strategies to halt their progression (Dawson and Dawson, 2002). There is indirect evidence that tachykinins may influence the survival of mesencephalic DA neurons: 1) in the brain, the substantia nigra is the richest area in SP and SP-containing axon terminals (Hokfelt et al., 1991); 2) striatal DA depletion in Parkinson's disease and in animal models of the disease results in profound changes in SP expression (Mauborgne et al., 1983; Betarbet and Greenamyre, 2004); and 3) excitatory stimuli, which are believed to play a key role in the survival of DA neurons during development and possibly in the adult brain (Douhou et al., 2001; Katsuki et al., 2001; Salthun-Lassalle et al., 2004), can be evoked by direct application of NK₁ or NK₃ agonists to mesencephalic DA neurons (Nalivaiko et al., 1997).

To examine the potential of tachykinins to promote the survival of DA neurons, we have used a model system in which these neurons die spontaneously and progressively as a function of time (Michel and Agid, 1996). More specifically, we wished to evaluate the neuroprotective potential of native and synthetic tachykinins for DA neurons; determine what receptor subtypes are involved; establish whether these mechanisms are shared with or related to that of GDNF, a prototypical neuroprotective factor for DA neurons (Choi-Lundberg et al., 1997); and explore the mechanisms underlying neuroprotection.

Materials and Methods

Peptide Agonists/Antagonists of Tachykinin Receptors and Other Pharmacological Agents. Native tachykinins SP, NKA, NKB, and synthetic peptide agonists/antagonists (GR73632, GR64349, [pro⁷]-NKB, GR71251, and R396) of tachykinin receptors were from NeoMPS (Strasbourg, France). The nonpeptide receptor antagonist SB218795 was from Bioblock Scientific (Strasbourg, France). Glial cell line-derived neurotrophic factor (GDNF) was purchased from AbCys (Paris, France), and the anti-GDNF neutralizing antibody (AB 212 NA) was from R&D Systems Europe (Lille, France). The pancaspase inhibitor *N*-tert-butoxycarbonyl-D-fluoromethylketone (Boc-D-FMK) was obtained from Calbiochem (Darmstadt, Germany). Others pharmacological reagents were from Sigma/RBI (Natick, MA). Tritiated neurotransmitters and tritiated methyl-thymidine were from GE Healthcare (Little Chalfont, Buckinghamshire, UK).

Mesencephalic Cell Cultures. Animals were treated in accordance with the *Guide for the Care and Use of Laboratory Animals* (National Research Council, 1996), the European Directive number 86/609, and the guidelines of the local institutional animal care and use committee. Cultures were prepared from the ventral mesencephalon of embryonic day 15.5 Wistar rat embryos (Janvier Breeding Center, Le Genest St Isle, France) as described previously (Michel and Agid, 1996). After dissection, pieces of mesencephalic tissue were dissociated mechanically with no enzymatic treatment using a Pipetman (Gilson, Villier Le Bel, France) set to 900 μ l and plated onto polyethylenimine (1 mg/ml; Sigma-Aldrich, St. Quentin Fallavier, France)-coated 24-well culture plates. Cells plated at a density of 1.5 to 2.0 $\times 10^5$ /cm² were then maintained at 37°C in a humidified incubator with a 5% CO₂ atmosphere using 500 μ l of N5 culture medium (Salthun-Lassalle et al., 2004) supplemented with 5 mM

glucose, 5% horse serum, and 0.5% fetal bovine serum, except for the first 3 days, when 2.5% fetal bovine serum was used to enhance cell attachment. The cultures were fed daily by replacing 350 μ l of the culture medium. They were then supplemented with appropriate pharmacological treatments, including native tachykinins and agonists/antagonists of tachykinin receptors.

Quantification of Neuronal Survival. The survival of DA neurons was quantified by counting the number of cells labeled with an antibody against tyrosine hydroxylase (TH), as described previously. In brief, the cultures were fixed with 4% formaldehyde in phosphate-buffered saline (PBS) for 15 min. Cells were washed three times with PBS and then incubated for 24 h at 4°C with a monoclonal anti-TH antibody (Diasorin, Stillwater, MN) diluted 1:5000 in PBS containing 0.2% Triton X-100. The TH antibody was then revealed with an anti-mouse IgG Cy3 conjugate (1:500; Sigma/RBI) for 2 h at room temperature. Mesencephalic cultures contained between 1 and 2% TH⁺ cells at the time of plating. All neurons, regardless of their neurotransmitter phenotype, were identified by labeling microtubule-associated protein-2 (MAP2) with a monoclonal antibody (AP-20, Sigma/RBI) diluted 1:100 in PBS revealed with a anti-mouse Alexa Fluor 488 conjugate.

Measurement of Neurotransmitter Uptake. The functional integrity of DA neurons was evaluated by their ability to take up DA by active transport (Douhou et al., 2001). Uptake was initiated by the addition of 50 nM [³H]DA (40 Ci/mmol) to cultures preincubated for 10 min in 500 μ l of PBS containing 5 mM glucose and 100 μ M ascorbic acid. It was terminated after 15 min by two rapid washes with cold PBS. Cells were scraped off the culture wells, and the accumulated tritium was counted by liquid scintillation spectrometry. The uptake of [³H]DA was also visualized by microautoradiography (Troade et al., 2002). In this case, the incubation time was prolonged to 30 min, and [³H]DA was used at 100 nM to improve detection. The cultures were then fixed with a mixture of glutaraldehyde/formaldehyde (0.5:4%) and dehydrated with ethanol. Incorporation of [³H]DA was detected with the Hypercoat LM-1 emulsion (GE Healthcare) after an exposure of 10 days in the dark at 4°C. In both paradigms, blank values were obtained in the presence of 0.5 μ M GBR-12909 (Sigma-Aldrich, St. Louis, MO). GABA and serotonin (5-HT) uptake was measured as described using 50 nM [³H]GABA and 20 nM [³H]5-HT, respectively (Salthun-Lassalle et al., 2004). Note that the concentrations of the three neurotransmitters were lower than the *K_m* values of their transporters to be within the linear portion of the uptake time course.

Quantification and Identification of Proliferating Cells. [methyl-³H]Thymidine, a marker of DNA synthesis, was used to label and quantify proliferating cells as described previously (Mourlevat et al., 2003). Mesencephalic cultures maintained up to 7 DIV in the presence of test treatments were exposed to 1 μ Ci of [methyl-³H]thymidine (GE Healthcare; 40 Ci/mmol) for 3 h at 37°C in serum-free N5 medium supplemented with 5 mM glucose. After three rapid washes, the cells were allowed to recover for 1 h in the same culture medium to remove unincorporated radioactivity. The cultures were fixed in 4% formaldehyde for 15 min and when necessary were processed for TH immunofluorescence detection subsequently. They were then dehydrated in ethanol and exposed to Hypercoat LM-1 emulsion (GE Healthcare) for 4 days at 4°C to detect the tritiated label.

Quantification of Intracellular Calcium Levels. Cytoplasmic-free calcium levels were measured in individual neurons using Calcium Green-1-AM (Molecular Probes, Eugene, OR) as described previously (Douhou et al., 2001; Salthun-Lassalle et al., 2004).

Reverse Transcription and Polymerase Chain Reaction. Total RNA from mesencephalic cultures or from rat uterus tissue was isolated with RNABle solution (Laboratories Eurobio, Les Ulis, France). Three types of cultures were used: mixed cultures containing neurons and glial cells, pure neuronal cultures in which glial cells were eliminated by treatment with 3 μ M araC, and pure astrocyte cultures. First-strand cDNA was synthesized from 2 μ g of total

RNA using the RT kit (QIAGEN S.A., Courtaboeuf, France). To analyze the expression of NK₁, NK₂, and NK₃ receptors, transcripts were amplified by polymerase chain reaction under the following conditions: 35 cycles of 94°C for 30 s, 58°C for 30 s, and 72°C for 30 s, with 3 μ l of the reverse transcription mixture. The primers used for tachykinin receptors were: NK₁: forward, (5'-CCTCCTGCCCTACA-TCAACCC-3'); and reverse, (5'-CTGTGTCTGGAGGTATCGGG-3'); NK₂: forward, (5'-CATCACTGTGGACGAGGGGG-3'); and reverse, (5'-TGTCTTCCTCAGTTGGTGTC-3'); and NK₃: forward, (5'-CATT-CTCACTGCGATCTACC-3'); and reverse (5'-CTTCTTGCGGCTGG-ATTTGG-3') (Pinto et al., 1999). The polymerase chain reaction products were visualized by electrophoresis in a 2% agarose gel containing 0.1 mg/ml ethidium bromide.

Statistical Analysis. Simple comparisons between two groups were performed with Student's *t* test. Multiple comparisons against a single reference group were made by one-way analysis of variance followed by Dunnett's test. When all pairwise comparisons were made, the Student-Newman-Keuls test was used. S.E.M. values were derived from at least three independent experiments.

Results

Native Tachykinins SP, NKA, and NKB Increase the Number of DA Neurons in Mesencephalic Cultures.

The native tachykinins SP, NKA, and NKB increased the number of TH⁺ DA neurons in mesencephalic cultures (Fig. 1, A–C and E). The effects of SP and NKB on the number of TH⁺ cells were restricted to a narrow range of concentrations between 1 and 50 nM and 1 and 5 nM, respectively, with peak

effects at 10 and 1 nM (Fig. 1, A and C). At higher concentrations, no effect of SP or NKB on the number of DA neurons was observed. NKA treatment increased the number of DA cells as well, but at much higher concentrations (100–1000 nM; Fig. 1, B and E).

Synthetic Peptide Agonists of NK₁ and NK₃ but Not NK₂ Receptors Mimic the Effects of Native Tachykinins. SP, NKA, and NKB are believed to act preferentially through NK₁, NK₂, and NK₃ receptors, respectively (Pennefather et al., 2004). To determine whether selective agonists of these receptors could reproduce the effects of native tachykinins on DA neurons, we treated mesencephalic cultures with the NK₁ agonist GR73632, the NK₂ agonist GR64349, and the NK₃ agonist [pro7]-NKB. Consistent with the results obtained with SP and NKB, the number of DA neurons increased in the presence of GR73632 and [pro7]-NKB (Fig. 1D), in a narrow range of concentrations. The peak effect was estimated at 10 nM for GR73632 and at 1 nM for [pro7]-NKB (Fig. 1D). We were surprised to find, however, that the NK₂ receptor agonist GR64349 did not affect the number of TH⁺ cells, indicating that the action of NKA was probably mediated by its nonpreferential receptors.

Selective Antagonists of NK₁ and NK₃ Receptors Prevent the Effects of Tachykinins on DA Neurons. We wished first to address the specificity of the interaction between synthetic tachykinin agonists and NK receptors. Therefore, specific antagonists of NK₁, NK₂, and NK₃ recep-





Fig. 1. Native tachykinins and selective agonists of NK₁ and NK₃ receptors increase the number of TH⁺ cells in mesencephalic cultures. Number of TH⁺ neurons as a function of the concentrations of SP (A), NKA (B), and NKB (C). D, number of TH⁺ neurons in cultures exposed continually to synthetic agonists of NK₁ (GR73632), NK₂ (GR64349), and NK₃ ([pro7]-NKB) receptors. All of the test compounds, except for GR64349, increased TH⁺ cell number. *, *p* < 0.05, different from corresponding control cultures. E, illustration of the effects of SP, NKA, and NKB on the number of TH⁺ cells. Cell counts and illustrations were made using cultures at 10 DIV. Scale bar, 50 μ m.



Lassalle et al., 2004). A long-term treatment with GR73632 (10 nM) or [pro7]-NKB (1 nM) also reduced TH⁺ cell loss substantially, indicating that tachykinins were probably acting by preventing the apoptotic mechanism as well. However, to exclude the possibility that NK receptor activation increased the number of TH⁺ neurons by stimulating cell division, we treated the cultures with the NK₁ receptor agonist GR73632 in the presence of [methyl-³H]thymidine. Nuclei positive for the tritiated label were never found in association with TH⁺ neurons, indicating that NK₁ receptor stimulation did not induce the proliferation of DA neuroblasts or their precursor cells (Fig. 3B). Note that similar results were obtained with [pro7]-NKB (data not shown).

Tachykinin-Dependent Survival of DA Neurons Is a Function of the Duration of Treatment. To determine whether shorter treatments also prevented the death of TH⁺ neurons, the addition of the NK₃ receptor agonist [pro7]-NKB (1 nM) was delayed after plating. Under these conditions, the number of TH⁺ neurons that could be rescued by the treatment diminished progressively (Fig. 4A). Cultures treated continuously for 10 days (0–10 DIV) with 1 nM [pro7]-NKB had ~120% more TH⁺ neurons than untreated cultures, whereas cultures only treated from 4 to 10 DIV had only ~40% more. On the other hand, the protective effect of [pro7]-NKB was rapidly reversible if the treatment was stopped prematurely, and the cultures maintained in control medium up to 10 DIV (Fig. 4B). If [pro7]-NKB was withdrawn at day 8, only 40% of the TH⁺ were rescued 2 days later; it was more remarkable that, if withdrawn at 6 DIV, [pro7]-NKB produced no net increase in the number of TH⁺ neurons at 10 DIV (Fig. 4B). Similar results were obtained with the NK₁ agonist GR73632 (data not shown).




Fig. 3. The activation of NK₁ or NK₃ receptors provides true neuroprotection for mesencephalic DA neurons. **A**, number of TH⁺ neurons as a function of the age of the cultures after long-term treatment with GR73632 (10 nM), [pro7]-NKB (1 nM), or the broad spectrum and cell-permeable caspase inhibitor Boc-D-FMK (60 μ M). \star , $p < 0.05$, different from corresponding age-matched control cultures. **B**, illustration showing that TH⁺ cells do not incorporate [methyl-³H]thymidine after a treatment with GR73632. Similar results were obtained in the presence of [pro7]-NKB. Scale bar, 50 μ m.

TH⁺ Neurons Rescued by Activation of NK₁ and NK₃ Receptors Possess a Functional DA Transporter.

To assess the function of the rescued DA neurons, we quantified the uptake of [³H]DA and combined microautoradiographic detection of the tritiated neurotransmitter with immunofluorescent labeling of TH. Treatments with either the NK₁ agonist GR73632 or the NK₃ agonist [pro7]-NKB increased the number of TH⁺ neurons and the number of cells accumulating [³H]DA, in the same proportion (Figs. 5A and 6). Under these conditions, virtually all TH⁺ neurons contained the tritiated label (Fig. 6), which indicates that both peptide agonists saved a population of neurons that was equipped with a functional DA transporter. It is interesting that the neurites of TH⁺ neurons treated with GR73632 and [pro7]-NKB were more intensely labeled with [³H]DA than in control cultures, suggesting that the uptake was also more efficient in these neurons. Confirming this observation, the accumulation rate of [³H]DA per TH⁺ neuron was augmented by approximately 2-fold in the presence of the test compounds compared with untreated cultures (Fig. 5B). In contrast, the agonist of voltage-gated Na⁺ channels, veratridine, which also provided robust neuroprotection to DA neurons in this model system (Salthun-Lassalle et al., 2004), had




Fig. 4. The effects of NK₁ and NK₃ receptor agonists depend on the duration of treatment and the time at which it is initiated. Number of TH⁺ neurons in mesencephalic cultures exposed to 1 nM [pro7]-NKB at various time intervals between 0 and 10 DIV. **A**, [pro7]-NKB still increased survival when the treatment was postponed after plating. **B**, the effects of [pro7]-NKB were short-lived after withdrawal of the treatment. \star , $p < 0.05$, different from corresponding control cultures. Similar results were obtained with the NK₁ receptor agonist GR73632.

no effect on the accumulation rate of DA by these neurons (Figs. 5B and 6). Note the virtual absence of cells accumulating [^3H]DA in cultures treated with the inhibitor of the DA transporter GBR-12909 (0.5 μM) before the uptake (Fig. 6).

Trophic Effects Mediated by NK₁ and NK₃ Receptors Are Selective for DA Neurons in Mesencephalic Cultures. TH⁺ cells represent approximately 1 to 2% of all neurons at the time of plating. To determine whether NK₁ (GR73632) and NK₃ ([pro7]-NKB) receptor agonists also affected the survival of non-DA neurons, which are predominantly GABAergic and to a lesser extent serotonergic, we labeled the entire population of mesencephalic neurons with an antibody against MAP2. The number of MAP2⁺ neurons remained unchanged after long-term treatment with GR73632 or [pro7]-NKB (Fig. 7A), as illustrated in Fig. 7B. Similar results were obtained when the uptake of tritiated GABA or 5-HT was used to assess the function of GABAergic and serotonergic neurons, respectively (Fig. 7C).

Survival Promotion by Stimulation of NK₁ and NK₃ Receptors Does Not Result from an Antiproliferative Effect on Glial Cells. DA neurons can be rescued efficiently in this culture model by reducing the proliferation of astro-

cytes or their precursor cells (Mourlevat et al., 2003). To determine whether stimulation of NK₁ and NK₃ receptors affected the proliferation of glial cells, we assessed the incorporation of [methyl- ^3H]thymidine in cultures treated continually for 7 DIV with GR73632 or [pro7]-NKB (Fig. 8). Neither GR73632 nor [pro7]-NKB had an influence on the number of thymidine⁺ nuclei in mesencephalic cultures (Fig. 8, B and C), which indicates that NK₁ and NK₃ agonists promoted the survival of DA neurons despite the glial proliferation. As expected, however, the synthetic deoxynucleoside cytosine (araC, 1 μM) caused an almost complete loss of [methyl- ^3H]thymidine nuclei and a robust increase in the number of TH⁺ neurons (Fig. 8).

The Rescue of DA Neurons Caused By Activation of NK Receptors Does Not Depend on the Secretion of a Trophic Factor. We wished to determine whether the effect mediated by NK₁ and NK₃ receptors was possibly mediated by GDNF, a prototypical trophic factor for DA neurons. Therefore, we tested the effects of GR73632 (10 nM) and [pro7]-NKB (1 nM) in the presence of an anti-GDNF antibody




Fig. 5. DA neurons rescued by NK₁ and NK₃ agonists accumulate [^3H]DA efficiently. A, the number of TH⁺ neurons (■) and neurons accumulating [^3H]DA (□) in 10 DIV mesencephalic cultures treated continually with the NK₁ agonist (GR73632, 10 nM) or the NK₃ agonist ([pro7]-NKB, 1 nM), or with the agonist of voltage-gated sodium channels veratridine (ver, 0.8 μM). All three treatments increased the number of neurons that express TH and incorporate [^3H]DA to the same extent. B, GR73632 and [pro7]-NKB, but not veratridine, substantially increased the uptake of [^3H]DA per TH⁺ neuron. *, $p < 0.05$, different from corresponding control cultures at 10 DIV. In control cultures, the number of TH⁺ neurons was estimated at 880, and the accumulation rates of [^3H]DA were estimated at 25 fmol/min/16-mm culture well and 2.8×10^{-2} fmol/min/TH⁺ neuron, respectively.




Fig. 6. Illustration of the effects of NK₁ and NK₃ agonists on DA cell numbers and DA uptake. DA neurons from 10 DIV mesencephalic cultures identified by TH immunofluorescent cytochemistry (left) and subsequently by microautoradiographic detection of [^3H]DA (right). Cultures were treated between 0 and 10 DIV with 10 nM GR73632, 1 nM [pro7]-NKB, or 0.8 μM veratridine as described above. Control cultures in which [^3H]DA uptake was performed in the presence of 0.5 μM concentration of the DA uptake inhibitor GBR-12909 (bottom). White arrows, cell bodies of neurons labeled with [^3H]DA. Scale bar, 50 μm .

(AB 212-NA; 5 μ g/ml) that neutralizes the biological activity of the trophic peptide (Fig. 9A). Whereas the antibody blocked the increase in DA cell survival produced by GDNF (10 ng/ml), it had no effect on neuronal survival promoted by tachykinins. This indicates that the effect of tachykinins was not dependent on the secretion of GDNF in the culture medium.

To exclude the effect of another putative factor secreted by glial or neuronal cells, we assessed the survival of TH⁺ cells maintained in a culture medium that had been conditioned previously by mesencephalic cultures exposed continually to GR73632 (10 nM) or [pro7]-NKB (1 nM). The two conditioned media protected the DA neurons but only because they con-

tained GR73632 or [pro7]-NKB, the effects of which were blocked by the corresponding NK₁ and NK₃ receptor antagonists GR71251 and SB218795 (Fig. 9B).

Neuroprotection of DA Neurons by Tachykinins: Role of Voltage-Gated Ionic Channels. We have shown previously that voltage-gated ionic channels are crucially involved in the survival of DA neurons in this model system (Doughou et al., 2001; Salthun-Lassalle et al., 2004). In particular, we showed that DA neurons can be rescued from death by low-level activation of voltage-gated Na⁺ channels using the alkaloid veratridine (Salthun-Lassalle et al., 2004), the effects of which were blocked by the Na⁺ channel antagonist TTX. The effects of the agonists of NK₁ (GR73632) and NK₃ ([pro7]-NKB) receptors on DA neurons were prevented by TTX (100 nM), indicating that Na⁺ channels were also implicated in the effects of tachykinins (Fig. 10, A and C). This finding and the fact that suboptimal amounts of veratridine (0.2 μ M) were able to improve the protective effect of optimal concentrations of [pro7]-NKB (Fig. 10A) and GR73632 (data not shown) suggested that the alkaloid and tachykinins promoted DA cell survival by a mechanism that was common. This was apparently not the case, because the T-type calcium-channel blocker flunarizine (5 μ M), which




Fig. 7. The survival and trophic effects of NK₁ and NK₃ agonists are restricted to DA neurons. A, number of TH⁺ and MAP2⁺ neurons in mesencephalic cultures exposed for 10 days to selective NK₁ (GR73632, 10 nM) or NK₃ ([pro7]-NKB, 1 nM) receptor agonists. B, illustration showing that neither GR73632 nor [pro7]-NKB affected the number of MAP2⁺ cells. C, treatments with GR73632 and [pro7]-NKB strongly promoted the uptake of [³H]DA but had no effect on the uptake of [³H]GABA or [³H]5-HT. *, $p < 0.05$ different from corresponding control cultures. Control values for the number of TH⁺ and MAP2⁺ neurons per 16-mm culture well at 10 DIV were 1150 and 420×10^3 . Control values at 10 DIV for the uptakes of [³H]DA, [³H]GABA, and [³H]5-HT were 19 fmol/min/well, 3.8 pmol/min/well, and 8.8 fmol/min/well, respectively.




Fig. 8. Treatment with NK₁ and NK₃ tachykinin receptor agonists does not affect glial cell proliferation in mesencephalic cultures. The number of TH⁺ neurons at 10 DIV (A) and the number of thymidine⁺ nuclei at 7 DIV (B) in mesencephalic cultures maintained in the presence of the NK₁ agonist GR73632 (10 nM), the NK₂ agonist GR64349 (100 nM), and the NK₃ agonist [pro7]-NKB (1 nM). araC was used at 1 μ M. *, $p < 0.05$, different from corresponding control cultures. C, illustration of the effects of [pro7]-NKB and araC on the number of thymidine⁺ cells.

prevented the survival-promoting effect of the alkaloid (Salthun-Lassalle et al., 2004), was unable to inhibit the effects of [pro7]-NKB or GR73632 (Fig. 10B). The neuroprotective effects of GR73632 and [pro7]-NKB were also resistant to nimodipine (10 μ M), an inhibitor of the L-type calcium channels. They were reduced, however, by the snail toxin ω -conotoxin MVIIA (0.5 μ M; Fig. 10, B and C), indicating that inward calcium currents through N-type calcium channels probably played a key role in the survival of DA neurons mediated by NK₁ and NK₃ receptors. As expected, the protective effects of native tachykinins SP and NKB were also blocked by treatment with TTX and ω -conotoxin MVIIA (Fig. 10, A and B). It is interesting that the rescuing effect of veratridine, which was prevented by TTX, remained unaffected by ω -conotoxin MVIIA (Fig. 10B), confirming that veratridine and tachykinins acted by mechanisms that were partly related but not identical.

To confirm that tachykinin-mediated neuroprotection was linked to activation of inward Ca²⁺ currents, we measured intracellular calcium levels in the presence of the various test treatments (Fig. 10D). At concentrations of NK₁ and NK₃




Fig. 9. Role of GDNF or other trophic factors in the neuroprotective effects of NK₁ and NK₃ receptor agonists. A, the number of TH⁺ neurons in mesencephalic cultures treated with GDNF (10 ng/ml), NK₁ (GR73632), and NK₃ ([pro7]-NKB) receptor agonists in the presence or not of a neutralizing anti-GDNF antibody (AB 212 NA; 5 μ g/ml). B, the culture media conditioned by GR73632 or [pro7]-NKB-treated cultures (CM) increased the survival of DA neurons when transferred to receiving cultures. The effects of these conditioned media were completely inhibited by antagonists of NK₁ (GR71251) and NK₃ (SB218795) receptors, respectively. Note that the NK₂ receptor antagonist failed to inhibit the effect of the CM. *, $p < 0.05$, lower than corresponding cultures treated with GDNF; **, $p < 0.05$, lower than cultures maintained with conditioned medium (CM) alone. Parentheses indicate that GR73632 and [pro7]-NKB were not added directly to the cultures but were transferred with the CM.

Fig. 10. Role of voltage-gated Na⁺ and Ca²⁺ channels in the neuroprotective effect of tachykinins. A, number of TH⁺ neurons at 10 DIV in cultures that were exposed to veratridine (ver), GR73632, and [pro7]-NKB in the presence or not of TTX, an antagonist of the voltage-gated sodium channels. B, number of TH⁺ neurons at 10 DIV in cultures that were exposed to veratridine, GR73632, and [pro7]-NKB in the presence or not of flunarizine (flun), nimodipine (nimo), and ω -conotoxin MVIIA (ω -conot), antagonists of T-type, L-type, and N-type voltage-gated calcium channels, respectively. Veratridine (0.2 or 0.8 μ M as indicated above); GR73632 (10 nM); [pro7]-NKB (1 nM); TTX (100 nM); flunarizine (5 μ M); ω -conotoxin MVIIA (0.5 μ M). C, immunofluorescent detection of TH⁺ neurons in control cultures and in cultures treated continually for 10 days with [pro7]-NKB in the presence or not of TTX or ω -conotoxin (ω -con). Scale bar, 50 μ m. D, calcium-green-1 fluorescence levels in 7 DIV cultures. Treatments with TTX or ω -conotoxin were used to modulate the effects of veratridine, GR73632, and [pro7]-NKB. *, $p < 0.05$, different from corresponding control cultures; **, $p < 0.05$, significant inhibition by the channel antagonist.

agonists that promote optimal survival of DA neurons, we observed an increase in $[Ca^{2+}]_i$ of ~60 to 80% greater than control levels. The capacity of tachykinins to increase $[Ca^{2+}]_i$ was abolished by TTX (0.1 μ M) or ω -conotoxin MVIIA (0.5 μ M), which indicates that voltage-gated Na^+ channels and N-type calcium channels played a key role in this effect (Fig. 10D). Unlike ω -conotoxin MVIIA, however, the L-type calcium-channel antagonist nimodipine (10 μ M) and the T-type calcium-channel antagonist flunarizine (5 μ M) failed to reduce the increase in $[Ca^{2+}]_i$ induced by tachykinins (data not shown). Note that veratridine-mediated increase in intracellular calcium was blocked by TTX as expected but not by ω -conotoxin MVIIA (Fig. 10D).

Discussion

We demonstrate here that stimulation of NK_1 and NK_3 but not NK_2 receptors by native tachykinins or synthetic peptide agonists provides robust and selective neuroprotection against DA cell death in mesencephalic cultures. Neuroprotection was mimicked by but independent of GDNF. It resulted from a depolarizing effect mediated through the activation of TTX-sensitive sodium channels and N-type voltage-gated calcium channels.

Native and Synthetic Tachykinins Increase the Number of DA Neurons through Activation of NK_1 and NK_3 Receptors. We observed that the native tachykinins SP, NKA, and NKB substantially increased the number of TH^+ neurons when applied continually to mesencephalic cultures. SP, NKA, and NKB are believed to act preferentially at the NK_1 , NK_2 , and NK_3 receptor sites, respectively. We wished to determine precisely, however, which of these receptors was responsible for the effects of the native tachykinins on DA neurons. The specific NK_1 receptor antagonist GR71251 suppressed entirely the effect of SP on DA neurons, whereas a cotreatment combining GR71251 and the specific NK_3 receptor antagonist SB218795 was required to prevent the action of NKB. This suggests, therefore, that the effect of SP occurred via NK_1 receptors whereas that of NKB required activation of both NK_1 and NK_3 receptors. The effect of NKA seemed to be mediated entirely by its nonpreferential receptors because its action on DA neurons was resistant to the selective NK_2 receptor antagonist R396 and was suppressed by a treatment combining the NK_1 and NK_3 receptor antagonists. Confirming that NK_1 and NK_3 receptors were crucial for the effects of tachykinins on DA neurons, the selective NK_1 agonist GR73632 and the selective NK_3 agonist [pro7]-NKB increased the number of DA neurons, whereas the NK_2 agonist GR64349 had no effect. Therefore, only mRNA transcripts encoding the NK_1 and NK_3 receptors were detected in mesencephalic cultures.

The Activation of NK_1 and NK_3 Receptors Is Truly Neuroprotective for DA Neurons. The increase in the number of TH^+ neurons produced by stimulation of NK_1 or NK_3 receptors may result from several mechanisms. The treatments may induce the proliferation of putative DA precursor cells, but this is unlikely because all TH^+ neuroblasts have exited the cell cycle in the gestational day 15.5 rat mesencephalon used to generate the cultures (Rothman et al., 1980). Furthermore, the number of TH^+ cells in treated cultures never exceeded that of TH^+ neuroblasts detected just after plating, and [methyl- 3H]thymidine, a marker of

DNA synthesis, was never associated with TH^+ cells in tachykinin-treated cultures. The effect of the tachykinin agonists might be simply to restore the expression of TH within neurons that contain undetectable amounts of the enzyme as they entered a premorbid state (Hirsch et al., 1988; Michel and Agid, 1996). Even though tachykinins were reported to stimulate TH protein expression (Friedman et al., 1988), this possibility is doubtful, because the tachykinin agonists failed to rescue the fraction of TH^+ neurons that had already disappeared when initiation of the treatments was delayed. The observation that the pancaspase inhibitor Boc-D-FMK was also able to increase the number of DA neurons in this model system indicates that the tachykinins probably reversed an ongoing apoptotic process that is occurring spontaneously (Salthun-Lassalle et al., 2004). The underlying mechanism, however, was not related to that described for the neurotransmitter adenosine and certain antimitotics, which were effective in this culture model through the repression of glial cell proliferation (Michel et al., 1999). Finally, it is worth noting that DA neurons treated with NK_1 and NK_3 receptor agonists efficiently accumulated DA by active transport, which indicates that these neurons were perfectly functional when protected by tachykinins.

TTX-Sensitive Sodium Channels and N-Type Voltage-Gated Calcium Channels Are Implicated in Tachykinin-Mediated Neuroprotection. Two observations indicate that inward sodium currents were critical for tachykinin-mediated neuroprotection. The effect of NK_1 and NK_3 receptor agonists was mimicked by veratridine, an agonist of the voltage-gated Na^+ channels (Salthun-Lassalle et al., 2004) and prevented by TTX, an irreversible antagonist of these channels (Catterall, 1980). This result is consistent with electrophysiological data showing that stimulation of NK_3 receptors evokes excitatory responses in DA neurons of the substantia nigra (Nalivaiko et al., 1997). We have established previously that low-level activation of voltage-gated Na^+ channels by the alkaloid veratridine protects DA neurons via a mechanism that requires calcium influx through low-threshold activated T-type calcium channels (Salthun-Lassalle et al., 2004). T-type calcium channels are reported to be activated in response to SP in certain types of neurons (Ikeda et al., 2003). In our study, however, the effects of the tachykinin agonists were resistant to flunarizine, an inhibitor of these channels (Santi et al., 2002). High-threshold-activated L- and N-type calcium channels that have been shown to contribute to the effects of tachykinins in several model systems (Bayguinov et al., 2003; Sculptoreanu and de Groat, 2003) might also participate to tachykinin-mediated neuroprotection in mesencephalic cultures. L-type calcium currents are implicated in the rescue of mesencephalic DA neurons by depolarizing concentrations of K^+ (Douchou et al., 2001), but a selective antagonist of these channels, nimodipine, failed to inhibit the action of tachykinins on these neurons. In contrast, N-type calcium channels were probably essential for the survival-promoting effect of tachykinins. Indeed, blockade of these channels by the snail toxin ω -conotoxin MVIIA (Hirata et al., 1997) abolished neuroprotection by tachykinins. Consistent with this observation, the moderate increase in $[Ca^{2+}]_i$ elicited by tachykinins was also reversed by ω -conotoxin MVIIA. This elevation was also prevented by blocking Na^+ channels with TTX, which provides further support for the idea that intracellular calcium levels

were crucially involved in the survival of these neurons. Finally, the efficacy of TTX to prevent the increase in calcium produced by tachykinins is an indication that Na^+ inward currents may precede the activation of N-type Ca^{2+} channels.

The present results demonstrate that N-type calcium channels possess in common with L- and T-type calcium channels (Douhou et al., 2001; Salthun-Lassalle et al., 2004) the ability to modulate DA cell survival. These channels have different voltage ranges and rates of activation and inactivation (Hammond, 2001), which probably explains why their recruitment depends largely on the nature and intensity of the excitatory stimuli applied on DA neurons. Overall, the crucial need of DA neurons for electrical stimulation suggests that they lack adequate excitatory inputs in culture and that tachykinins could act as substitutes. It is interesting to note that only DA neurons in this model system seem dependent on such input. Finally, the rapid reversal of the effects of NK_1 and NK_3 receptor agonists when they are withdrawn indicates that tachykinin-dependent excitatory signals were a permanent requirement for long-term survival of DA neurons.

Is There a Link between the Survival Promotion Mediated by Tachykinins and GDNF? We wished to determine whether NK_1 and NK_3 receptor agonists could operate through a mechanism involving GDNF, a major neurotrophic factor for DA neurons (Choi-Lundberg et al., 1997). This was excluded because a neutralizing antibody that prevented the rescue of DA neurons by GDNF did not interfere with the effects of the NK_1 or NK_3 receptor agonists. Other autocrine or paracrine trophic factors were probably not implicated either, as shown by our experiments with conditioned media. These experiments support the idea that tachykinins and synthetic peptide agonists acted directly on DA neurons, which express both NK_1 (Futami et al., 1998) and NK_3 receptors (Chen et al., 1998; Friedman et al., 2002). Doubt remains, however, because the presence of NK_1 receptors on DA neurons is still a question of debate (Futami et al., 1998), and the experiments using conditioned culture media might have failed to detect molecules having a short-range diffusion or a limited lifespan. Finally, because GDNF is known to stimulate the synthesis of substance P (Ogun-Muyiwa et al., 1999), particularly in the striatum (Humpel et al., 1996), we envisaged the possibility that some of its effects might be mediated by tachykinins. This was unlikely because TTX, which prevented the rescuing effect of tachykinins on DA neurons, failed to reduce that of GDNF on the same neurons (Salthun-Lassalle et al., 2004). Therefore, as expected, the antagonists of NK_1 and NK_3 receptors failed to reduce the survival-promoting activity of GDNF on DA neurons as well (data not shown).

In conclusion, we have shown that stimulation of the NK_1 and NK_3 receptors by native or synthetic agonists exerts neuroprotective effects that are highly selective for DA neurons in mesencephalic cultures. This indicates that these neurons may depend in part on tachykinins for their survival during development. It is interesting that survival promotion mediated by tachykinins in mesencephalic cultures was mimicked by GDNF, a trophic factor that prevents the death of DA neurons during development (Oo et al., 2003). Because GDNF is still effective when applied exogenously to injured DA neurons in the adult brain (Choi-Lundberg et al., 1997;

Dauer and Przedborski, 2003), it is reasonable to ask whether tachykinins could be neuroprotective in neurodegenerative conditions in which DA neurons die selectively, such as Parkinson's disease and related disorders.

Acknowledgments

We are grateful to Merle Ruberg for helpful suggestions on the manuscript.

References

- Agid Y (1991) Parkinson's disease: pathophysiology. *Lancet* **337**:1321–1324.
- Bayguinov O, Hagen B, and Sanders KM (2003) Substance P modulates localized calcium transients and membrane current responses in murine colonic myocytes. *Br J Pharmacol* **138**:1233–1243.
- Betarbet R and Greenamyre JT (2004) Regulation of dopamine receptor and neuropeptide expression in the basal ganglia of monkeys treated with MPTP. *Exp Neurol* **189**:393–403.
- Calvo N, Reiriz J, Perez-Navarro E, and Alberch J (1996) Tachykinins protect cholinergic neurons from quinolinic acid excitotoxicity in striatal cultures. *Brain Res* **740**:323–328.
- Catterall WA (1980) Neurotoxins that act on voltage-sensitive sodium channels in excitable membranes. *Annu Rev Pharmacol Toxicol* **20**:15–43.
- Chen LW, Guan ZL, and Ding YQ (1998) Mesencephalic dopaminergic neurons expressing neurexin K receptor (NK_3): a double immunocytochemical study in the rat. *Brain Res* **780**:150–154.
- Choi-Lundberg DL, Lin Q, Chang YN, Chiang YL, Hay CM, Mohajeri H, Davidson BL, and Bohn MC (1997) Dopaminergic neurons protected from degeneration by GDNF gene therapy. *Science (Wash DC)* **275**:838–841.
- Dauer W and Przedborski S (2003) Parkinson's disease: mechanisms and models. *Neuron* **39**:889–909.
- Dawson TM and Dawson VL (2002) Neuroprotective and neurorestorative strategies for Parkinson's disease. *Nat Neurosci* **5** (Suppl):1058–1061.
- Douhou A, Troade JD, Ruberg M, Raisman-Vozari R, and Michel PP (2001) Survival promotion of mesencephalic dopaminergic neurons by depolarizing concentrations of K^+ requires concurrent inactivation of NMDA or AMPA/kainate receptors. *J Neurochem* **78**:163–174.
- Friedman WJ, Dreyfus CF, McEwen B, and Black IB (1988) Presynaptic transmitters and depolarizing influences regulate development of the substantia nigra in culture. *J Neurosci* **8**:3616–3623.
- Friedman Y, Richter A, Raymond R, Loscher W, and Norega JN (2002) Regional decreases in NK_3 , but not NK_1 tachykinin receptor binding in dystonic hamster (dt/sz) brains. *Neuroscience* **112**:639–645.
- Futami T, Hatanaka Y, Matsushita K, and Furuya S (1998) Expression of substance P receptor in the substantia nigra. *Brain Res Mol Brain Res* **54**:183–198.
- Hammond C (2001) Subliminal voltage-gated currents of the somato-dendritic membrane, in *Cellular and Molecular Neurobiology*, pp 358–373, Academic Press, London.
- Hirata H, Albillos A, Fernandez F, Medrano J, Jurkiewicz A, and Garcia AG (1997) omega-Conotoxins block neurotransmission in the rat vas deferens by binding to different presynaptic sites on the N-type Ca^{2+} channel. *Eur J Pharmacol* **321**:217–223.
- Hirsch E, Graybiel AM, and Agid YA (1988) Melanized dopaminergic neurons are differentially susceptible to degeneration in Parkinson's disease. *Nature (Lond)* **334**:345–348.
- Hokfelt T, Reid M, Herrera-Marschitz M, Ungerstedt U, Terenius L, Hakanson R, Feng DM, and Polkers K (1991) Tachykinins and related peptides in the substantia nigra and neostriatum. *Ann NY Acad Sci* **632**:192–197.
- Humpel C, Marksteiner J, and Saria A (1996) Glial-cell-line-derived neurotrophic factor enhances biosynthesis of substance P in striatal neurons in vitro. *Cell Tissue Res* **286**:249–255.
- Ikeda H, Heinke B, Ruscheweyh R, and Sandkuhler J (2003) Synaptic plasticity in spinal lamina I projection neurons that mediate hyperalgesia. *Science (Wash DC)* **299**:1237–1240.
- Katsuki H, Takenaka C, Kume T, Kaneko S, and Akaike A (2001) Requirement of neural activity for the maintenance of dopaminergic neurons in rat midbrain slice cultures. *Neurosci Lett* **300**:166–170.
- Khawaja AM and Rogers DF (1996) Tachykinins: receptor to effector. *Int J Biochem Cell Biol* **28**:721–738.
- Kowall NW, Beal MF, Busciglio J, Duffy LK, and Yankner BA (1991) An in vivo model for the neurodegenerative effects of β amyloid and protection by substance P. *Proc Natl Acad Sci USA* **88**:7247–7251.
- Lallemend F, Lefebvre PP, Hans G, Rigo JM, Van de Water TR, Moonen G, and Malgrange B (2003) Substance P protects spiral ganglion neurons from apoptosis via PKC-Ca^{2+} -MAPK/ERK pathways. *J Neurochem* **87**:508–521.
- Mauborgne A, Javoy-Agud F, Legrand JC, Agid Y, and Cesselin F (1983) Decrease of substance P-like immunoreactivity in the substantia nigra and pallidum of parkinsonian brains. *Brain Res* **268**:167–170.
- Michel PP and Agid Y (1996) Chronic activation of the cyclic AMP signaling pathway promotes development and long-term survival of mesencephalic dopaminergic neurons. *J Neurochem* **67**:1633–1642.
- Michel PP, Marien M, Ruberg M, Colpaert F, and Agid Y (1999) Adenosine prevents the death of mesencephalic dopaminergic neurons by a mechanism that involves astrocytes. *J Neurochem* **72**:2074–2082.
- Mourlevat S, Troade JD, Ruberg M, and Michel PP (2003) Prevention of dopaminergic neuronal death by cyclic AMP in mixed neuronal/glia mesencephalic cul-

- tures requires the repression of presumptive astrocytes. *Mol Pharmacol* **64**:578–586.
- Nalivaiko E, Michaud JC, Soubrie P, Le Fur G, and Feltz P (1997) Tachykinin neurokinin-1 and neurokinin-3 receptor-mediated responses in guinea-pig substantia nigra: an in vitro electrophysiological study. *Neuroscience* **78**:745–757.
- National Research Council (1996) *Guide for the Care and Use of Laboratory Animals*, National Academies Press, Washington, DC.
- Ogun-Muyiwa P, Helliwell R, McIntyre P, and Winter J (1999) Glial cell line derived neurotrophic factor (GDNF) regulates VR1 and substance P in cultured sensory neurons. *Neuroreport* **10**:2107–2111.
- Oo TF, Kholodilov N, and Burke RE (2003) Regulation of natural cell death in dopaminergic neurons of the substantia nigra by striatal glial cell line-derived neurotrophic factor in vivo. *J Neurosci* **23**:5141–5148.
- Pennefather JN, Lecci A, Candenas ML, Patak E, Pinto FM, and Maggi CA (2004) Tachykinins and tachykinin receptors: a growing family. *Life Sci* **74**:1445–1463.
- Pinto FM, Armesto CP, Magraner J, Trujillo M, Martin JD, and Candenas ML (1999) Tachykinin receptor and neutral endopeptidase gene expression in the rat uterus: characterization and regulation in response to ovarian steroid treatment. *Endocrinology* **140**:2526–2532.
- Rothman TP, Specht LA, Gershon MD, Joh TH, Teitelman G, Pickel VM, and Reis DJ (1980) Catecholamine biosynthetic enzymes are expressed in replicating cells of the peripheral but not the central nervous system. *Proc Natl Acad Sci USA* **77**:6221–6225.
- Salthun-Lassalle B, Hirsch EC, Wolfart J, Ruberg M, and Michel PP (2004) Rescue of mesencephalic dopaminergic neurons in culture by low-level stimulation of voltage-gated sodium channels. *J Neurosci* **24**:5922–5930.
- Santi CM, Cayabyab FS, Sutton KG, McRory JE, Mezeyova J, Hamming KS, Parker D, Stea A, and Snutch TP (2002) Differential inhibition of T-type calcium channels by neuroleptics. *J Neurosci* **22**:396–403.
- Sculptoreanu A and de Groat WC (2003) Protein kinase C is involved in neurokinin receptor modulation of N- and L-type Ca^{2+} channels in DRG neurons of the adult rat. *J Neurophysiol* **90**:21–31.
- Severini C, Improta G, Falconieri-Erspamer G, Salvadori S, and Erspamer V (2002) The tachykinin peptide family. *Pharmacol Rev* **54**:285–322.
- Stacey AE, Woodhall GL, and Jones RS (2002) Neurokinin-receptor-mediated depolarization of cortical neurons elicits an increase in glutamate release at excitatory synapses. *Eur J Neurosci* **16**:1896–1906.
- Troade JD, Marien M, Mourlevat S, Debeir T, Ruberg M, Colpaert F, and Michel PP (2002) Activation of the mitogen-activated protein kinase ($\text{ERK}_{1/2}$) signaling pathway by cyclic AMP potentiates the neuroprotective effect of the neurotransmitter noradrenaline on dopaminergic neurons. *Mol Pharmacol* **62**:1043–1052.
- Yang YL, Yao KH, and Li ZW (2003) Similarities of SP-, NKA- and NKB-induced currents in rat dorsal root ganglion neurons. *Brain Res* **991**:18–25.

Address correspondence to: Dr. Patrick P. Michel, INSERM 679, Bâtiment Pharmacie, Hôpital de la Salpêtrière, 47 bd de l'hôpital, 75013, Paris, France. E-mail: ppmichel@ecr.jussieu.fr
