

# A Direct Interaction between the N Terminus of Adenylyl Cyclase AC8 and the Catalytic Subunit of Protein Phosphatase 2A

Andrew J. Crossthwaite, Antonio Ciruela, Timothy F. Rayner, and Dermot M. F. Cooper

*Department of Pharmacology, University of Cambridge, Cambridge, United Kingdom*

Received August 18, 2005; accepted October 28, 2005

## ABSTRACT

Although protein scaffolding complexes compartmentalize protein kinase A (PKA) and phosphodiesterases to optimize cAMP signaling, adenylyl cyclases, the sources of cAMP, have been implicated in very few direct protein interactions. The N termini of adenylyl cyclases are highly divergent, which hints at isoform-specific interactions. Indeed, the  $\text{Ca}^{2+}$ -sensitive adenylyl cyclase 8 (AC8) contains a  $\text{Ca}^{2+}$ /calmodulin binding site on the N terminus that is essential for stimulation of activity by the capacitative entry of  $\text{Ca}^{2+}$  in the intact cell. Here, we have used the N terminus of AC8 as a bait in a yeast two-hybrid screen of a human embryonic kidney (HEK) 293 cell cDNA library and identified the catalytic subunit of the serine/threonine protein phosphatase 2A (PP2A<sub>C</sub>) as a binding partner. Confirming the highly specific nature of this novel interaction, glutathione-S-transferase fusion proteins containing the full-length N terminus

of AC8 affinity precipitated catalytically active PP2A<sub>C</sub> from both HEK293 and mouse forebrain membranes—the latter a normal source of AC8. The scaffolding subunit of PP2A (PP2A<sub>A</sub>; 65 kDa) was also precipitated by the N terminus of AC8, indicating that AC8 may occur in a complex with the PP2A core dimer. The interaction between the N terminus of AC8 and PP2A<sub>C</sub> was antagonized by  $\text{Ca}^{2+}$ /calmodulin. However, PP2A<sub>C</sub> and  $\text{Ca}^{2+}$ /calmodulin did not share identical binding specificities in the N terminus of AC8. PKA-mediated phosphorylation did not influence either calmodulin or PP2A<sub>C</sub> association with AC8. In addition, both PP2A<sub>C</sub> and AC8 occurred in lipid rafts. These findings are the first demonstration of an association between adenylyl cyclase and any downstream element of cAMP signaling.

Many of what have later turned out to be generalities in signal transduction were first established in cAMP signaling. Thus, it is surprising that although phosphodiesterases and PKA have all been shown to be involved in scaffolding complexes, AC, the instigator of the pathway, has not been found until now to be directly associated with any downstream regulatory element. The organization of regulatory proteins into macromolecular complexes is now expected to be encountered as a device to optimize the efficiency of signal transduction, particularly between different regulatory pathways (Smith and Scott, 2002). The interaction between  $\text{Ca}^{2+}$ - and

cAMP-regulated pathways occurs at multiple levels, with the earliest modulation occurring at the site of cAMP synthesis. Of the nine membrane-bound isoforms of adenylyl cyclase (AC), five either are stimulated or inhibited by physiological increases in  $[\text{Ca}^{2+}]_i$ . AC1 and AC8 are activated by  $[\text{Ca}^{2+}]_i$  in a calmodulin-dependent manner, whereas AC5 and AC6 are inhibited by  $\text{Ca}^{2+}$  independently of calmodulin (Cooper, 2003). In the intact cell,  $\text{Ca}^{2+}$ -regulated adenylyl cyclases selectively respond to  $\text{Ca}^{2+}$  entry through either capacitative calcium entry (CCE) channels or voltage-gated calcium channels (VGCCs) (Fagan et al., 2000a). A growing body of evidence suggests that  $\text{Ca}^{2+}$ -sensitive adenylyl cyclases and CCE channels are functionally colocalized (Cooper et al., 1994; Chiono et al., 1995; Fagan et al., 1996, 1998, 2000a; Smith et al., 2002), a situation which is reinforced by their compartmentalization in plasma membrane domains rich in cholesterol and sphingolipids, known as lipid rafts (Fagan et

This work was funded by the Wellcome Trust and by National Institutes of Health grant GM32483 (to D.M.F.C.).

A.J.C. and A.C. contributed equally to this work.

Article, publication date, and citation information can be found at <http://molpharm.aspetjournals.org>.  
doi:10.1124/mol.105.018275.

**ABBREVIATIONS:** PKA, protein kinase A; AC, adenylyl cyclase; CCE, capacitative  $\text{Ca}^{2+}$  entry; GFP, green fluorescent protein; GST, glutathione S-transferase; mAb, monoclonal antibody; NR, N-methyl-D-aspartate receptor subunit; PSD, postsynaptic density; PP2A, protein phosphatase 2A; PP2A<sub>A</sub>, protein phosphatase 2A scaffolding subunit A; PP2A<sub>B</sub>, protein phosphatase 2A regulatory subunit B; PP2A<sub>C</sub>, protein phosphatase 2A catalytic subunit; TRP, transient receptor potential; VGCC, voltage-gated calcium channel; PCR, polymerase chain reaction; PAGE, polyacrylamide gel electrophoresis; DIV, days in vitro; TNE buffer, NaCl/EDTA/Tris; TBS, Tris-buffered saline; TTBS, Tris-buffered saline/Tween 20; NP40, Nonidet P-40; TRP, transient receptor potential.

al., 2000b; Smith et al., 2002). The presence of  $\text{Ca}^{2+}$ -sensitive adenylyl cyclases in such domains may permit specific interactions with other signaling proteins present in rafts (Davare et al., 2001; Lavigne et al., 2002; Foster et al., 2003). However, very few proteins have been identified that specifically interact with adenylyl cyclases.

AC8 is predominantly expressed in brain areas associated with learning and memory (Mons et al., 1998). Indeed, AC8 knockout mice display deficits in hippocampal long-term potentiation, a cellular correlate of memory formation, which involves dynamic alterations in  $[\text{Ca}^{2+}]_i$  and cAMP (Frey et al., 1993; Poser and Storm, 2001; Wang et al., 2003). The molecular basis whereby an increase in  $[\text{Ca}^{2+}]_i$  regulates AC8 activity is becoming clearer (Gu and Cooper, 1999). The N terminus of AC8 contains an amphipathic  $\alpha$ -helical calmodulin-binding domain, which is absolutely essential for stimulation by  $\text{Ca}^{2+}$  in the intact cell (Gu and Cooper, 1999; Smith et al., 2002). Deletion of the N terminus results in an activity that, although still able to be stimulated by  $\text{Ca}^{2+}$ /calmodulin in vitro, is no longer able to be stimulated by CCE in the intact cell (Smith et al., 2002). The N termini of all adenylyl cyclase isoforms are highly divergent and might be expected to function in isozyme-specific interactions. Indeed, recently, using a yeast two-hybrid approach, the N terminus of AC6 was found to interact with snapin, a component of the soluble N-ethylmaleimide-sensitive factor attached protein receptor complex (Chou et al., 2004). Because we were convinced that the N terminus of AC8 played a critical role in regulation in the intact cell, we adopted a yeast two-hybrid strategy to search for regulatory partners of AC8. We found that, in addition to associating with calmodulin, the N terminus of AC8 interacts with the catalytic subunit of protein phosphatase 2A (PP2A<sub>C</sub>). The N terminus of AC8 pulled down PP2A<sub>C</sub> not only from HEK293 cell lysates, but also from brain homogenates. In addition, AC8 and PP2A<sub>C</sub> were both detected in lipid rafts, which suggests that PP2A may be situated to efficiently control downstream kinase activity, initiated by the  $\text{Ca}^{2+}$ -dependent activation of AC8. These findings identify a potentially important direct interaction between adenylyl cyclase and a downstream component of the cAMP signaling cascade.

## Materials and Methods

**Materials.** Monoclonal flotillin and PP2A catalytic subunit antibodies were obtained from BD Biosciences Transduction Laboratories (Erembodegem, Belgium). Monoclonal calmodulin antibody was from Upstate Biotechnology (Lake Placid, NY). Polyclonal PP2A structural subunit A and the polyclonal pan-PP2A<sub>B</sub> antibody were a gift from Dr. Brian Wadzinski (Department of Pharmacology, Vanderbilt University, Nashville, TN). Polyclonal  $\text{G}\alpha_{\text{olf}}$  and  $\text{PKA}\alpha$  catalytic subunit (C-20) antibodies were from Santa Cruz Biotechnology (Santa Cruz, CA). Horseradish peroxidase-conjugated goat anti-rabbit IgG was from GE Healthcare (Little Chalfont, Buckinghamshire, UK), and horseradish peroxidase-conjugated goat anti-mouse IgG conjugated was from Promega (Madison, WI). Paraformaldehyde was obtained from TAAB Laboratories (Aldermaston, UK). Tissue culture media and mammalian protease inhibitor cocktail were purchased from Sigma Chemical (Poole, Dorset, UK). His-Select cobalt affinity gel was from Sigma, and glutathione-Sepharose was from GE Healthcare.

**Yeast 2-Hybrid System.** The N terminus of rat AC8 (residues 1–179) was PCR-amplified from a plasmid containing full-length AC8 (construct 8Nt). The following oligonucleotide primers were

used: 5'-ccg aat tca tgg aac tct cgg atg tgc act gcc tta g-3' (primer r\_AC8-Nt-F) and 5'-atg gat ccc tcc gat ttg cgc ctc tgg ccc agg aa-3' (primer r\_AC8-Nt-R). The resulting PCR product was digested with EcoRI and BamHI restriction enzymes and subcloned between the EcoRI and BamHI sites of plasmid pGBK-T7 (BD Biosciences). The PCR-amplified insert (pGBK-8Nt) was sequenced to check for errors, and then the vectors were tested for self-activation in the yeast two-hybrid system alongside the empty transcription-activation domain vector pGAD-T7 (BD Biosciences). Plasmids pGBK-8Nt failed to self-activate. The AC8 N terminus encoding the DNA fragment was also subcloned between the EcoRI and XhoI sites of plasmid pGAD-T7 (generating pGAD-8Nt1) to allow interactions detected in the screen to be confirmed in both directions. Yeast strain AH109 (genotype, *MATa trp1-901 leu2-3,112 ura3-52 his3-200 gal4Δgal80Δ LYS2::GAL1<sub>UAS</sub>-GAL1<sub>TATA</sub>-HIS3 MEL1 GAL2<sub>UAS</sub>-GAL2<sub>TATA</sub>-ADE2, URA3::MEL1<sub>UAS</sub>-MEL1<sub>TATA</sub>-lacZ*), a human kidney cDNA library (with  $2.5 \times 10^6$  independent clones and average cDNA size of 1.5 kilobases), as the source of cDNA and plasmids used in this study were obtained from BD Biosciences as part of their Matchmaker 3 system. For the stringent selection of interacting clones, media lacking histidine, leucine, tryptophan, and adenine were used.

**Production and Expression of Recombinant Proteins.** The N terminus of rat AC8 was PCR-amplified from a plasmid containing full-length AC8 from residues 1 to 179 (construct 8Nt). The following oligonucleotide primers were used: 5'-ccg aat tca tgg aac tct cgg atg tgc act gcc tta g-3' and 5'-cgc gtc gac tta ctc cga ttt gcg cct ctg g-3'. The resulting PCR product was digested with EcoRI and SalI restriction enzymes and subcloned between the EcoRI and SalI sites of plasmid pGEX4T (GE Healthcare) to produce a fusion protein between glutathione-S-transferase (GST) and NtAC8. The C terminus of rat AC8 was PCR-amplified from a plasmid containing full-length AC8, from residues 1106 to 1248 (construct C2bAC8). The following oligonucleotide primers were used: 5'-gcg age tgc aca ttt ggg gta aaa ctg-3' and 5'-acg cgt cga ctt atg gca aat cgg att tg-3'. The resulting PCR product was digested with SacI and SalI restriction enzymes and subcloned between the SacI and SalI sites of plasmid pQE30 (QIAGEN, Valencia, CA) to produce a fusion protein containing 6× histidines at the Nt of C2bAC8. Both fusion proteins were expressed in *Escherichia coli* XL10 Gold and purified either on glutathione-Sepharose (GST-NtAC8) or cobalt affinity gel (His-C2bAC8).

**Preparation of HEK293 Crude Membranes.** HEK293 cells were detached with phosphate-buffered saline containing 0.03% EDTA and centrifuged at 195g for 5 min. The supernatant was removed and the pellet resuspended in hypotonic lysis buffer (10 mM Tris, 1 mM EDTA, 1 mM EGTA, and protease inhibitors, pH 8.0). After 10 min, cells were homogenized at 4°C by 50 strokes in a tight-fitting Dounce homogenizer followed by centrifugation (195g at 4°C for 5 min). The supernatant was centrifuged at 17,257g (15 min, 4°C). The supernatant was removed, and the pellet, representing crude membranes, was collected.

**Preparation of Mouse Forebrain Crude Membranes.** Adult mouse forebrains were dissected into ice-cold buffer consisting of 50 mM Tris, 1 mM  $\text{MgCl}_2$ , 1 mM EDTA, 1 mM 4-(2-aminoethyl) benzenesulfonyl fluoride, 1 mM benzamidine, and 1  $\mu\text{g}$  of DNAase, pH 7.4, and lysed by passing through a 0.22-gauge needle 20 times. After further centrifugation (195g at 4°C for 5 min) and dissociation, the lysate was centrifuged at 17,257g (15 min, 4°C). The supernatant was removed, and the pellet, representing crude membranes, was collected.

**AC8 N- and C-Terminal Affinity Precipitation.** Isolated crude membranes were solubilized in 2% SDS solubilization buffer (50 mM Tris, 150 mM NaCl, 1 mM EGTA, 1 mM EDTA, and protease inhibitors, pH 7.4) (Leonard et al., 1998) followed by centrifugation (17,257g at 4°C for 15 min). The supernatant was diluted (1:20) in binding buffer (50 mM phosphate buffer, pH 7.4, 150 mM NaCl, 0.2% Triton X-100, 1 mM EDTA, and 1 mM EGTA) and precleared for 30 min or more with 5  $\mu\text{l}$  of a 50% suspension of GST glutathione-Sepharose. GST-NtAC8 glutathione-Sepharose that had been

washed three times in dilution buffer was added to the precleared sample and rotated for 3 h at 4°C. Otherwise, for in vitro pull-downs, GST or cobalt affinity gels were used as controls, as noted in the figure legends. The Sepharose/cobalt beads were collected and washed four times in ice-cold binding buffer, resuspended in 2× boiling buffer (final concentration, 125 mM Tris, 300 mM dithiothreitol, 20% glycerol, and 0.004% bromophenol blue, pH 6.8) and heated at 100°C for 5 min. Supernatant was collected, and samples were stored at –80°C. The results shown are representative of at least five experiments with similar results.

**Phosphorylation Assay.** GST-NtAC8, His-C2bAC8, or myelin basic protein, as positive control, were incubated with either a cell-free extract of forskolin (20 μM) and prostaglandin E<sub>1</sub> (100 nM)-induced HEK293 cells or the catalytic subunit of PKA (Sigma) in phosphorylation buffer incubated with [ $\gamma$ -<sup>32</sup>P]ATP (10 μCi; GE Healthcare) (50 mM Tris-HCl, pH 7.5, 10 mM MgCl<sub>2</sub>, 1 mM EDTA, 10 μM ATP, 2 mM dithiothreitol, 0.01% Triton X-100, 20 μg/ml aprotinin, 20 μg/ml leupeptin, and 10 μg/ml pepstatin A), with or without six units of PKA (Sigma) at 30°C for 20 min. After separation by SDS-PAGE, phosphorylated proteins were detected by autoradiography.

**Dephosphorylation Assay.** myelin basic protein (250 μg) was phosphorylated as described above with 24 units of PKA in phosphorylation buffer incubated with [ $\gamma$ -<sup>32</sup>P]ATP. The reaction was terminated by filtration in a Microcon YM-3 filter (Millipore Corporation, Billerica, MA) and washed with 500 μl of 20 mM Tris-HCl and 140 mM NaCl, pH 7.6. The phosphorylated myelin basic protein was resuspended in 100 μl of 12.5 mM Tris-HCl and 25 μM CaCl<sub>2</sub>, pH 7.6, and 25-μl fractions were mixed with the resultant affinity precipitates (GST or GST-NtAC8) from Triton X-100-solubilized HEK293 cells. After incubation (30 min at 30°C), the reaction was terminated by the addition of 6 μl of 5× boiling buffer and heated at 100°C for 5 min. Samples were run on a 13% SDS-PAGE gel and exposed to Hyperfilm film for 17 h (GE Healthcare).

**Immunofluorescence.** Rat hippocampal neurons were cultured as described previously (Chawla et al., 2003) and transfected with N-terminally tagged GFP-AC8 (0.4 μg of cDNA) after 8 DIV using Lipofectamine 2000 (Invitrogen, Carlsbad, CA). Forty-eight hours after transfection, neurons were washed with phosphate-buffered saline (12.1 mM Na<sub>2</sub>HPO<sub>4</sub>, 4 mM KH<sub>2</sub>PO<sub>4</sub>, and 130 mM NaCl, pH 7.4) and fixed using 4% paraformaldehyde (1 h, 20°C). Otherwise, before fixing, neurons were prepermeabilized in 0.5% Triton X-100 at 4°C for 30 min (Hering et al., 2003). Coverslips were mounted in Antifade (Invitrogen) according to the manufacturer's procedures and the cells were visualized on a Zeiss Axiovert LSM510 confocal microscope (Carl Zeiss GmbH, Jena, Germany), using 40× and 63× oil immersion objectives. The scale bar in all images represents 10 μM.

**Detergent-Resistant Membrane Preparation.** Mouse forebrain crude membranes were incubated with ice-cold Triton X-100 in TNE buffer (150 mM NaCl, 5 mM EDTA, and 25 mM Tris) containing 0.1 M sodium carbonate, pH 11 (Ostermeyer et al., 1999). The suspension was transferred to a Dounce homogenizer and homogenized with 20 strokes and left on ice for 30 min. The homogenate was adjusted to 40% sucrose by the addition of 60% sucrose in TNE buffer, pH 7.4. The extract was placed below a 5 and 30% discontinuous sucrose gradient prepared in cold TNE buffer and centrifuged in a Beckman SW55 rotor at 24,000 rpm for 16 h at 4°C (Beckman Coulter, Fullerton, CA). Fractions (10 × 0.5 ml) were collected from the top of the gradient. The isolated fractions were diluted in TNE buffer and centrifuged in a Beckman SW55 rotor at 50,000 rpm for 1 h at 4°C. The pelleted membranes were resuspended in 1% SDS. Finally, samples were suspended in boiling buffer heated to 100°C for 5 min and stored at –80°C.

**Immunoblotting.** Proteins were resolved using SDS-PAGE 7.5% and 12% SDS-polyacrylamide gels and transferred to a polyvinylidene difluoride membrane. The polyvinylidene difluoride membrane was incubated in TBS blocking buffer (20 mM Tris, pH 7.5, 150 mM

NaCl) containing 5% nonfat dry milk for 30 min, followed by two 10-min washes in TBS supplemented with 0.05% (v/v) Tween 20 (TTBS). Membranes were incubated overnight at room temperature with anti-PP2A<sub>C</sub> mAb (1:5000), anti-PKAc<sub>at</sub> pAb (1:1000), anti-flo-tillin mAb (1:5000), anti-calmodulin mAb (1:5000), mAb anti-G $\alpha$ s-olf (1:1000), in TTBS containing 1% nonfat dry milk (antibody buffer). The membranes were washed (2 × 10 min) in TTBS and then incubated with goat anti-rabbit IgG conjugated to horseradish peroxidase (1:5000 dilution of stock) or goat anti-mouse IgG conjugated to horseradish peroxidase (1:3000) in antibody buffer for 1 h. Finally, the membranes were washed in TTBS (2 × 10 min), rinsed in TBS, and treated with enhanced chemiluminescence plus reagent and exposed to Hyperfilm (GE Healthcare).

## Results

**The N Terminus of AC8 Interacts Specifically with PP2A Catalytic Subunit.** AC8 lacking the N terminus is insensitive to regulation by Ca<sup>2+</sup> in the intact cell, despite being targeted to lipid rafts, which is a property that is essential for regulation by CCE (Smith et al., 2002). However, AC8 is fully stimulated by Ca<sup>2+</sup>/calmodulin in vitro (Smith et al., 2002). This observation suggests that the N terminus of AC8 might engage with elements of the CCE regulatory apparatus. To determine whether such proteins would interact with the N terminus of AC8, amino acids 1 to 179 (i.e., the full-length N terminus) was cloned into a bait vector to screen an HEK293 cDNA library using the yeast two-hybrid system. The first screen identified 77 colonies that survived on synthetic media lacking histidine. The plasmids were recovered into *E. coli* and sequenced. Duplicate clones and common false-positives (transcription factors and “housekeeping” genes, such as those involved in metabolism) were discarded, leaving a pool of 22 genes of potential interest. The bait and prey vectors were reversed and tested positive in every case. Of these positive interacting plasmids, the one with most potential interest was a peptide sequence (Ser<sup>120</sup> to Leu<sup>309</sup>) corresponding to the C terminus of the  $\alpha$ -catalytic subunit of the serine/threonine PP2A<sub>C</sub>. Other proteins that may yet be of some interest included a G protein, G $\beta$ <sub>2</sub>, a laminin receptor, and a WD repeat domain protein; the others were rather obscure.

The finding that PP2Ac interacted with the N terminus of the Ca<sup>2+</sup>/calmodulin-stimulatable AC8 was deemed extremely interesting because PP2Ac participates in a number of signaling complexes (Lebrin et al., 1999; Boudreau et al., 2002). To be specific, PP2Ac interacts with the C-terminal region of the  $\alpha$ <sub>1C</sub> L-type VGCC, which is known to be in a functional signaling complex in the rat forebrain with the  $\beta$ <sub>2</sub>-adrenergic receptor and an unidentified adenylyl cyclase isoform (Davare et al., 2000, 2001). No obvious candidates that might have been involved in regulating CCE were identified; therefore, further investigations centered on the association of the N terminus with PP2A<sub>C</sub>.

PP2Ac is a member of the PPP family of serine/threonine phosphatases, which have conserved catalytic regions. Evidence from the crystal structure of the PPP family member PP1c suggests that regulation, either through post-translational modification or binding proteins, occurs at the C terminus (Egloff et al., 1997; Barford et al., 1998). In an effort to narrow down more precisely the domain of PP2Ac that interacted with the N terminus of AC8, two halves of the positive interacting PP2Ac fragment (Ser<sup>120</sup> to Arg<sup>215</sup>) and (Arg<sup>215</sup> to



Leu<sup>309</sup>) were expressed as histidine fusion proteins in *E. coli*. However, the peptide fragments were only present as an aggregated form within inclusion bodies, preventing further binding studies.

AC8 is predominantly neuronal in expression, with the highest levels of mRNA detected in the cortex, hippocampus, and cerebellum (Cali et al., 1994). Therefore, solubilized mouse cortical membranes were used to confirm the selectivity of the interaction between the N terminus of AC8 and PP2A<sub>C</sub>, using a GST recombinant fusion protein containing the full-length N terminus of AC8 (GST-NtAC8). Initial GST pull-downs were carried out on cortical membranes solubilized with 1% Nonidet P-40 (NP40). Incubating GST or GST-NtAC8 with solubilized mouse forebrain membranes demonstrated that the full-length, native PP2A<sub>C</sub> (36 kDa) was affinity-precipitated by GST-NtAC8, compared with a far less extent by a gross excess of GST (Fig. 1A).


Other solubilization procedures were also investigated; GST-NtAC8 additionally affinity-precipitated PP2A<sub>C</sub> from whole-cell HEK293 lysates solubilized in 1% Triton X-100 (data not shown). The postsynaptic density (PSD) is the proposed localization of AC8 (Mons et al., 1995; Wang et al., 2003), and proteins in this domain are not solubilized by NP40 or Triton X-100 (Lau et al., 1996). Therefore we used 2% SDS, which solubilizes PSD proteins and enables the coimmunoprecipitation of interacting proteins in the PSD as a means of investigating whether any additional proteins interacted with GST-NtAC8 (Lau et al., 1996; Mehta et al., 2001). The dilution of SDS and addition of Triton X-100 reduce the potential of SDS to dissolve or denature proteins (Leonard et al., 1998). However, we observed no differences in the protein profile between pull-downs using cortical membranes solubilized with either SDS or NP40, as determined by in-gel silver nitrate staining (data not shown). There was a consistent reduction in nonspecific binding of PP2A<sub>C</sub> to GST when membranes were solubilized with SDS, and therefore, we opted for SDS solubilization for further studies (Fig. 1B).

GST-NtAC8 also affinity-precipitated PP2A<sub>C</sub> from solubilized HEK293 membranes (Fig. 1C). We believe in vitro GST

pull-downs to be a more accurate method than coimmunoprecipitation experiments to ascertain genuine interactions involving AC8 and native PP2A<sub>C</sub>. This is because it is extremely difficult to solubilize adenylyl cyclases to any degree of relative purity. Just how difficult this can be is clearly demonstrated with Western blots of adenylyl cyclases, which typically run at a much higher than expected molecular mass. Indeed genuine interactions between adenylyl cyclase and interacting proteins are believed to be lost during solubilization procedures (Chou et al., 2004). Taken together, the results obtained with the yeast two-hybrid analysis, which depend on correctly folded protein fragments for the detection of positive interactions and results from the GST-NtAC8 pull-down experiments, which examine interactions with full-length native PP2A<sub>C</sub>, clearly demonstrate that the N terminus of AC8 can efficiently interact with native PP2A catalytic subunit, even in the presence of a highly diverse range of membrane proteins.

**The N Terminus of AC8 Interacts with the PP2A Core Enzyme.** The core enzyme of PP2A is a dimer consisting of the catalytic subunit and a scaffolding subunit of 65 kDa (PP2A<sub>A</sub>), referred to as PP2A<sub>D</sub> (Janssens and Goris, 2001). A third regulatory B subunit, of which there are four separate families, can associate with the core enzyme to target the trimeric holoenzyme to specific subcellular locations. However, the core enzyme is functional without the B subunit (Kremmer et al., 1997). To determine whether the N terminus of AC8 associated with the PP2A core enzyme complex, GST-NtAC8 affinity precipitates were probed with antibodies to both PP2A<sub>C</sub> and PP2A<sub>A</sub>. GST-NtAC8 pulled down both PP2A<sub>A</sub> (Fig. 2A) and PP2A<sub>C</sub> (Fig. 2B) from the same solubilized brain membrane preparation, which demonstrated that the N terminus of AC8 interacts with the PP2A core enzyme complex. However, no immunoreactivity against the B subunit was detected from aliquots that were positive for the core enzyme (data not shown).


**The N Terminus of AC8 Interacts with Active PP2A<sub>C</sub>.** The PP2A enzyme complex interacts with a wide variety of proteins, suggesting that it may act as a scaffolding protein, in addition to its well-defined role as a serine/threonine phos-



**Fig. 1.** The N terminus of AC8 interacts with the catalytic subunit of protein phosphatase 2A. Mouse cortical membranes were solubilized by either 1% NP40 (A) or 2% SDS (B), precleared with an excess of GST, and incubated with a GST fusion protein containing the full-length N terminus of AC8 immobilized on glutathione-Sepharose beads. The affinity precipitates were resolved by SDS-PAGE and were immunoblotted with an antibody raised against the catalytic subunit of protein phosphatase 2A (anti-PP2A<sub>C</sub>). Aliquots from the same experiments were examined for protein loading by Coomassie staining of membranes or parallel in-gel silver nitrate staining. C, HEK293 membranes were solubilized by 2% SDS, precleared with an excess of GST, and incubated with a GST fusion protein containing the full-length N terminus of AC8 immobilized on glutathione-Sepharose beads. The affinity precipitates were resolved by SDS-PAGE and were immunoblotted with an antibody raised against the catalytic subunit of protein phosphatase 2A (anti-PP2A<sub>C</sub>). Aliquots from the same experiments were examined for protein loading by parallel in-gel silver nitrate staining.


phatase (Hsu et al., 1999; Davare et al., 2000; Chan and Sucher, 2001; Boudreau et al., 2002). Therefore, we believed that it was important to ask whether the fraction of PP2A<sub>C</sub> that interacts with the N terminus of AC8 is catalytically active by examining its ability to dephosphorylate myelin basic protein. HEK293 cells solubilized in 1% Triton X-100 were incubated with GST or GST-NtAC8, and the resultant pull-downs were incubated with PKA-phosphorylated myelin basic protein. There was a clear reduction in the phosphorylation state of myelin basic protein after incubation with GST-NtAC8 pull-downs compared with GST pull-down controls (Fig. 3). These results clearly show that the N terminus of AC8 interacts with catalytically active PP2A<sub>C</sub>.

**Mutually Exclusive Binding of Calmodulin and PP2A<sub>C</sub> to the N Terminus of AC8.** To further address the nature of the association between catalytically active PP2A<sub>C</sub> and the N terminus of AC8, we investigated the putative regulation of the interaction. The N terminus of AC8 contains a calmodulin binding site (Gu and Cooper, 1999) that is absolutely necessary for efficient stimulation by CCE in the intact cell (Smith et al., 2002). As expected, exogenous calmodulin was precipitated by GST-NtAC8 *in vitro* but only weakly by a great excess of GST (Fig. 4A). We therefore asked whether endogenous PP2A could bind to the N terminus of AC8 in the presence of added calmodulin. The preparations of solubilized brain membranes used in our experiments would reduce Ca<sup>2+</sup> to very low levels, because they contain the chelators EGTA and EDTA (at 1 mM), to prevent protease activation. As an initial exploration of the effects of Ca<sup>2+</sup>/calmodulin on the ability of GST-NtAC8 to interact with PP2A, EGTA and EDTA were omitted, which slightly reduced the affinity precipitation of PP2A<sub>C</sub> (Fig. 4B). Inclusion of 20 μM Ca<sup>2+</sup> further reduced the ability of GST-NtAC8 to interact with PP2A<sub>C</sub>. The inclusion of 0.5 μM exogenous calmodulin and 20 μM Ca<sup>2+</sup> eliminated any detectable interaction between GST-NtAC8 and PP2A<sub>C</sub>. At this concentration of exogenous calmodulin, GST-NtAC8 bound preferentially to calmodulin (Fig. 4C). Therefore, the binding of PP2A<sub>C</sub> and calmodulin to the N terminus of AC8 is mutually exclusive.



**Fig. 2.** The N terminus of AC8 interact with the PP2A core enzyme. A GST fusion protein containing the full-length N terminus of AC8 immobilized on glutathione-Sepharose beads was incubated with GST pre-cleared brain membranes. The affinity precipitates were resolved by SDS-PAGE and immunoblotted with an antibody raised against the scaffolding subunit of protein phosphatase 2A (anti-PP2A<sub>A</sub>) (A) or an antibody raised against the catalytic subunit of protein phosphatase 2A (anti-PP2A<sub>C</sub>) (B). The Coomassie-stained membrane is shown at the bottom.

**The Amino Acids in the N Terminus of AC8 that Are Critical for Calmodulin Binding Are Not Critical for PP2A<sub>C</sub> Binding.** Point mutations in the helical calmodulin binding domain between amino acids 34 and 51 in the N terminus of AC8 prevent Ca<sup>2+</sup> stimulation of AC8 in the whole cell (Smith et al., 2002). We generated a GST recombinant fusion protein containing the full-length N terminus of AC8 with the same six critical amino acids mutated to alanine (GST-Nt8M34). To verify that this prevented calmodulin binding, GST-Nt8M34, GST, and GST-NtAC8 were incubated in the presence of exogenous Ca<sup>2+</sup> and calmodulin *in vitro*. GST-Nt8M34 clearly showed no interaction with calmodulin compared with GST-NtAC8 (Fig. 5A). To determine the relative amounts of fusion proteins present in the individual incubations, the blot was stripped and reprobed for GST immunoreactivity (Fig. 5A, bottom blot). Densitometric analysis of this immunoblot demonstrated that for equal levels of GST-fusion protein, GST-Nt8M34 is unable to interact with calmodulin compared with GST-NtAC8 (Fig. 5A). Therefore, GST-Nt8M34, which is unable to bind calmodulin *in vitro*, was examined for its ability to interact with endogenous PP2A<sub>C</sub>. Solubilized mouse forebrain membranes were pre-cleared with an excess of GST and divided into two aliquots. GST-NtAC8 was added to one and GST-Nt8M34 to the other. Equal volumes from the resulting pull-down were compared for PP2A<sub>C</sub> immunoreactivity. Both GST-Nt8M34 and GST-NtAC8 could clearly affinity-precipitate PP2A<sub>C</sub> (Fig. 5B). This demonstrates that the amino acids critical for binding calmodulin are not critical for binding PP2A<sub>C</sub>. Therefore, if GST-Nt8M34 does not bind calmodulin, then the addition of exogenous calmodulin to solubilized mouse forebrain membranes would not be expected to prevent interaction of GST-



**Fig. 3.** The N terminus of AC8 interacts with catalytically active PP2A<sub>C</sub>. GST or a GST fusion protein containing the full-length N terminus of AC8 immobilized on glutathione-Sepharose beads was incubated with Triton X-100-solubilized HEK293 cell lysates. The resulting washed affinity precipitates were incubated with myelin basic protein that had been previously phosphorylated with the catalytic subunit of PKA in buffer containing [ $\gamma$ -<sup>32</sup>P]ATP. A, proteins were resolved by SDS-PAGE and the level of myelin basic protein dephosphorylation determined by autoradiography. B, the Coomassie-stained membrane is shown at the bottom.

Nt8M34 with PP2A<sub>C</sub>. Indeed, including 0.5  $\mu$ M calmodulin and 20  $\mu$ M Ca<sup>2+</sup> in the pull-down assay did not prevent the affinity precipitation of PP2A<sub>C</sub> by GST-Nt8M34 (Fig. 5C). Taken together, these results demonstrate that calmodulin and PP2A<sub>C</sub> have overlapping binding domains on the N terminus of AC8, but the precise amino acids that are essential for binding calmodulin are not essential for binding PP2A<sub>C</sub>.


**Calmodulin Binding to AC8 Is Not Regulated by Phosphorylation.** The ability of calmodulin to bind to its target sequence is regulated by phosphorylation in a wide range of proteins (Hofmann et al., 1994; Williams and Coluccio, 1995; Enyedi et al., 1997; Turner et al., 2004). For instance, endothelial nitric-oxide synthase, which contains an  $\alpha$ -helical calmodulin binding domain similar to that present on the N terminus of AC8, contains a threonine residue, which, upon phosphorylation, prevents calmodulin binding (Fleming et al., 2001). In addition, this site is dephosphorylated by PP2A, and this reversible phosphorylation regulates Ca<sup>2+</sup>/calmodulin-stimulated endothelial nitric-oxide synthase activity (Fleming et al., 2001; Greif et al., 2002). AC8 contains two calmodulin binding domains, both of which are essential for in vivo stimulation (Gu and Cooper, 1999); therefore the possibility can be considered that reversible phosphorylation of either site could regulate calmodulin binding and, hence, activity. There are putative PKA-phosphorylation sites at positions 46 and 66 in the N terminus and positions 1156 and 1164 in the C terminus. The second calmodulin-binding domain of AC8 is an IQ motif that is situated in the C2b domain. We generated recombinant C2b domain protein containing a hexa-His tag at the N terminus (His-C2bAC8) to determine whether this would bind calmodulin in an in vitro assay as a prelude to conducting phosphorylation experiments. Incubating His-C2bAC8 with exogenous Ca<sup>2+</sup> and calmodulin in vitro showed that the C2b domain specifically bound calmodulin (Fig. 6).

Therefore, we asked whether putative phosphorylation of the N terminus or C2b calmodulin binding domains could regulate calmodulin association and thereby provide a rationale for the interaction with PP2A. Incubating GST-NtAC8 or His-C2bAC8 with either the catalytic subunit of PKA

(PKAc<sub>at</sub>) or a lysate prepared from forskolin- and prostaglandin E<sub>1</sub>-stimulated HEK293 cells failed to demonstrate any ability of either domain to be phosphorylated. Myelin basic protein used as a positive control was phosphorylated by both PKAc<sub>at</sub> and the stimulated HEK293 lysate (data not shown). In a further experiment, mutating Ser66 (a putative PKA phosphorylation site in the N terminus immediately upstream of one calmodulin binding domain) to aspartate did not prevent calmodulin association (data not shown). We conclude that neither of the calmodulin binding domains of AC8 is directly regulated by PKA-mediated phosphorylation—and by implication, of course, that it is not a substrate for PP2A.

**Lipid Raft Localization of AC8 and PP2A<sub>C</sub>.** If reversible phosphorylation of AC8 is not a major regulatory influence on calmodulin binding, the association of AC8 with PP2A might represent a novel mechanism for regulation and/or localization of the phosphatase. When heterologously expressed in nonexcitable cells, AC8 occurs in lipid rafts. This specific membrane compartmentalization is essential for regulating AC8 activity in vivo (Smith et al., 2002). In the whole animal, AC8 is largely confined to the brain, and we therefore examined whether in primary hippocampal neurons a membrane compartmentalization occurs similar to that observed in non-neuronal cells. Hippocampal neurons that were transfected with AC8 tagged with GFP at the N terminus (GFP-AC8) displayed predominant plasma membrane labeling at the soma, with fluorescence extending into the dendritic network (Fig. 7B). Higher magnification of axons demonstrated a clear plasma membrane localization of GFP-AC8 (Fig. 7B, inset). This was in obvious contrast to neurons transfected with GFP alone, in which fluorescence was clearly intracellular in the soma and confined to intracellular puncta throughout the processes (Fig. 7, A and inset).


To examine the plasma membrane domain in which GFP-AC8 resides, neurons were prepermeabilized with Triton X-100 at 4°C before fixation. This procedure demonstrated the lipid raft targeting of the calmodulin binding protein GAP-43/neuromodulin in PC12 cells (Arni et al., 1998). In



**Fig. 4.** A competitive interaction between Ca<sup>2+</sup>/calmodulin and PP2A<sub>C</sub> for association with the N terminus of AC8. A, GST or GST fusion proteins containing the full-length N terminus of AC8 immobilized on glutathione-Sepharose were incubated in the absence (–) or presence (+) of 0.5  $\mu$ M calmodulin and 20  $\mu$ M Ca<sup>2+</sup>. The pull-downs were resolved by SDS-PAGE and immunoblotted with an antibody raised against calmodulin (anti-calmodulin). The Coomassie-stained membrane is shown at the bottom. B, a GST fusion protein containing the full-length N terminus of AC8 immobilized on glutathione-Sepharose beads was incubated with GST-purified brain membranes containing either 1 mM EDTA/EGTA, no exogenous additions, 20  $\mu$ M Ca<sup>2+</sup>, 0.5  $\mu$ M calmodulin, or 20  $\mu$ M Ca<sup>2+</sup> and 0.5  $\mu$ M calmodulin. The affinity precipitates were resolved by SDS-PAGE and immunoblotted with an antibody raised against the catalytic subunit of protein phosphatase 2A (anti-PP2A<sub>C</sub>). The Coomassie-stained membrane is shown at the bottom. C, a GST fusion protein containing the full-length N terminus of AC8 immobilized on glutathione-Sepharose beads was incubated with GST-purified brain membranes containing either 1 mM EDTA/EGTA or 20  $\mu$ M Ca<sup>2+</sup> and 0.5  $\mu$ M calmodulin. The affinity precipitates were resolved by SDS-PAGE and immunoblotted with an antibody raised against calmodulin (anti-calmodulin). The Coomassie-stained membrane is shown at the bottom.




addition, with the use of specific fluorophore lipid markers, cold Triton X-100 preferentially solubilizes unsaturated lipids in primary hippocampal neurons, leaving a subset of less-soluble cholesterol/sphingolipid-rich domains at the plasma membrane (Hering et al., 2003). Confocal imaging revealed that GFP-AC8 was localized in detergent-resistant membrane domains along the dendrites of positively labeled neurons (Fig. 7C and inset). There was no detectable fluorescence from prepermeabilized GFP control neurons (data not shown), most probably because the soluble GFP protein was washed out. It should be noted that prepermeabilization with Triton X-100 would not be expected to remove nonraft plasma membrane proteins associated with the cytoskeleton.




**Fig. 5.** Amino acids critical for  $\text{Ca}^{2+}$ /calmodulin association are not essential for binding  $\text{PP2A}_C$ . **A**, GST, a GST fusion protein containing the full-length N terminus of AC8, or a GST fusion protein containing the full-length N terminus of AC8 with point mutations in the calmodulin binding region (8M34) that were immobilized on glutathione-Sepharose beads were incubated in the presence or absence of  $20 \mu\text{M}$   $\text{Ca}^{2+}$  and  $0.5 \mu\text{M}$  calmodulin. After washing with 1% Triton X-100, the pull-downs were resolved by SDS-PAGE and immunoblotted with an antibody raised against calmodulin (anti-calmodulin). The same blot stripped and reprobed with an antibody raised against GST (anti-GST) is shown at the bottom. The relative amount of the three fusion proteins was quantified by densitometric analysis of the GST immunoblot. **B**, GST fusion proteins containing the full-length N terminus of AC8 or the full-length N terminus of AC8 with point mutations in the calmodulin binding region (8M34) immobilized on glutathione-Sepharose beads were incubated with equivalent volumes of GST precleared brain membranes. The affinity precipitates were resolved by SDS-PAGE and immunoblotted with an antibody raised against the catalytic subunit of protein phosphatase 2A (anti- $\text{PP2A}_C$ ). The Coomassie-stained membrane is shown at the bottom. **C**, a GST fusion protein containing the full-length N terminus of AC8 with point mutations in the calmodulin binding region (8M34) immobilized on glutathione-Sepharose beads was incubated with GST precleared brain membranes containing either 1 mM EDTA/EGTA or  $20 \mu\text{M}$   $\text{Ca}^{2+}$  and  $0.5 \mu\text{M}$  calmodulin. The affinity precipitates were resolved by SDS-PAGE and were immunoblotted with an antibody raised against the catalytic subunit of protein phosphatase 2A (anti- $\text{PP2A}_C$ ). The Coomassie-stained membrane is shown at the bottom.

However, AC8 also localizes to cholesterol-rich membranes in HEK293 cells, and its regulation by CCE is dependent on the presence of membrane cholesterol.

This result clearly suggests that, as with nonexcitable cells, AC8 resides in lipid rafts in primary neurons. A component of cellular  $\text{PP2A}$  might be expected to occur in lipid rafts to permit interaction with the N terminus of AC8.  $\text{PP2A}$  is a highly abundant protein (Goldberg, 1999), and incubating hippocampal neurons with an antibody raised against  $\text{PP2A}_C$  demonstrated strong labeling throughout the entire neuronal structure, with no clear specific compartmentalization (data not shown). To identify whether  $\text{PP2A}_C$  was found in lipid rafts, a clearer method for the separation of cellular compartments was required. Lipid rafts resist extraction



**Fig. 6.** The calmodulin binding domains of AC8 are not regulated by phosphorylation. Cobalt affinity gel (control) or a histone-fusion protein containing the C terminus of AC8 immobilized on cobalt affinity gel (C2bAC8) was incubated in the presence or absence of  $20 \mu\text{M}$   $\text{Ca}^{2+}$  and  $0.5 \mu\text{M}$  calmodulin. The affinity precipitates were resolved by SDS-PAGE and immunoblotted with an antibody raised against calmodulin (anti-calmodulin). The Coomassie-stained membrane is shown at the bottom.



**Fig. 7.** AC8 and  $\text{PP2A}_C$  are present in lipid rafts. Hippocampal neurons at 10 DIV 48 h after transfection with GFP (**A**) or GFP-AC8 (**B**). **C**, hippocampal neurons at 10 DIV 48 h after transfection with GFP-AC8 were extracted with 0.5% Triton X-100 at  $4^\circ\text{C}$ . The scale bar in all images represents  $10 \mu\text{M}$ . **D**, crude mouse forebrain membranes were incubated in cold Triton X-100 at a 10:1 detergent-to-protein ratio and prepared for sucrose equilibrium density gradient centrifugation as described under *Materials and Methods*. Fractions (3–10) were resolved by SDS-PAGE and immunoblotted with antibodies raised against flotillin (anti-flotillin), PKA catalytic subunit (anti-PKA),  $\text{G}\alpha_{\text{s}}$ -olf (anti- $\text{G}\alpha_{\text{s}}$ ), or the catalytic subunit of protein phosphatase 2A (anti- $\text{PP2A}_C$ ).

with cold Triton X-100 and have a higher lipid-to-protein ratio, which permits their isolation because of their increased buoyancy in sucrose density gradients (Pike et al., 2002).

Mouse brain membranes were fractionated at an optimized detergent-to-protein ratio as described in *Materials and Methods*. Sodium carbonate was included in the extraction procedure to reduce the binding of raft-associated proteins to high-density Triton-insoluble material that pellets during sucrose centrifugation, thus retaining a greater fraction of genuine raft proteins in the buoyant fraction (Arni et al., 1998). Flotillin immunoreactivity was used to identify the lipid raft fraction (Lang et al., 1998). At low detergent-to-protein ratios, flotillin was distributed in both the raft and nonraft fractions, whereas at ratios of 20:1 and greater, flotillin immunoreactivity was only observed in the pellet, demonstrating complete membrane solubilization (data not shown) (Lang et al., 1998). At a 10:1 detergent-to-protein ratio, flotillin was clearly enriched in fraction 4, which corresponded to 20% sucrose (Fig. 7D).  $G\alpha_{\text{olf}}$  was also enriched in the light membrane fraction (Fig. 7D) (Rybin et al., 2000). Because both of these lipid raft markers were clearly enriched in the buoyant membrane fraction, this indicated a high degree of separation between lipid rafts and bulk membranes. Lipid rafts constitute only a small amount of total protein, with the bulk of the protein content residing in fractions 6 to 10 (approximately 80% of total protein; data not shown). The catalytic subunit of PKA was present in lipid rafts (fraction 4) and in the bulk membrane, corresponding to fraction 7 (~35% sucrose) (Fig. 7D) (Razani et al., 1999). This clear separation of lipid rafts allowed us to search for PP2A. Immunoblotting with an antibody raised against PP2A<sub>C</sub> demonstrated PP2A<sub>C</sub> in lipid rafts (Fig. 7D). However, as might be expected for a highly expressed protein with diverse regulatory subunits, PP2A<sub>C</sub> immunoreactivity was also present in the bulk membrane and particulate fractions. It is noteworthy, however, that a fraction of PP2A<sub>C</sub> occurs in an environment that would permit interaction with AC8.

## Discussion

When heterologously expressed in nonexcitable cells, AC8 is exclusively regulated by CCE (Smith et al., 2002). This observation, along with other data, has led to the suggestion that AC8 and CCE channels reside in close proximity at the plasma membrane (Cooper et al., 1994; Chiono et al., 1995; Fagan et al., 1996, 1998, 2000a; Murthy and Makhoul, 1998; Watson et al., 2000; Smith et al., 2002). The targeting of AC8 to specialized domains of the plasma membrane, enriched in cholesterol and sphingolipids, termed lipid rafts, is essential but not sufficient for enabling regulation by CCE (Smith et al., 2002). The molecular identity of CCE channels remains uncertain, although mammalian homologues of the *Drosophila melanogaster* transient receptor potential (TRP) proteins are putative candidates (Putney and McKay, 1999). Of the TRP proteins, TRP1 at least is enriched in lipid rafts (Lockwich et al., 2000). An N-terminally truncated form of AC8 is unresponsive to CCE regulation, although the enzyme remains localized in lipid rafts and is fully stimulated by  $\text{Ca}^{2+}$ /calmodulin in vitro (Smith et al., 2002). This led us to consider that the N terminus of AC8 might associate with elements of the cellular CCE apparatus.

Therefore, in this study, we used the N terminus of AC8 in

a yeast two-hybrid screen of an HEK293 cDNA library to try to identify interacting proteins that might contribute to CCE regulation of AC8. It turned out that none of the positive candidates identified were proteins that might obviously be involved in regulating CCE. However, the catalytic subunit of PP2A emerged as one compelling interacting protein. The failure to find more obvious candidates in this screen does not exclude the possibility that the N terminus targets AC8 directly to elements of the CCE apparatus, although it suggests that more indirect interactions should be countenanced.

Extending and confirming the interaction identified by the yeast two-hybrid screen, an N-terminal AC8 GST-fusion protein affinity precipitated the full-length PP2A<sub>C</sub> from both brain—the natural source of AC8—and HEK293 membranes. This underlined the highly specific nature of the interaction between PP2A<sub>C</sub> and the N terminus of AC8 against a large background of other proteins.

PP2A is one of the four major types of serine/threonine protein phosphatase and associates in vivo with a scaffolding subunit of 65 kDa (PP2A<sub>A</sub>). This core dimer can further associate with one of a class of regulatory B subunits, ranging in molecular mass from 55 to 130 kDa. The B subunits apparently play a role in specifying substrate selection and subcellular location (Goldberg, 1999; Janssens and Goris, 2001). We identified PP2A<sub>A</sub> in pull-downs that were positive for PP2A<sub>C</sub>, which suggested that the N terminus of AC8 interacted with the PP2A core dimer. However the possibility cannot be ruled out that PP2A<sub>A</sub> and PP2A<sub>C</sub> interact at individual sites on the N terminus of AC8. We did not detect any B subunits when immunoblots were probed with a pan-specific PP2A<sub>B</sub> antibody. This may indicate that the core dimer preferentially associates with the N terminus of AC8 or that PP2A<sub>B</sub> was not present in sufficient quantity to allow signal detection by enhanced chemiluminescence on immunoblotting with a pan-PP2A<sub>B</sub> antibody.

Although the regulation of PP2A is believed to occur through the A and B subunits, additional proteins that bind specifically to the PP2A catalytic subunit have been identified. Axin, a component of the Wnt signal transduction system, PKC- $\alpha$ , the  $\alpha 1_c$  subunit of the L-type VGCC, and the NR3A subunit of the N-methyl-D-aspartate receptor all display a specific interaction with PP2A<sub>C</sub> (Hsu et al., 1999; Davare et al., 2000; Chan and Sucher, 2001; Boudreau et al., 2002). Such observations suggest that the catalytic subunit may itself confer a degree of regulation to PP2A activity. Among proteins shown to bind to PP2A<sub>C</sub>, there seems to be no immediately obvious motif responsible for PP2A<sub>C</sub> association. Indeed, it is possible that PP2A<sub>C</sub> associates with low stringency to its respective binding sites (Ma and Sucher, 2004).

To begin exploring the putative physiological meaning of the interaction between PP2A<sub>C</sub> and the N terminus of AC8, we investigated whether PP2A<sub>C</sub> was catalytically active. Indeed, the fraction of PP2A<sub>C</sub> that associated with the N terminus of AC8 dephosphorylated myelin basic protein that had been phosphorylated by PKA. This finding of an interaction between AC8 and catalytically active PP2A<sub>C</sub> opens a very interesting possibility and a means of drawing together previous speculations that AC, VGCCs, PKA,  $\beta_2$ -adrenergic receptors, and PP2A might form a regulatory complex (Davare et al., 2001).

$\text{Ca}^{2+}$ /calmodulin binding to the single  $\alpha$  helical domain in



the N terminus of AC8 is an essential step in the activation of AC8 by increases in  $[Ca^{2+}]_i$  and is considered to occur on a 1:1 ratio (Gu and Cooper, 1999; Smith et al., 2002). We found that the interaction of the AC8 N terminus with PP2A<sub>C</sub> was prevented by  $Ca^{2+}$ /calmodulin, which suggests that PP2A also interacted with the N terminus of AC8 on a 1:1 ratio. This competition between  $Ca^{2+}$ /calmodulin and PP2A<sub>C</sub> for the N terminus of AC8 is comparable with the binding of PP2A to the autoregulatory domain of CaMKIV, which also occurs in a mutually exclusive manner with respect to  $Ca^{2+}$ /calmodulin (Anderson et al., 2004). However, there is no sequence homology between the autoregulatory domains of CaMKIV and the N terminus of AC8. Although binding of PP2A<sub>C</sub> to the N terminus of AC8 was prevented by  $Ca^{2+}$ /calmodulin, PP2A<sub>C</sub> was not binding to the identical amino acid sequence that bound  $Ca^{2+}$ /calmodulin, because mutations of amino acids within the  $\alpha$  helical domain that were essential for  $Ca^{2+}$ /calmodulin association were not required for binding of PP2A<sub>C</sub>. It is therefore likely that the respective domains overlap.

The association of calmodulin with its target sequence in proteins can often be regulated by dynamic phosphorylation of residues within or adjacent to the calmodulin binding domain (Black et al., 2004). When we explored the functional consequence of catalytically active PP2A association with AC8 in the context of regulating phosphorylation of the N-terminal or C2b calmodulin binding domains, we found that neither domain was phosphorylated by PKA or a forskolin- and prostaglandin  $E_1$ -induced HEK293 cell lysate. This suggests that PKA-mediated phosphorylation is not involved in regulating calmodulin binding to AC8. However, this does not rule out the possibility that other sites on AC8 are regulated by phosphorylation, either by PKA or additional kinases. Therefore, PP2A, in association with AC8, if not involved in the regulation of calmodulin binding to AC8, may be involved in regulating the phosphorylation status of proteins in the vicinity of AC8. For example, the association of PP2A with NR3A directly regulates the phosphorylation state, not of NR3A, but of the adjacent NR1 subunit (Chan and Sucher, 2001). In this context, both AC8 and PP2A<sub>C</sub> are present in lipid raft microdomains in which PP2A<sub>C</sub> is positioned in an environment with other signaling molecules, including putative CCE channels and voltage-gated calcium channels (Fagan et al., 2000a,b).

The association of PP2A<sub>C</sub> with the NR3A subunit of the N-methyl-D-aspartate receptor is disrupted by  $Ca^{2+}$  entry through the receptor ion channel; however, the NR3A subunit has not been shown to bind calmodulin (Chan and Sucher, 2001). It is conceivable that a similar scenario occurs in relation to the association of the N terminus of AC8 with PP2A. AC8 selectively responds to increases in  $[Ca^{2+}]_i$  through either CCE channels or L-type VGCC. These modes of  $Ca^{2+}$  entry are believed to give rise to microdomains of elevated  $[Ca^{2+}]_i$  in the vicinity of the  $Ca^{2+}$ -sensitive AC, an essential element for regulation within intact cells. Thus the high level of  $[Ca^{2+}]_i$  at these sites would be expected to recruit calmodulin, which in turn may displace catalytically active PP2A<sub>C</sub> from the N terminus of AC8, facilitating activation of the cyclase and the dephosphorylation of target proteins (Persechini and Cronk, 1999).

This study failed to detect any interaction between the N terminus of AC8 and putative CCE channel proteins, despite

the fact that the N terminus is an essential component in enabling AC8 to respond to increases in  $[Ca^{2+}]_i$  in vivo. Thus, it would seem that the N terminus of AC8 is not responsible for a direct association with CCE channels. However, the coimmunoprecipitation of an unidentified adenylyl cyclase isoform with an L-type VGCC has been described in rat forebrain membranes, which suggests a close association between endogenous  $Ca^{2+}$ -sensitive adenylyl cyclases and L-type VGCCs. This latter interaction formed part of a larger signaling complex, in which PP2A<sub>C</sub>, the  $\beta_2$ -adrenergic receptor, PKA, and  $G\alpha_s$  were all identified (Davare et al., 2000, 2001). Thus, it is conceivable that the interaction of catalytically active PP2A<sub>C</sub> with AC8 may function as part of a larger signaling complex coordinating increases in  $[Ca^{2+}]_i$  with the generation of cAMP, with concomitant alterations in the phosphorylation levels of key signaling intermediates by either  $Ca^{2+}$ - or cAMP-activated protein kinases. Despite the number of potential PP2A substrates, the in vivo regulation of PP2A by extracellular signals is not well understood; indeed, its regulation may rely on protein-protein interactions that position PP2A at sites of required activity (Hsu et al., 1999; Sim and Scott, 1999).

AC8 links  $[Ca^{2+}]_i$  increases to elevations in cAMP and, as such, is intimately involved in hippocampal long-term potentiation (Wang et al., 2003). This first demonstration of the association of an adenylyl cyclase with an active protein phosphatase provides a key intermediate in the organization of a dynamic signaling network. This may be particularly important in neuronal contexts involving  $Ca^{2+}$ /calmodulin, L-type VGCC, PKA, AC8, and PP2A, in which such signaling complexes may play an important role in regulating synaptic plasticity (Frey et al., 1993; Wang et al., 2003). Future experimental exploration of the physiological significance of the AC8/PP2A interaction described may be profitably considered in such neuronal contexts, in which AC8 naturally occurs. Fluorescence resonance energy transfer studies involving cyan fluorescent protein- and yellow fluorescent protein-labeled components could provide a dynamic means for studying the interactions of the full-length proteins in living cells.

#### Acknowledgments

We thank Dr. Brian Wadzinski for the PP2A<sub>A</sub> and pan-PP2A<sub>B</sub> polyclonal antibodies, Dr. Sangeeta Chawla and Johanna Belfield for culture and transfection of hippocampal neurons, and Dr. Debbie Willoughby for assistance with confocal imaging and useful comments on the manuscript.

#### References

- Anderson KA, Noeldner PK, Reece K, Wadzinski BE, and Means AR (2004) Regulation and function of the calcium/calmodulin-dependent protein kinase IV/protein serine/threonine phosphatase 2A signaling complex. *J Biol Chem* **279**:31708–31716.
- Arni S, Keilbaugh SA, Ostermeyer AG, and Brown DA (1998) Association of GAP-43 with detergent-resistant membranes requires two palmitoylated cysteine residues. *J Biol Chem* **273**:28478–28485.
- Barford D, Das AK, and Egloff MP (1998) The structure and mechanism of protein phosphatases: insights into catalysis and regulation. *Annu Rev Biophys Biomol Struct* **27**:133–164.
- Black DJ, Tran QK, and Persechini A (2004) Monitoring the total available calmodulin concentration in intact cells over the physiological range in free  $Ca^{2+}$ . *Cell Calcium* **35**:415–425.
- Boudreau RT, Garduno R, and Lin TJ (2002) Protein phosphatase 2A and protein kinase C $\alpha$  are physically associated and are involved in *Pseudomonas aeruginosa*-induced interleukin 6 production by mast cells. *J Biol Chem* **277**:5322–5329.
- Cali JJ, Zwaagstra JC, Mons N, Cooper DMF, and Krupinski J (1994) Type VIII adenylyl cyclase. A  $Ca^{2+}$ /calmodulin-stimulated enzyme expressed in discrete regions of rat brain. *J Biol Chem* **269**:12190–12195.

- Chan SF and Sucher NJ (2001) An NMDA receptor signaling complex with protein phosphatase 2A. *J Neurosci* **21**:7985–7992.
- Chawla S, Vanhoutte P, Arnold FJ, Huang CL, and Bading H (2003) Neuronal activity-dependent nucleocytoplasmic shuttling of HDAC4 and HDAC5. *J Neurochem* **85**:151–159.
- Chiono M, Mahey R, Tate G, and Cooper DMF (1995) Capacitative  $\text{Ca}^{2+}$  entry exclusively inhibits cAMP synthesis in C6–2B glioma cells. Evidence that physiologically evoked  $\text{Ca}^{2+}$  entry regulates  $\text{Ca}^{2+}$ -inhibitable adenylyl cyclase in non-excitable cells. *J Biol Chem* **270**:1149–1155.
- Chou JL, Huang CL, Lai HL, Hung AC, Chien CL, Kao YY, and Chern Y (2004) Regulation of type VI adenylyl cyclase by Snapin, a SNAP25-binding protein. *J Biol Chem* **279**:46271–46279.
- Cooper DMF (2003) Regulation and organization of adenylyl cyclases and cAMP. *Biochem J* **375**:517–529.
- Cooper DMF, Yoshimura M, Zhang Y, Chiono M, and Mahey R (1994) Capacitative  $\text{Ca}^{2+}$  entry regulates  $\text{Ca}^{2+}$ -sensitive adenylyl cyclases. *Biochem J* **297**:437–440.
- Davare MA, Avdonin V, Hall DD, Peden EM, Burette A, Weinberg RJ, Horne MC, Hoshi T, and Hell JW (2001) A beta2 adrenergic receptor signaling complex assembled with the  $\text{Ca}^{2+}$  channel Cav1.2. *Science (Wash DC)* **293**:98–101.
- Davare MA, Horne MC, and Hell JW (2000) Protein phosphatase 2A is associated with class C L-type calcium channels (Cav1.2) and antagonizes channel phosphorylation by cAMP-dependent protein kinase. *J Biol Chem* **275**:39710–39717.
- Egloff MP, Johnson DF, Moorhead G, Cohen PT, Cohen P, and Barford D (1997) Structural basis for the recognition of regulatory subunits by the catalytic subunit of protein phosphatase 1. *EMBO (Eur Mol Biol Organ) J* **16**:1876–1887.
- Enyedi A, Elwess NL, Filoteo AG, Verma AK, Paszty K, and Penniston JT (1997) Protein kinase C phosphorylates the “a” forms of plasma membrane  $\text{Ca}^{2+}$  pump isoforms 2 and 3 and prevents binding of calmodulin. *J Biol Chem* **272**:27525–27528.
- Fagan KA, Graf RA, Tolman S, Schaack J, and Cooper DMF (2000a) Regulation of a  $\text{Ca}^{2+}$ -sensitive adenylyl cyclase in an excitable cell. Role of voltage-gated versus capacitative  $\text{Ca}^{2+}$  entry. *J Biol Chem* **275**:40187–40194.
- Fagan KA, Mahey R, and Cooper DMF (1996) Functional co-localization of transfected  $\text{Ca}^{2+}$ -stimulable adenylyl cyclases with capacitative  $\text{Ca}^{2+}$  entry sites. *J Biol Chem* **271**:12438–12444.
- Fagan KA, Mons N, and Cooper DMF (1998) Dependence of the  $\text{Ca}^{2+}$ -inhibitable adenylyl cyclase of C6–2B glioma cells on capacitative  $\text{Ca}^{2+}$  entry. *J Biol Chem* **273**:9297–9305.
- Fagan KA, Smith KE, and Cooper DMF (2000b) Regulation of the  $\text{Ca}^{2+}$ -inhibitable adenylyl cyclase type VI by capacitative  $\text{Ca}^{2+}$  entry requires localization in cholesterol-rich domains. *J Biol Chem* **275**:26530–26537.
- Fleming I, Fisslthaler B, Dimmeler S, Kemp BE, and Busse R (2001) Phosphorylation of Thr<sub>95</sub> regulates  $\text{Ca}^{2+}$ /calmodulin-dependent endothelial nitric oxide synthase activity. *Circ Res* **88**:E68–E75.
- Foster LJ, De Hoog CL, and Mann M (2003) Unbiased quantitative proteomics of lipid rafts reveals high specificity for signaling factors. *Proc Natl Acad Sci USA* **100**:5813–5818.
- Frey U, Huang YY, and Kandel ER (1993) Effects of cAMP simulate a late stage of LTP in hippocampal CA1 neurons. *Science (Wash DC)* **260**:1661–1664.
- Goldberg Y (1999) Protein phosphatase 2A: who shall regulate the regulator? *Biochem Pharmacol* **57**:321–328.
- Greif DM, Kou R, and Michel T (2002) Site-specific dephosphorylation of endothelial nitric oxide synthase by protein phosphatase 2A: evidence for crosstalk between phosphorylation sites. *Biochemistry* **41**:15845–15853.
- Gu C and Cooper DMF (1999) Calmodulin-binding sites on adenylyl cyclase type VIII. *J Biol Chem* **274**:8012–8021.
- Hering H, Lin CC, and Sheng M (2003) Lipid rafts in the maintenance of synapses, dendritic spines and surface AMPA receptor stability. *J Neurosci* **23**:3262–3271.
- Hofmann F, Anagli J, Carafoli E, and Vorherr T (1994) Phosphorylation of the calmodulin binding domain of the plasma membrane  $\text{Ca}^{2+}$  pump by protein kinase C reduces its interaction with calmodulin and with its pump receptor site. *J Biol Chem* **269**:24298–24303.
- Hsu W, Zeng L, and Costantini F (1999) Identification of a domain of Axin that binds to the serine/threonine protein phosphatase 2A and a self-binding domain. *J Biol Chem* **274**:3439–3445.
- Janssens V and Goris J (2001) Protein phosphatase 2A: a highly regulated family of serine/threonine phosphatases implicated in cell growth and signalling. *Biochem J* **353**:417–439.
- Kremmer E, Ohst K, Kiefer J, Brewis N, and Walter G (1997) Separation of PP2A core enzyme and holoenzyme with monoclonal antibodies against the regulatory A subunit: abundant expression of both forms in cells. *Mol Cell Biol* **17**:1692–1701.
- Lang DM, Lommel S, Jung M, Ankerhold R, Petrusch B, Laessing U, Wiechers MF, Plattner H, and Stuermer CA (1998) Identification of reggie-1 and reggie-2 as plasma membrane-associated proteins which cocluster with activated GPI-anchored cell adhesion molecules in non-caveolar micropatches in neurons. *J Neurobiol* **37**:502–523.
- Lau LF, Mammen A, Ehlers MD, Kindler S, Chung WJ, Garner CC, and Haganir RL (1996) Interaction of the N-methyl-D-aspartate receptor complex with a novel synapse-associated protein, SAP102. *J Biol Chem* **271**:21622–21628.
- Lavigne N, Ethier N, Oak JN, Pei L, Liu F, Trieu P, Rebois RV, Bouvier M, Hebert TE, and Van Tol HH (2002) G protein-coupled receptors form stable complexes with inwardly rectifying potassium channels and adenylyl cyclase. *J Biol Chem* **277**:46010–46019.
- Lebrin F, Bianchini L, Rabilloud T, Chambaz EM, and Goldberg Y (1999) CK2alpha-protein phosphatase 2A molecular complex: possible interaction with the MAP kinase pathway. *Mol Cell Biochem* **191**:207–212.
- Leonard AS, Davare MA, Horne MC, Garner CC, and Hell JW (1998) SAP97 is associated with the  $\alpha$ -amino-3-hydroxy-5-methylisoxazole-4-propionic acid receptor GluR1 subunit. *J Biol Chem* **273**:19518–19524.
- Lockwich TP, Liu X, Singh BB, Jadowiec J, Weiland S, and Ambudkar IS (2000) Assembly of Trp1 in a signaling complex associated with caveolin-scaffolding lipid raft domains. *J Biol Chem* **275**:11934–11942.
- Ma OK and Sucher NJ (2004) Molecular interaction of NMDA receptor subunit NR3A with protein phosphatase 2A. *Neuroreport* **15**:1447–1450.
- Mehta S, Wu H, Garner CC, and Marshall J (2001) Molecular mechanisms regulating the differential association of kainate receptor subunits with SAP90/PSD-95 and SAP97. *J Biol Chem* **276**:16092–16099.
- Mons N, Harry A, Dubourg P, Premont RT, Iyengar R, and Cooper DMF (1995) Immunohistochemical localization of adenylyl cyclase in rat brain indicates a highly selective concentration at synapses. *Proc Natl Acad Sci USA* **92**:8473–8477.
- Mons N, Yoshimura M, Ikeda H, Hoffman PL, and Tabakoff B (1998) Immunological assessment of the distribution of type VII adenylyl cyclase in brain. *Brain Res* **788**:251–261.
- Murthy KS and Makhlof GM (1998) Regulation of adenylyl cyclase type V/VI in smooth muscle: interplay of inhibitory G protein and  $\text{Ca}^{2+}$  influx. *Mol Pharmacol* **54**:122–128.
- Ostermeyer AG, Beckrich BT, Ivarson KA, Grove KE, and Brown DA (1999) Glycosphingolipids are not essential for formation of detergent-resistant membrane rafts in melanoma cells. Methyl- $\beta$ -cyclodextrin does not affect cell surface transport of a GPI-anchored protein. *J Biol Chem* **274**:34459–34466.
- Persechini A and Cronk B (1999) The relationship between the free concentrations of  $\text{Ca}^{2+}$  and  $\text{Ca}^{2+}$ -calmodulin in intact cells. *J Biol Chem* **274**:6827–6830.
- Pike LJ, Han X, Chung KN, and Gross RW (2002) Lipid rafts are enriched in arachidonic acid and plasmenylethanolamine and their composition is independent of caveolin-1 expression: a quantitative electrospray ionization/mass spectrometric analysis. *Biochemistry* **41**:2075–2088.
- Poser S and Storm DR (2001) Role of  $\text{Ca}^{2+}$ -stimulated adenylyl cyclases in LTP and memory formation. *Int J Dev Neurosci* **19**:387–394.
- Putney JW and McKay RR (1999) Capacitative calcium entry channels. *Bioessays* **21**:38–46.
- Razani B, Rubin CS, and Lisanti MP (1999) Regulation of cAMP-mediated signal transduction via interaction of caveolins with the catalytic subunit of protein kinase A. *J Biol Chem* **274**:26353–26360.
- Rybin VO, Xu X, Lisanti MP, and Steinberg SF (2000) Differential targeting of beta-adrenergic receptor subtypes and adenylyl cyclase to cardiomyocyte caveolae. A mechanism to functionally regulate the cAMP signaling pathway. *J Biol Chem* **275**:41447–41457.
- Sim AT and Scott JD (1999) Targeting of PKA, PKC and protein phosphatases to cellular microdomains. *Cell Calcium* **26**:209–217.
- Smith FD and Scott JD (2002) Signaling complexes: junctions on the intracellular information super highway. *Curr Biol* **12**:R32–R40.
- Smith KE, Gu C, Fagan KA, Hu B, and Cooper DMF (2002) Residence of adenylyl cyclase type 8 in caveolae is necessary but not sufficient for regulation by capacitative  $\text{Ca}^{2+}$  entry. *J Biol Chem* **277**:6025–6031.
- Turner JH, Gelasco AK, and Raymond JR (2004) Calmodulin interacts with the third intracellular loop of the serotonin 5-hydroxytryptamine1A receptor at two distinct sites: putative role in receptor phosphorylation by protein kinase C. *J Biol Chem* **279**:17027–17037.
- Wang H, Pineda VV, Chan GC, Wong ST, Muglia LJ, and Storm DR (2003) Type 8 adenylyl cyclase is targeted to excitatory synapses and required for mossy fiber long-term potentiation. *J Neurosci* **23**:9710–9718.
- Watson EL, Jacobson KL, Singh JC, Idzerda R, Ott SM, DiJulio DH, Wong ST, and Storm DR (2000) The type 8 adenylyl cyclase is critical for  $\text{Ca}^{2+}$  stimulation of cAMP accumulation in mouse parotid acini. *J Biol Chem* **275**:14691–14699.
- Williams R and Coluccio LM (1995) Phosphorylation of myosin-I from rat liver by protein kinase C reduces calmodulin binding. *Biochem Biophys Res Commun* **216**:90–102.

**Address correspondence to:** Dr. Dermot M. F. Cooper, Department of Pharmacology, University of Cambridge, Tennis Court Rd, Cambridge, CB2 1PD, United Kingdom. E-mail: dmfc2@cam.ac.uk