

Identification of Key Residues Coordinating Functional Inhibition of P2X₇ Receptors by Zinc and Copper^S

Xing Liu, Annmarie Surprenant, Hong-Ju Mao, Sebastien Roger, Rong Xia, Helen Bradley, and Lin-Hua Jiang

Institute of Membrane and Systems Biology, Faculty of Biological Sciences, University of Leeds, Leeds, United Kingdom (X.L., H.-J.M., R.X., H.B., L.-H.J.); and Department of Biomedical Science, University of Sheffield, United Kingdom (A.S., S.R.)

Received July 4, 2007; accepted October 23, 2007

ABSTRACT

P2X₇ receptors are distinct from other ATP-gated P2X receptors in that they are potently inhibited by submicromolar concentrations of zinc and copper. The molecular basis for the strong functional inhibition by zinc and copper at this purinergic ionotropic receptor is controversial. We hypothesized that it involves a direct interaction of zinc and copper with residues in the ectodomain of the P2X₇ receptor. Fourteen potential metal interacting residues are conserved in the ectodomain of all mammalian P2X₇ receptors, none of which is homologous to previously identified sites in other P2X receptors shown to be important for functional potentiation by zinc. We introduced alanine substitutions into each of these residues, expressed

wild-type and mutated receptors in human embryonic kidney 293 cells, and recorded resulting ATP and BzATP-evoked membrane currents. Agonist concentration-response curves were similar for all 12 functional mutant receptors. Alanine substitution at His⁶² or Asp¹⁹⁷ strongly attenuated both zinc and copper inhibition, and the double mutant [H62A/D197A] mutant receptor was virtually insensitive to inhibition by zinc or copper. Thus, we conclude that zinc and copper inhibition is due to a direct interaction of these divalent cations with ectodomain residues of the P2X₇ receptor, primarily involving combined interaction with His⁶² and Asp¹⁹⁷ residues.

The P2X₇ receptor (previously termed P2Z) is the last member (Surprenant et al., 1996) of the ATP-gated P2X receptor family and is found primarily on hematopoietic lineage cells, such as lymphocytes, monocytes, and macrophages, as well as brain microglial cells and astrocytes (North, 2002; Duan and Neary, 2006). Studies using different approaches, including gene knockout, have underscored P2X₇ receptors as a crucial component in a diversity of biological processes, such as inflammation, elimination of intracellular pathogens, release of pro-inflammatory cytokines by macrophages, and release of neurotransmitters by astrocytes, and chronic inflammatory and neuropathic pain (e.g., Lammas et al., 1997; Solle et al., 2001; Chessell et al., 2005; Duan and Neary, 2006; Ferrari et al., 2006).

P2X₇ receptors show several unique functional features that differ strikingly from those of other members of the P2X

receptor family (North, 2002). P2X₇ receptors display remarkable functional plasticity; brief activation opens small cation permeable channels, whereas prolonged stimulation induces formation of “large molecule” permissive pores, rapid membrane and mitochondrial morphological changes, cytoskeletal rearrangement, and eventual cell death (Surprenant et al., 1996; North, 2002; Morelli et al., 2003). The most upstream signaling event consequent to the P2X₇ receptor activation, the ionic current, also shows properties uniquely different from those of other P2X receptors. Currents are activated by submillimolar concentrations of ATP, which are 10 to 100 times higher than those required for activation of other P2X receptors. Furthermore, P2X₇ receptors are potently inhibited by submicromolar concentrations of zinc and copper, whereas other P2X receptors are strongly potentiated or unaffected (Wildman et al., 1998, 1999a,b; Xiong et al., 1999; Acuña-Castillo et al., 2000; Clyne et al., 2002; Lorca et al., 2005). P2X₇ receptors are also substantially inhibited by magnesium and calcium, albeit with much less potency than zinc and copper. Earlier studies examining the effects of calcium and magnesium have led to the view that ATP⁴⁻ is the effective agonist form of ATP and these divalent cations inhibit the P2X₇ receptors by reducing the effective agonist concentrations (Cockcroft and Gomperts,

This work was supported by the Biotechnology and Biological Sciences Research Council (BBSRC) (L.-H.J., A.S.), the Royal Society (H.-J. M., L.-H.J.), a UK Overseas Research Scholarship (R.X.), and a BBSRC doctoral training grant (H.B.).

Article, publication date, and citation information can be found at <http://molpharm.aspetjournals.org>.
doi:10.1124/mol.107.039651.

^S The online version of this article (available at <http://molpharm.aspetjournals.org>) contains supplemental material.

ABBREVIATIONS: BzATP, 2',3'-O-(4-benzoylbenzoyl) adenosine 5'-triphosphate; HEK, human embryonic kidney.

1979). This view is still prevailing and implied in the inhibition by other divalent cations such as zinc and copper (North, 2002). Our previous indirect evidence demonstrates that the divalent cation that induces inhibition of P2X₇ currents is due primarily to direct binding to and subsequent allosteric modulation of the P2X₇ receptor (Virginio et al., 1997).

Histidine, cysteine, lysine, aspartic acid, and glutamic acid residues are all known to be capable of coordinating zinc inhibition/potential of different ion channels, such as GABA_C receptors (Wang et al., 1995), glycine receptors (Laube et al., 2000), *N*-methyl-D-aspartate receptors (Paoletti et al., 2000), and acid-sensing ion channels (Chu et al., 2004). Sequence analysis of the ectodomain of the seven P2X receptor subunits led us to the identification of 14 potential zinc interacting residues that are conserved in mouse, rat, and human P2X₇ receptors but not the other P2X receptor family members. We introduced alanine substitution by site-directed mutagenesis into these 14 positions in the rat P2X₇ receptor and found that mutation of His⁶² or Asp¹⁹⁷ residues strongly attenuated the inhibition by zinc and copper. Thus, the present study provides direct evidence supporting the view that the potent functional inhibition of the P2X₇ receptor by zinc and copper results primarily from an interaction with the receptor, in which His⁶² and Asp¹⁹⁷ residues are critical.

Materials and Methods

Cell and Molecular Biology. Human embryonic kidney (HEK293) cells were cultured and transfected with plasmids as described previously (Jiang et al., 2001). Wild-type and mutant rat P2X₇ receptors were engineered to contain a C-terminal EYMPME epitope (EE tag). Tagging had no detectable effect on the functional properties of wild-type P2X₇ receptors (L.-H. Jiang and A. Surprenant, unpublished data). Alanine substitution was introduced by site-directed mutagenesis and confirmed by sequencing.

Electrophysiological Recording. Whole-cell recordings were carried out using an Axopatch 200B (Molecular Devices, Sunnyvale, CA) or EPC10 (HEKA, Lambrecht/Pfalz, Germany) amplifier at room temperature as described previously (Jiang et al., 2001). Cells were held at -60 mV. Patch pipettes were fabricated from borosilicate glass capillaries with a resistance of 4 to 6 MΩ. Standard external solution contained 147 mM NaCl, 2 mM KCl, 1 mM MgCl₂, 2 mM CaCl₂, 10 mM HEPES, and 13 mM glucose. Intracellular solution contained 145 mM NaCl, 10 mM EGTA, and 10 mM HEPES. Copper chloride and zinc sulfate salts were dissolved in standard external solution. Application of the agonists, zinc and copper was carried out via a rapid solution changer RSC-160 system (Bio-Logic Science Instruments, Claix, France), which has limitations in studying very fast kinetics (Spelta et al., 2002), possibly contributing to the variable rebound currents. All the chemicals were purchased commercially (Sigma, St. Louis, MO).

Immunocytochemistry. Transfected cells were stained as described previously (Jiang et al., 2001) using mouse primary anti-EE antibody at a dilution of 1:1000 (BAbCO, Richmond, CA) and secondary fluorescein isothiocyanate-conjugated anti-mouse IgG antibody at a dilution of 1:200 (Sigma), or using rabbit anti-EE antibody at a dilution of 1:1000 (Bethyl Laboratories, Montgomery, TX) and secondary Alexa Fluor 488 anti-rabbit IgG antibody at a dilution of 1:2000 (Invitrogen, Carlsbad, CA). The images were captured using a Zeiss AxioVert 200M confocal microscope and LSM510 META software (Carl Zeiss GmbH, Jena, Germany).

Data Analysis. All data, where appropriate, are presented as mean ± S.E.M. The EC₅₀ values for the agonists BzATP and ATP were determined by least-squares curve fitting to the Hill equation:

$I/I_{\max} = 1/[1 + (EC_{50}/[A])^{n_H}]$, where I is the current as a fraction of the maximum (I_{\max}), $[A]$ is the agonist concentration, and n_H is the Hill coefficient. Likewise, the IC₅₀ values for the inhibitors zinc and copper were derived by fitting to $I/I_0 = 1/[1 + ([B]/IC_{50})^{n_H}]$, where I is the current as a fraction of the control current (I_0) in the absence of the inhibitors under examination and $[B]$ is the inhibitor concentration. It is noteworthy that in the present study (except Fig. 1C), zinc or copper was coapplied with agonists for 4 s, and in most experiments, the inhibition occurred quickly and reached steady state by the end of the 4-s application. Figures show the curves fitted to the mean of all experiments. Curve fit was done using Origin software (OriginLab Corp, Northampton, MA). Statistical analysis was carried out using Student's t test.

Results

Zinc Inhibition of BzATP-Evoked Currents at the Wild-Type P2X₇ Receptor. We used BzATP at 30 μM, which evoked 30 to 50% of maximum current, and three




Fig. 1. Inhibition of P2X₇ receptor currents by zinc and rebound currents. A and B, representative BzATP-evoked currents in the presence of increasing concentrations of zinc. Note that there were robust rebound currents accompanying simultaneous washout of zinc and BzATP, as indicated by the arrows (the first protocol, A), which were absent in washout of BzATP in the persistent presence of zinc (the second protocol, B). C, zinc concentration-current response curves from all protocols as shown in A and B and described in text ($n = 3-8$ for each point). ●, data from the first protocol ($IC_{50} = 4.6 \pm 0.9 \mu\text{M}$, $n = 8$); ■, data from the second protocol ($IC_{50} = 5.9 \pm 1.2 \mu\text{M}$, $n = 4$); ▲, data from preapplication of zinc ($IC_{50} = 5.6 \pm 1.4 \mu\text{M}$, $n = 4$). These values were not significantly different from each other.

distinct protocols to examine zinc inhibition of the BzATP-evoked currents: simultaneous application and washout of agonist and zinc (Fig. 1A), simultaneous application of agonist/zinc with sustained presence of zinc upon washout of BzATP (Fig. 1B), or preapplication of zinc for 30 to 90 s followed by either of the former protocols (Fig. 1C). The zinc inhibition curves were not significantly different using any of these protocols ($IC_{50} = 4.6 \pm 0.9 \mu\text{M}$, $n = 8$; $5.9 \pm 1.2 \mu\text{M}$, $n = 4$, and $5.6 \pm 1.4 \mu\text{M}$, $n = 4$, respectively). It is noteworthy that we observed rebound currents upon simultaneous washout of agonist and zinc irrespective of the duration of preapplication of zinc (Fig. 1A). These rebound currents were particularly prominent at high concentrations of zinc but were never observed upon washing agonist in the persistent presence of zinc (Fig. 1B). No quantitative characterization of the rebound currents was attempted because there was significant variation in their amplitude that was probably due to the limited solution exchange rate of the system (see *Materials and Methods*). However, the presence of rebound currents upon simultaneous agonist/antagonist washout suggests that they result from the faster dissociation of zinc from its binding site than that of BzATP from its binding site. This further supports our hypothesis that zinc interacts directly with sites on the receptor protein rather than acting by chelating free ATP^{4-} . We thus turned to site-directed mutagenesis studies to seek direct evidence to corroborate this view.

Effects of Alanine Substitution on Functional Expression. Sequence analysis of the ectodomains of the seven P2X receptor subunits identified 14 potential interacting residues (five histidines, three glutamic acids, two aspartic acids, and four lysines) uniquely located in the ectodomain of the P2X₇ receptor but not P2X₁₋₆ receptors (Supplemental Fig. 1). This analysis was based on the discriminating actions at the P2X₇ receptor versus other P2X receptors and the known roles of histidine, cysteine, lysine, aspartic acid, and glutamic acid in zinc-mediated inhibition/potential of several ion channels (see Introduction). We substituted the individual residues with alanine, expressed the resultant mutant receptors in HEK293 cells, and measured the BzATP-evoked currents by patch clamp recording. The current amplitudes to maximum concentration of BzATP were not significantly different from wild-type currents at 11 of the mutant receptors (Table 1). Either no currents or minimal (< 50 pA) currents were recorded at the P2X₇ mutant receptors [K54A] and [D156A], and immunostaining using an anti-EE epitope Ab showed significantly reduced membrane localization (Supplemental Fig. 2). Therefore, these mutant receptors were not further investigated. The [H201A] mutant receptor showed significantly lower maximum current amplitudes (4-fold lower) without a significant shift in the agonist EC_{50} (Table 1). Of the 12 functional mutant receptors examined in this study, only one, [K145A], showed a significantly different BzATP EC_{50} value, which was shifted to the right by approximately 4-fold (Table 1). These results suggest that alanine substitution at all sites introduced minimal global conformational changes.

Effects of Zinc. First, we examined the wild-type and 12 functional mutant channel currents elicited by 30 μM BzATP (producing ~20–50% of the maximum current as shown in Supplemental Fig. 3, except [K145A] at which 200 μM BzATP was used). We used this concentration of BzATP to

minimize the potential functional regulation by endogenous P2Y receptors (see *Discussion*). Furthermore, we used 30 μM zinc (resulting in nearly maximum inhibition) in the presence of BzATP to ask which residues, upon substitution with alanine, would give rise to pronounced reduction or loss of zinc inhibition. No significant differences in zinc inhibition of the BzATP-evoked currents were observed at 10 of the 12 mutant receptors examined (Fig. 2); however, we noted that zinc inhibition at the [H201A] mutant receptor showed much greater variability compared with the other receptors (Fig. 2B). Upon simultaneous washout of BzATP and zinc, rebound currents continued to be present at all of these mutant receptors and were not qualitatively different from those observed for the wild-type receptor. In contrast, zinc inhibition of BzATP-evoked currents was strongly attenuated (by 65–85%) at both [H62A] and [D197A] mutant receptors (Fig. 2). We constructed zinc inhibition curves for the wild-type receptor and [H62A], [H201A], and [D197A] mutant P2X₇ receptors (Fig. 3). IC_{50} values in these experiments were $6.6 \pm 0.7 \mu\text{M}$ ($n = 15$), $5.9 \pm 1.4 \mu\text{M}$ ($n = 4$), $52 \pm 2.6 \mu\text{M}$ ($n = 10$), and $171 \pm 27 \mu\text{M}$ ($n = 7$) for the wild-type receptor and [H201A], [D197A], and [H62A] mutant receptors, respectively. We also generated a P2X₇ receptor double mutant, [H62A/D197A], and found that the maximum current amplitude was reduced approximately 3-fold; however, the BzATP concentration response curve was shifted significantly to the left at this mutant receptor, resulting in an approximately 4-fold decrease in BzATP EC_{50} value (Table 1). In contrast to these relatively modest alterations in receptor function, the [H62A/D197A] double mutant receptor was virtually resistant to inhibition by zinc at concentrations up to 100 μM (Fig. 3). In addition, no rebound currents were observed upon simultaneous washout of BzATP and zinc for the [H62A] and [H62A/D197A] mutant receptors; however, rebound currents were reduced, but not abolished, for the [D197A] mutant receptor (Figs. 2A and 3A).

Effects of Copper. Copper also inhibits P2X₇ receptors with even greater potency than zinc (Virginio et al., 1997). Here, we also examined copper inhibition of BzATP-evoked currents for the wild-type receptor and 12 functional mutant receptors, using protocols similar to those for zinc inhibition (i.e., 30 μM BzATP and 10 μM copper, a concentration having nearly maximum inhibition). We observed little or no re-

TABLE 1
Summary of BzATP EC_{50} values and maximal currents of wild-type and mutant P2X₇ receptors

Receptors	BzATP EC_{50} μM	n_H	Maximum Currents nA	Cell No
WT	53 ± 4	1.9 ± 0.2	1.8 ± 0.4	9
H62A	68 ± 9	2.1 ± 0.5	1.4 ± 0.2	10
H85A	79 ± 6	1.6 ± 0.1	1.8 ± 0.6	6
H201A	38 ± 3	1.9 ± 0.4	$0.5 \pm 0.1^*$	8
H219A	42 ± 7	1.9 ± 0.2	2.1 ± 0.5	6
H267A	32 ± 2	2.4 ± 0.3	1.5 ± 0.3	4
E70A	57 ± 8	2.7 ± 0.3	1.8 ± 0.3	4
K110A	98 ± 7	4.4 ± 0.7	2.1 ± 0.6	5
E115A	65 ± 15	2.2 ± 0.5	1.1 ± 0.1	5
K137A	35 ± 4	2.4 ± 0.3	1.5 ± 0.3	4
K145A	$196 \pm 11^*$	3.5 ± 0.1	0.6 ± 0.1	6
D197A	88 ± 7	2.7 ± 0.2	3.6 ± 0.7	7
E255A	33 ± 10	1.8 ± 0.3	1.3 ± 0.2	4
H62A/D197A	$14 \pm 1^*$	2.1 ± 0.2	$0.4 \pm 0.1^*$	6

* $P < 0.01$, compared with the wild-type receptor (WT).

bound currents upon simultaneous washout of BzATP/copper from the wild-type or mutant receptors (e.g., Figs. 4A and 5A). Otherwise, results with copper were very similar to those observed for zinc. For the [H62A] and [D197A] but not the other mutant receptors, we observed significantly attenuated inhibition by copper and an even greater decrease in copper inhibition at the [H62A/D197A] double mutant receptor (Fig. 4). We then constructed the copper concentration inhibition curves for the wild-type receptor and the [H62A], [D197A] and [H62A/D197A] mutant receptors, which produced IC₅₀ values of $2.3 \pm 0.2 \mu\text{M}$ ($n = 13$), $12 \pm 1.6 \mu\text{M}$ ($n = 8$), $17 \pm 1.6 \mu\text{M}$ ($n = 8$), and $114 \pm 25 \mu\text{M}$ ($n = 9$) respectively (Fig. 5).

Effects of Zinc and Copper on ATP-Evoked Currents.

Finally, we repeated these experiments using 1 mM ATP (producing ~20–50% of the maximum current of the wild-type and mutant receptors) as the agonist. It was noted that both zinc and copper were slightly less potent in inhibiting the wild-type P2X₇ receptors activated by ATP than by BzATP; the IC₅₀ values for zinc ($19 \pm 1.1 \mu\text{M}$, $n = 4$) and

copper ($6.4 \pm 1.9 \mu\text{M}$, $n = 4$) were 2–3 fold greater than those observed with BzATP as the agonist. The concentrations of zinc (100 μM) and copper (10 μM) used in these experiments resulted in inhibition of the wild-type receptor currents by approximately 70%. The offset kinetics of ATP at the P2X₇ receptor were significantly faster than those observed with BzATP, most likely because BzATP has a higher affinity for this receptor. We also observed no rebound current upon simultaneous washout of ATP with zinc or copper at the wild-type or any of the mutant receptors (Fig. 6). Similar to the results obtained using BzATP as a agonist, we found that zinc and copper inhibition of ATP-evoked currents was potently attenuated or abolished for the [H62A], [D197A], and [H62A/D197A] P2X₇ mutant receptors (Fig. 6). In addition, there was a small but nonetheless significant attenuation of




Fig. 2. Mutational effects on inhibition of BzATP-evoked P2X₇ receptor currents by zinc. A, representative BzATP-evoked currents in the absence or presence of 30 μM zinc at wild-type (WT) and indicated mutant P2X₇ receptors. BzATP concentration was 30 μM except at [K145A] where 200 μM was used. B, summary of the results for wild-type and all the mutant receptors examined. The number of cells examined in each receptor is indicated in brackets. The inhibition was significantly reduced for the [D197A], [H62A], and [H62A/D197A] mutant receptors. *, $p < 0.01$, compared with WT.




Fig. 3. Zinc concentration-current response curves for wild-type and mutant P2X₇ receptors. A, representative BzATP-evoked currents in the presence of increasing concentrations of zinc for wild type (WT), [D197A], [H62A], and [H62A/D197A] mutant P2X₇ receptors. B, concentration-current response curves from experiments as shown in A ($n = 3$ –15 for each point). The IC₅₀ values are $6.6 \pm 0.7 \mu\text{M}$ ($n = 15$), $52 \pm 2.6 \mu\text{M}$ ($n = 10$), and $171 \pm 27 \mu\text{M}$ ($n = 7$) for the WT receptor, [D197A], and [H62A] mutant receptors. The Hill slopes are not significantly different between the WT and mutant receptors.

both zinc and copper inhibition of the [H201A] and [H267A] receptors currents (Fig. 6) that was not observed when BzATP was used as the agonist (compare Figs. 2 and 4).

Discussion

The key finding from this study is the identification of residues His⁶² and Asp¹⁹⁷, located in the ectodomain of the P2X₇ receptor, that are critical for inhibition by the divalent cations zinc and copper. This is of much conceptual significance, because it provides direct evidence against the long-standing view that divalent cation inhibition of P2X₇ receptors results from chelation of ATP⁴⁻, which is considered to be the effective agonist (North, 2002). Indeed, single or combined mutation of His⁶² and Asp¹⁹⁷ results in minimal effects on agonist sensitivity (Table 1) but strongly reduces zinc and copper inhibition, which quite conclusively excludes such an interpretation and substantiates the idea that direct binding is the primary mechanism underlying the inhibition of the P2X₇ receptor by divalent cations. The rebound currents recorded upon simultaneous washout of BzATP and zinc lend

further support. Copper inhibition (IC₅₀ ~ 2–5 μM) was more potent than zinc inhibition (IC₅₀ ~ 5–10 μM), and rebound currents were observed only upon washout of zinc with BzATP but not ATP. The simplest explanation could be that the rebound currents depend upon the relative dissociation rate of the divalent cation and the agonist used. Thus, if the dissociation rate of ATP is > BzATP and that of zinc > BzATP ≥ copper, rebound currents could occur only with the BzATP/zinc combination.

Zinc and copper in submicromolar concentrations have either opposite effects or no actions at other P2X receptors. For example, zinc potently facilitates the P2X₂ and P2X₄ receptor currents (see Introduction). Similar mutational analysis by




Fig. 4. Mutational effects on inhibition of BzATP-evoked P2X₇ receptor currents by copper. A, representative BzATP-evoked currents in the absence or presence of 10 μM copper at wild-type (WT) and indicated mutant P2X₇ receptors. B, summary of the results for all experiments as illustrated in A. The number of cells examined in each receptor is indicated in brackets. Significant attenuation in copper inhibition was seen at [D197A], [H62A] and [H62A/D197A] mutant receptors. *, $p < 0.01$, compared with WT.




Fig. 5. Copper concentration-current response curves for wild-type and mutant P2X₇ receptors. A, representative BzATP-evoked current responses to increasing concentrations of copper at wild-type (WT), [H62A], [D197A], and [H62A/D197A] mutant P2X₇ receptors. B, concentration-current response curves from experiments shown in A ($n = 5-13$ for each point). The IC₅₀ values are $2.3 \pm 0.2 \mu\text{M}$ ($n = 13$), $12 \pm 1.6 \mu\text{M}$ ($n = 8$), $17 \pm 1.6 \mu\text{M}$ ($n = 8$), and $114 \pm 25 \mu\text{M}$ ($n = 9$) for the WT, [H62A], [D197A], and [H62A/D197A] mutant receptors. The Hill slopes are not significantly different between the WT and mutant receptors.

Nagaya et al. (2005) in the ectodomain of P2X₂ receptors identified His¹²⁰ and His²¹³ as the key residues mediating zinc facilitation of P2X₂ receptor. Subsequently, a series of elegant experiments on concatenated P2X₂ trimers convincingly demonstrated that these two residues are located closely at the interface between subunits and form an inter-subunit zinc binding site. There is no information regarding how close His⁶² and Asp¹⁹⁷ residues are in the P2X₇ receptor, but similar approaches may be expected to determine whether these two key residues form interior intrasubunit zinc/copper binding sites. It is noteworthy that our functional characterization (Table 1) showed that alanine substitution of both residues together ([H62A/D197A] double mutant), but not individually, resulted in a significant increase in the agonist sensitivity. This is reminiscent of the mutational effects on the close apposition and direct interaction between the N-terminal ligand binding domain and the extracellular loops between transmembrane segments in the pore forming domain, which underlie the gating of the GABA_A (Kash et al., 2003) and nACh receptors (Lee and Sine, 2005). Although the mutational effects on agonist sensitivity are not fully understood, one possibility is that His⁶² and Asp¹⁹⁷ may be closely positioned. His⁶² is located between Lys⁶⁴, the critical resi-

due for P2X₇ receptor activation (Wilkinson et al., 2006), and the first transmembrane domain, which is also implicated in channel gating (Jiang et al., 2001). An alternative explanation is that replacement of both His⁶² and Asp¹⁹⁷ with alanine (a residue that contains a small side chain) may facilitate, to some extent, the channel gating. Consistent with this idea, occupation by divalent cations would disfavor channel gating and thereby results in a rightward shift and suppression of maximal receptor responses in the agonist concentration response curves as previously observed (Virginio et al., 1997). Finally, the present study has also shown a critical role of the nonhistidine residue (Asp¹⁹⁷) in coordinating zinc and copper inhibition of the P2X₇ receptor, similar to zinc inhibition of the *N*-methyl-D-aspartate receptor (Paoletti et al., 2000).

We observed modest, but significant differences depending on the agonists used: first, zinc and copper were 2- to 3-fold more potent at inhibiting BzATP-evoked currents than ATP-evoked currents. Second, when BzATP was used to evoke currents, only single and double mutations of His⁶² and Asp¹⁹⁷ resulted in significant attenuation of zinc and copper inhibition. These mutant receptors also exhibited the strongest attenuation of zinc and copper inhibition when ATP was




Fig. 6. Inhibition of ATP-evoked P2X₇ receptor currents by zinc and copper. A, left, representative currents of wild-type (WT) and mutant P2X₇ receptors activated by ATP (1 mM) in the absence and presence of zinc (100 μM). Right, summary of the results as shown on the left. B, left, representative currents of wild-type (WT) and mutant P2X₇ receptors activated by ATP (1 mM) in the absence and presence of copper (10 μM). Right, summary of the results as shown on the left. The number of cells examined in each case is indicated in brackets. *, *p* < 0.05; **, *p* < 0.01, compared with WT. Note the differential inhibition of [H62A] and [D197A] mutant receptors by zinc and copper.

the agonist, but we also observed significant attenuation at the [H201A] and [H267A] mutant receptors (compare Figs. 2B, 4B, and 7). We have identified two residues in the ectodomain that could account for the differential agonist (BzATP versus ATP) sensitivity at the P2X₇ receptor. We demonstrated that Lys¹²⁷ and Asn²⁸⁴ together are required to maintain BzATP potency, whereas Asn²⁸⁴ alone is involved in ATP potency. We therefore suggest that these findings may indicate overlapping but distinct regions of the P2X₇ receptor that bind BzATP and ATP (Young et al., 2007). If binding of BzATP involves residues in addition to those required for ATP binding, it is reasonable to imagine a similar overlapping but distinct binding of divalent cations in the presence of each of these agonists. In any event, the actions of zinc and copper share the same two key residues (His⁶² and Asp¹⁹⁷), both of which are required to inhibit currents evoked by either agonist. Another possible contribution to the modest agonist-dependent difference seen here is the potential regulation of the P2X₇ receptor by the endogenous P2Y receptors (Schachter et al., 1997) as shown for the P2X₁ receptor (Vial et al., 2004). However, such an effect should be minimal when BzATP is used (Wilson et al., 2002).

During the preparation of the present study, an article by Acuña-Castillo et al. (2007) appeared in which they examined the role of histidine residues in zinc and copper inhibition of ATP-evoked rat P2X₇ receptor currents in *Xenopus laevis* oocytes expressing single alanine mutant receptors. In agreement with our findings, this group also observed no significant effects on agonist sensitivity by mutating the histidine residues and reached the same conclusion that zinc and copper inhibition of the P2X₇ receptor has little to do with changes in the active form of the agonist. However, their results differ considerably from ours in terms of the contribution these residues make. Acuña-Castillo et al. (2007) showed that the [H201] and [H267A] mutant receptors had either a significant reduction in or loss of inhibition by copper, whereas the [H219A] and [H267A] mutant receptors were both insensitive to inhibition by zinc. We also demonstrated a reduced zinc or copper inhibition of the [H201A] and [H267A] mutant receptors when ATP was used as the agonist, but clearly the reduction was far less pronounced compared with [H62A] and [D197A] mutants (the latter mutant was not studied by Acuña-Castillo et al., 2007) (Fig. 6). The inhibition became variable and insignificantly different from the wild-type receptor when BzATP was the agonist (Fig. 2B), probably because of the relatively higher potency of zinc and copper (see *Results*). The most surprising difference between the two studies is that we found the most prominent decrease in zinc and copper inhibition of P2X₇ receptor currents at the [H62A] mutant receptor, regardless of whether we used BzATP (Figs. 2–5) or ATP (Fig. 6), whereas Acuña-Castillo et al. (2007) observed no effects evoked by ATP. Inconsistent results have been noted for P2X₇ receptors expressed in mammalian cells and oocytes. The most noticeable difference is that just as in native P2X₇ receptor expressing cells, formation of the large dye uptake pore was persistently observed when the P2X₇ receptors were expressed in mammalian cells (e.g., Surprenant et al., 1996) but not in oocytes (Petrou et al., 1997; Klapperstück et al., 2000). In addition, loss of function associated with a naturally occurring mutation E496A in monocytes and lymphocytes (Gu et al., 2001) was confirmed in HEK293 cells (Cabrini et al., 2005). How-

ever, the [E496A] mutant receptor was functional and insignificantly differed from the wild-type receptor in oocytes as well as in HEK293 cells in another study (Boldt et al., 2003). Thus, utilization of the expression cells and in particular different cells (mammalian cell versus oocyte) may contribute to the discrepancy observed in the different studies.

Inhibition of P2X₇ receptors by zinc and copper has significant physiological relevance. For example, P2X₇ receptors are expressed in brain astrocytes and glial cells where they are increasingly known to engage in neural signaling processes such as neurotransmitter release (Duan and Neary, 2006). Local concentrations of zinc and copper released from nerve terminals are in the range of 10 to 100 μM (Kardos et al., 1989; Li et al., 2001) and can easily reach astrocytes and glial cells, which are in close vicinity and thus inhibit P2X₇ receptors. P2X₇ receptors are highly expressed in immune cells and important to the physiology and pathology of these cells (see Introduction). Under conditions of infection and/or inflammation, copper is found at micromolar concentrations and is essential for both T-cell proliferation and the ability of neutrophils to generate superoxide to kill ingested microorganisms (Percival, 1998). The present study has revealed, at the molecular level, how zinc and copper modulate the P2X₇ receptor.

In summary, we have provided direct evidence showing that zinc and copper inhibition is due to a direct interaction with ectodomain residues of the P2X₇ receptor, primarily involving combined interaction with His⁶² and Asp¹⁹⁷. The present study enhances our understanding of how this multifunctional ionotropic purinergic receptor can be activated by agonist, and in particular modulated by trace metals.

Acknowledgments

We are grateful to Dr. C. Milligan for critical comments on the manuscript.

References

- Acuña-Castillo C, Coddou C, Bull P, Brito J, and Huidobro-Toro JP (2007) Differential role of extracellular histidines in copper, zinc, magnesium and proton modulation of the P2X₇ purinergic receptor. *J Neurochem* **101**:17–26.
- Acuña-Castillo C, Morales B, and Huidobro-Toro JP (2000) Zinc and copper modulate differentially the P2X₄ receptor. *J Neurochem* **74**:1529–1537.
- Boldt W, Klapperstück M, Buttner C, Sadtler S, Schmalzing G, and Markwardt F (2003) Glu496Ala polymorphism of human P2X₇ receptor does not affect its electrophysiological phenotype. *Am J Physiol Cell Physiol* **284**:C749–C756.
- Cabrini G, Falzoni S, Forchop SL, Pellegatti P, Balboni A, Agostini P, Cuneo A, Castoldi G, Baricordi OR, and Di Virgilio F (2005) A His-155 to Tyr polymorphism confers gain-of-function to the human P2X₇ receptor of human leukemic lymphocytes. *J Immunol* **175**:82–89.
- Chessell IP, Hatcher JP, Bountra C, Michel AD, Hughes JP, Green P, Egerton J, Murfin M, Richardson J, Peck WL, et al. (2005) Disruption of the P2X₇ purinoceptor gene abolishes chronic inflammatory and neuropathic pain. *Pain* **114**:386–396.
- Chu XP, Wemmie JA, Wang WZ, Zhu XM, Saugstad JA, Price MP, Simon RP, and Xiong ZG (2004) Subunit-dependent high-affinity zinc inhibition of acid-sensing ion channels. *J Neurosci* **24**:8678–8689.
- Clyne JD, LaPointe LD, and Hume RI (2002) The role of histidine residues in modulation of the rat P2X₂ purinoceptor by zinc and pH. *J Physiol* **539**:347–359.
- Cockcroft S and Gomperts BD (1979) Activation and inhibition of calcium-dependent histamine secretion by ATP ions applied to rat mast cells. *J Physiol* **296**:229–243.
- Duan S and Neary JT (2006) P2X₇ receptors: properties and relevance to CNS function. *Glia* **15**:738–746.
- Ferrari D, Pizzirani C, Adinolfi E, Lemoli RM, Curti A, Idzko M, Panther E, and Di Virgilio F (2006) The P2X₇ receptor: a key player in IL-1 processing and release. *J Immunol* **176**:3877–3883.
- Gu BJ, Zhang W, Worthington RA, Sluyter R, Dao-Ung P, Petrou S, Barden JA, and Wiley JS (2001) A Glu-496 to Ala polymorphism leads to loss of function of the human P2X₇ receptor. *J Biol Chem* **276**:11135–11142.
- Jiang LH, Rassendren F, Spelta V, Surprenant A, and North RA (2001) Amino acid residues involved in gating identified in the first membrane-spanning domain of the rat P2X₂ receptor. *J Biol Chem* **276**:14902–14908.
- Kash TL, Jenkins A, Kelley JC, Trudell JR, and Harrison NL (2003) Coupling of agonist binding to channel gating in the GABA(A) receptor. *Nature* **421**:272–275.

- Kardos J, Kovacs I, Hajos F, Kalman M, and Simonyi M (1989) Nerve endings from rat brain tissue release copper upon depolarization: A possible role in regulating neuronal excitability. *Neurosci Lett* **103**:139–144.
- Klapperstück M, Buttner C, Bohm T, Schmalzing G, and Markwardt F (2000) Characteristics of P2X₇ receptors from human B lymphocytes expressed in *Xenopus* oocytes. *Biochim Biophys Acta* **1467**:444–456.
- Laube B, Kuhse J, and Betz H (2000) Kinetic and mutational analysis of Zn²⁺ modulation of recombinant human inhibitory glycine receptors. *J Physiol* **522**:215–230.
- Lammas DA, Stober C, Harvey CJ, Kendrick N, Panchalingam S, and Kumararatne DS (1997) ATP-induced killing of mycobacteria by human macrophages is mediated by purinergic P2Z(P2X7) receptors. *Immunity* **7**:433–444.
- Lee WY and Sine SM (2005) Principal pathway coupling agonist binding to channel gating in nicotinic receptors. *Nature* **438**:243–247.
- Li Y, Hough CJ, Suh SW, Sarvey JM, and Frederickson CJ (2001) Rapid translocation of Zn²⁺ from presynaptic terminals into postsynaptic hippocampal neurons after physiological stimulation. *J Neurophysiol* **86**:2597–2604.
- Lorca RA, Coddou C, Gazitua MC, Bull P, Arredondo C, and Huidobro-Toro JP (2005) Extracellular histidine residues identify common structural determinants in the copper/zinc P2X₂ receptor modulation. *J Neurochem* **95**:499–512.
- Morelli A, Chiozzi P, Chiesa A, Ferrari D, Sanz JM, Falzoni S, Pinton P, Rizzuto R, Olson MF, and Di Virgilio F (2003) Extracellular ATP causes ROCK I-dependent bleb formation in P2X₇-transfected HEK293 cells. *Mol Biol Cell* **14**:2655–2664.
- Nagaya N, Tittle RK, Saar N, Dellal SS, and Hume RI (2005) An intersubunit zinc binding site in rat P2X₂ receptors. *J Biol Chem* **280**:25982–25993.
- North RA (2002) Molecular physiology of P2X receptors. *Physiol Rev* **82**:1013–1067.
- Paoletti P, Perin-Dureau F, Fayyazuddin A, Le Goff A, Callebaut I, and Neyton J (2000) Molecular organization of a zinc binding N-terminal modulatory domain in a NMDA receptor subunit. *Neuron* **28**:911–925.
- Percival SS (1998) Copper and immunity. *Am J Clin Nutr* **67**:1064S–1068S.
- Petrou S, Ugur M, Drummond RM, Singer JJ, and Walsh JV Jr (1997) P2X₇ purinoceptor expression in *Xenopus* oocytes is not sufficient to produce a pore-forming P2Z-like phenotype. *FEBS Lett* **411**:339–345.
- Schachter JB, Sromek SM, Nicholas RA, and Harden TK (1997) HEK293 human embryonic kidney cells endogenously express the P2Y₁ and P2Y₂ receptors. *Neuropharmacology* **36**:1181–1187.
- Solle M, Labasi J, Perregaux DG, Stam E, Petrushova N, Koller BH, Griffiths RJ, and Gabel CA (2001) Altered cytokine production in mice lacking P2X(7) receptors. *J Biol Chem* **276**:125–132.
- Spelta V, Jiang LH, Surprenant A, and North RA (2003) Kinetics of agonist actions at rat P2X_{2/3} heteromeric receptors. *Br J Pharmacol* **135**:1524–1530.
- Surprenant A, Rassendren F, Kawashima E, North RA, and Buell G (1996) The cytolytic P2Z receptor for extracellular ATP identified as a P2X receptor (P2X₇). *Science* **272**:735–738.
- Vial C, Tobin AB, and Evans RJ (2004) G-protein-coupled receptor regulation of P2X₁ receptors does not involve direct channel phosphorylation. *Biochem J* **382**:101–110.
- Virginio C, Church D, North RA, and Surprenant A (1997) Effects of divalent cations, protons and calmidazolium at the rat P2X₇ receptor. *Neuropharmacology* **36**:1285–1294.
- Wang TL, Hackam A, Guggino WB, and Cutting GR (1995) A single histidine residue is essential for zinc inhibition of GABA ρ 1 receptors. *J Neurosci* **15**:7684–7691.
- Wildman SS, King BF, and Burnstock G (1998) Zn²⁺ modulation of ATP-responses at recombinant P2X₂ receptors and its dependence on extracellular pH. *Br J Pharmacol* **123**:1214–1220.
- Wildman SS, King BF, and Burnstock G (1999a) Modulation of ATP-responses at recombinant rP2X₄ receptors by extracellular pH and zinc. *Br J Pharmacol* **126**:762–768.
- Wildman SS, King BF, and Burnstock G (1999b) Modulatory activity of extracellular H⁺ and Zn²⁺ on ATP-responses at rP2X₁ and rP2X₃ receptors. *Br J Pharmacol* **128**:486–492.
- Wilkinson WJ, Jiang LH, Surprenant A, and North RA (2006) Role of ectodomain lysines in the subunits of the heteromeric P2X_{2/3} receptor. *Mol Pharmacol* **70**:1159–1163.
- Wilson HL, Wilson SA, Surprenant A, and North RA (2002) Epithelial membrane proteins induce membrane blebbing and interact with the P2X₇ receptor C terminus. *J Biol Chem* **277**:34017–34023.
- Xiong K, Peoples RW, Montgomery JP, Chiang Y, Stewart RR, Weight FF, and Li C (1999) Differential modulation by copper and zinc of P2X₂ and P2X₄ receptor function. *J Neurophysiol* **81**:2088–2094.
- Young MT, Pelegrin P, and Surprenant A (2007) Amino acid residues in the P2X₇ receptor that mediate differential sensitivity to ATP and BzATP. *Mol Pharmacol* **71**:92–100.

Address correspondence to: Dr. Lin-Hua Jiang, Institute of Membrane and Systems Biology, Faculty of Biological Sciences, University of Leeds, Leeds LS2 9JT, UK. E-mail: l.h.jiang@leeds.ac.uk
