

Aging-Resistant Organophosphate Bioscavenger Based on Polyethylene Glycol-Conjugated F338A Human Acetylcholinesterase

Ohad Mazor, Ofer Cohen, Chanoch Kronman, Lily Raveh, Dana Stein, Arie Ordentlich, and Avigdor Shafferman

Departments of Biochemistry and Molecular Genetics (O.M., O.C., C.K., D.S., A.O., A.S.) and Pharmacology (L.R.), Israel Institute for Biological Research, Ness-Ziona, Israel

Received March 25, 2008; accepted June 2, 2008

ABSTRACT

The high reactivity of cholinesterases (ChEs) toward organophosphorus (OP) compounds has led to propose recombinant ChEs as bioscavengers of nerve agents. The bioscavenging potential of recombinant ChEs can be enhanced by conjugation of polyethylene glycol (PEG) moieties, to extend their circulatory residence. However, the ability of exogenously administered ChEs to confer long-term protection against repeated exposures to nerve agents is still limited due to the aging process, whereby organophosphate-ChE adducts undergo irreversible dealkylation, which precludes oxime-mediated reactivation of the enzyme. To generate an optimal acetylcholinesterase (AChE)-based OP bioscavenger, the F338A mutation, known to decelerate the rate of aging of AChE-OP conjugates, was incorporated into polyethylene glycol-conjugated (PEGylated) human AChE. The PEGylated F338A-AChE displayed unaltered rates of hydrolysis, inhibition, phosphorylation, and reactivation and could effectively protect mice against exposure to soman (pinacolylmethyl phosphonofluoridate), sarin (O-isopropyl methylphosphonofluoridate), or O-ethyl-S-

(2-isopropylaminoethyl) methylphosphonothioate (VX). Unlike PEGylated wild-type (WT)-AChE, the PEGylated F338A-AChE exhibits significantly reduced aging rates after soman inhibition and can be efficiently reactivated by the 1-[[[4(aminocarbonyl)pyridinio]methoxy]methyl]-2(hydroxyimino)methyl]pyridinium dichloride (HI-6) oxime, both in vitro and in vivo. Accordingly, oxime administration to PEG-F338A-AChE-pretreated mice enabled them to withstand repeated soman exposure (5.4 and 4 LD₅₀/dose), whereas same regime treatment of non-PEGylated F338A-AChE- or PEGylated WT-AChE-pretreated mice failed to protect against the second challenge, due to rapid clearance or irreversible aging of the latter enzymes. Thus, judicious incorporation of selected mutations into the AChE mold in conjunction with its chemical modification provides means to engineer an optimal ChE-based OP bioscavenger in terms of circulatory longevity, resistance to aging, and reduced doses required for protection, even against repeated exposures to nerve agents, such as soman.

Acetylcholinesterase plays a central role in the cholinergic system, where it hydrolyzes the neurotransmitter acetylcholine, thereby terminating nerve impulse transmission. This enzyme is the major target for organophosphorus nerve agents, such as soman (pinacolylmethyl phosphonofluoridate), sarin (O-isopropyl methylphosphonofluoridate), and O-ethyl-S-(2-isopropylaminoethyl) methylphosphonothioate

(VX), which irreversibly phosphorylate the catalytic serine residue within the enzyme active site. Inactivation of AChE leads to continuous stimulation of cholinergic receptors, resulting in symptoms of toxicity such as salivation, tremors, and miosis and in severe cases, respiratory paralysis and death. Current drug treatment against nerve agent exposure is based on administration of cholinolytic drugs (e.g., atropine) and oximes, which displace the phosphyl group to restore AChE activity, and of anticonvulsants, which reduce OP intoxication-associated convulsions. Yet, although the combined postexposure treatment is effective in preventing lethality, it does not prevent severe incapacitation, including

This work was supported by the U.S. Army Research and Development Command Contract DAMD17-03-C-0012 (to A.S.)

Article, publication date, and citation information can be found at <http://molpharm.aspetjournals.org>.
doi:10.1124/mol.108.047449.

ABBREVIATIONS: VX, O-ethyl-S-(2-isopropylaminoethyl) methylphosphonothioate; AChE, acetylcholinesterase; OP, organophosphorus; ChE, cholinesterase; rHu, recombinant human; PEG, polyethylene glycol; PEGylated, polyethylene glycol-conjugated; WT, wild type; PBS, phosphate-buffered saline; MEPQ, 7-(methylethoxyphosphinyloxy)-1-methylquinolinium iodide; CBDP, cresylbenzodioxaphosphorin oxide; HI-6, 1-[[[4(aminocarbonyl)pyridinio]methoxy]methyl]-2(hydroxyimino)methyl]pyridinium dichloride; BW284c51, 1,5-bis(4-allyldimethylammoniumphenyl)-pentan-3-one dibromide; BChE, butyrylcholinesterase; MRT, mean residence time; ATC, acetylthiocholine iodide.

behavioral difficulties and permanent brain damage (Dawson, 1994).

The high reactivity of ChEs toward OP nerve agents has led to propose their use as exogenous biological scavengers for the sequestering of toxic OP agents before they reach their physiological target (Raveh et al., 1993, 1997; Cohen et al., 2006; Kronman et al., 2007; Lenz et al., 2007). Being a stoichiometric scavenger, it was estimated that approximately 200 mg of exogenous enzyme is needed to protect adult humans against 2 LD₅₀ of soman (Ashani et al., 1998). This requirement for large quantities of enzyme encouraged the development of production systems for the generation of recombinant ChEs on a large scale (Kronman et al., 1992; Fischer et al., 1993; Masson et al., 1993). However, pharmacokinetic studies revealed that these recombinant enzymes display rapid clearance profiles, which limits their use as efficient bioscavengers that could confer long-term protection (Kronman et al., 1995). To overcome this limitation, chemical modification of recombinant human-AChE (rHuAChE) via controlled appendage of polyethylene glycol (PEG) to lysine side chains was performed (Cohen et al., 2001, 2004). The PEGylated rHuAChE exhibited very long retention times in mice and monkeys and allowed effective protection of mice against OP nerve agents (Cohen et al., 2004, 2006; Kronman et al., 2007).

Upon exposure to OP agents, phosphorylation of the inhibited AChE-active site serine results in the formation of a stable conjugate, which can further undergo a unimolecular process (Fleisher and Harris, 1965), resulting in an "aged" enzyme form that is refractive to oxime reactivation (Scheme 1) (see also Shafferman et al., 1996). This aging process is most pronounced for OPs containing a branched alkyl group, such as soman (Barak et al., 1997). Inhibition of exogenously administered scavenger ChEs, by aging, limits their ability to confer long-term protection against repeated exposures to nerve agents. Such a limitation may be alleviated by the development of AChE-based bioscavengers with reduced aging rates, which can thereby be effectively reactivated by oximes.

Over the past two decades, extensive research was carried out to understand the basic aspects of the mechanism of catalysis, phosphorylation, and aging of AChE. Such studies included X-ray crystallography (Sussman et al., 1991; Millard et al., 1999), site-directed mutagenesis and molecular modeling, together with kinetic studies of the AChE mutants with substrates and reversible inhibitors (Ordentlich et al., 1993, 1996; Radić et al., 1993; Barak et al., 1994; Shafferman et al., 1996, 2005). These studies revealed several functional subsites in the AChE active center gorge that are essential for accommodation of the various groups of AChE ligands, as

well as for the pronounced activity of the enzyme (for review, see Shafferman et al., 2005). Among other findings, these studies allowed to identify the amino acid residues that play a role in the enzyme aging process (Ordentlich et al., 1993, 1996). Most notably, replacement of the Phe338 by alanine (F338A-AChE) resulted in a pronounced decrease (160-fold) in the rate of aging after soman inhibition (Shafferman et al., 1996) compared with the WT enzyme. Consequently, the adduct of F338A HuAChE with soman could be reactivated much in the same way as adducts of HuAChE with sarin.


In the present study, we generated a PEGylated human AChE enzyme based upon the F338A mutant (PEG-F338A-AChE), which exhibits an improved pharmacokinetic profile and could be effectively reactivated after inhibition by various OP compounds. In a series of *in vivo* experiments, we demonstrate that PEG-F338A-AChE, unlike PEG-WT-AChE, effectively protects mice against repeated soman exposure in conjunction with oxime treatment. Thus, incorporation of the F338A mutation into the PEGylated AChE template may provide the means to generate a superior prophylactic agent that can confer long-term protection against OP compound exposures.

Materials and Methods

Production of Enzymes. Transfection of the human embryonic kidney 293 cell line with expression vectors coding for the C-terminally truncated WT- or F338A-HuAChE enzymes (Grosfeld et al., 1996; Cohen et al., 2001) and the generation of stable cell clones expressing high levels of recombinant product were described previously (Kronman et al., 1992). AChEs were purified as described previously (Kronman et al., 1995). Attachment of PEG chains to primary amines of AChEs was performed using succinimidyl propionate-activated methoxy-PEG (Nektar Inc., San Carlos, CA) as described previously (Cohen et al., 2001). In brief, purified AChE (5 μM) was incubated with PEG-20000 at a ratio of 30:1 (mol/mol) [PEG]/[AChE primary amines] in 50 mM borate buffer, pH 8.5, for 2 h at room temperature. The chemically modified products were dialyzed extensively against PBS and visualized on 6% SDS-polyacrylamide gel electrophoresis gels.

Organophosphate Inhibitors. Soman, sarin, VX, 7-(methoxyphosphoryloxy)-1-methylquinolinium iodide (MEPQ), and cresylbenzodioxaphosphorin oxide (CBDP) were prepared as described in procedures reported previously (Levy and Ashani, 1986; Cohen et al., 2006). The purity of the OPs (>95%) was determined by ¹H and ³¹P NMR spectroscopy and by a titration with a defined amount of HuAChE. Stock solutions were kept at -20°C and diluted in 0.9% saline to the desired concentration before use.

Kinetic Studies. HuAChE activity was assayed according to Ellman et al. (1961) in the presence of 0.2 mg/ml bovine serum albumin, 0.3 mM 5,5'-dithio-bis-(2-nitrobenzoic acid, 50 mM sodium phosphate buffer, pH 8.0, and 0.5 mM acetylthiocholine iodide (ATC) at 27°C and monitored with a Thermomax microplate reader (Molecular Devices, Sunnyvale, CA). Measurements of phosphorylation rates were carried out by monitoring residual activity (E) at various time points, after incubation of the enzyme in the presence of at least four different concentrations of OP inhibitor (I). The apparent bimolecular phosphorylation rate constants (*k_i*) determined under pseudo-first-order conditions were computed from the plot of slopes of ln(E) versus time at different inhibitor concentrations (Ordentlich et al., 1996, 1999). Rate constants under second-order conditions were determined from plots of ln[E/(I₀ - (E₀ - E))] versus time. Stereoselectivity of the enzymes toward soman was determined by active site titrations, comparing residual activities of enzymes inhibited by the appropriate racemic phosphonate to that of MEPQ (Shafferman et al., 1996).



Scheme 1. Phosphorylation of ChEs (EOH), reactivation, and aging.

Measurements of aging rates were performed essentially as described previously (Shafferman et al., 1996), where the reactivatable fraction of the OP-inhibited AChEs was determined after reactivation with 0.5 mM HI-6. The activity of the reactivated enzyme (E_r) was routinely corrected for the inhibitory effect of the reactivator. The first-order rate constants of aging (k_a) were determined from the slopes of $\ln(E_r/E_0)$ against time.

Measurements of the reactivation kinetics rates were performed as described previously (Grosfeld et al., 1996). In brief, reactivation reactions of the OP-inhibited AChEs (in 50 mM phosphate buffer, pH 8.0) were initiated by the addition of HI-6 at various concentrations (0.005–1 mM) and incubated at 27°C. Regeneration of AChE activity was then monitored, and compared with noninhibited enzyme preparations that served as control for the expected enzyme activity at maximal reactivation. For the in vitro reactivation of soman-inhibited AChE, heparinized blood of CBDP-treated mice was spiked with either PEGylated WT- or F338A-AChE (0.8 μ M) and then inhibited with soman, at a concentration (1.6 μ M) that leads to 90% inactivation. HI-6 (0.5 mM) was added to the enzyme/soman mixture after incubation for 20 min at 37°C. Reactivation of AChE in the blood samples was determined as described above.

Animal Studies. Male outbred ICR mice (Charles River Laboratories, Inc., Wilmington, MA) were maintained at 20–22°C and a relative humidity of 50 \pm 10% on a 12-h light/dark cycle, fed with commercial rodent chow (Koffolk, Inc., Tel Aviv, Israel), and provided with tap water ad libitum. Treatment of animals was in accordance with regulations outlined in the USDA Animal Welfare Act and the conditions specified in Guide for Care and Use of Laboratory Animals (National Institute of Health, 1996). Animal studies were approved by the local ethical committee on animal experiments.

In vivo reactivation of the soman-inhibited AChEs was initiated 20 min after soman intoxication, by the intravascular administration of 50 mg/kg HI-6, a well tolerated dose without any signs of distress. Residual AChE activity was determined in blood samples withdrawn at various time points throughout the experiments.

For challenge experiments, mice intravenously administered with AChEs were exposed intravenously to OPs. Before exposure to soman or sarin, mice were pretreated with 2 mg/kg CBDP 1 h before challenge to inhibit endogenous carboxylesterases, as described previously (Cohen et al., 2006). In experiments where repeated soman challenge was performed, a second administration of CBDP (1 mg/kg) was performed immediately before AChE treatment. Mice were monitored for toxic signs and mortality for 4 h after OP administration.

Clearance experiments in mice (three to six mice per enzyme sample) were carried out essentially as described previously (Cohen et al., 2006). Mice were injected intravenously with 200 units/mouse of native or PEGylated AChEs, in 0.2 ml of PBS. At different time points, blood samples (5 μ l) were drawn from the tail vein, diluted 20-fold in PBS, and centrifuged for 3 min at 3000 rpm for the removal of red blood cells. Residual AChE activity in the serum was measured, and values were corrected for background hydrolytic activity in the blood. Pharmacokinetic parameters were calculated using the PK Solutions software (Summit Research Services, Ashland, OH).

Results

Reactivity of PEGylated F338A-AChE toward Active-Center Inhibitors and OP Agents

F338A mutated HuAChE was conjugated to polyethylene glycol under conditions that should promote high-level PEGylation without compromising the catalytic activity of the chemically modified enzyme product (Cohen et al., 2001). Kinetic analyses of PEGylated F338A-AChE, allowed us to determine that this version of the enzyme hydrolyzes ATC, with a K_m value of 0.2 mM, similar to its non-PEGylated counterpart (0.14 mM; Table 1). Likewise, the PEGylated and non-PEGylated F338A-AChE exhibit comparable values of k_{cat} and k_{app} . These values were also similar to those exhibited by both the non-PEGylated and PEGylated versions of the WT enzyme (Table 1). Thus, the appendage of PEG chains to F338A-AChE did not deleteriously affect the catalytic efficiency of the enzyme.

We next determined the inhibition kinetic constants of PEG-F338A-AChE toward several AChE active-center inhibitors. PEGylation of both F338A AChE and WT enzymes did not affect their interactions with edrophonium, with both exhibiting K_i values (1.2 μ M) similar to those of the nonmodified enzymes (0.8 μ M; Table 1). Likewise, binding of PEG chains to the enzyme surface did not affect the inhibition constant of PEG-F338A-AChE or PEG-WT-AChE toward the peripheral anionic site ligand propidium or the bisquaternary inhibitor BW284c51; and in both cases, the measured K_i values were in the same range as determined earlier for the non-PEGylated enzyme forms (Table 1).

The kinetic properties of F338A AChEs were further examined by determining the reactivity of this enzyme toward various OP compounds. Determination of the apparent bimolecular rate constant (k_i) of PEG-F338A-AChE for soman phosphorylation, demonstrated that the chemically modified enzyme retained its reactivity toward soman (Table 2). Likewise, the inhibition rate constant of the PEG-F338A-AChE toward sarin was found to be similar to that of the non-PEGylated F338A-AChE form and of WT-AChE (Table 2). In the case of VX, the reactivity of both non-PEGylated and PEGylated F338A-AChE were also similar; yet, these values were found to be slightly lower (\sim 4-fold) than those displayed by the non-PEGylated and PEGylated WT enzymes (Table 2). This difference in reactivity toward VX has been observed previously, and it can be related to a minor disruption within the active center environment caused by the Phe338 to Ala replacement, which affects the accommodation of positively charged OP molecules such as VX, within the active center (Ordentlich et al., 2005). PEGylation of F338A AChE did not

TABLE 1
ATC hydrolysis rates and inhibition constants (k_i) of WT- and F338A-AChEs
Values represents mean of triplicate determination with S.D. not exceeding 20%.

AChE Species	ATC			K_i		
	K_m	k_{cat}	k_{app}	Edrophonium	Propidium	BW284c51
	mM	$\times 10^{-5} \text{ min}^{-1}$	$\times 10^8 \text{ M}^{-1} \text{ min}^{-1}$		μM	
WT ^a	0.14	4	29	0.75	1.4	10
F338A ^a	0.14	1.7	12	0.8	1.5	10
PEG-WT	0.17	6.4	38	1.1	1.6	12.5
PEG-F338A	0.2	2.7	14	1.2	1.1	16.5

^a Values compiled from our previous studies (Ordentlich et al., 1993) are presented here for comparison.

reduce its reactivity toward the organophosphate compound paraoxon, compared with its non-PEGylated counterpart (Table 2). The reactivity of PEG-F338A-AChE was about 10-fold lower than that of PEG-WT-AChE, analogous to the difference in reactivities toward paraoxon of the nonmodified forms of F338A-AChE and WT-AChE (Table 2). This was previously shown to be related specifically to this organophosphate compound, where in the generation of the AChE-OP adduct, the leaving group is aromatic. Indeed, for diethyl phosphorofluoridate, which generates the same AChE adduct as paraoxon but in which the leaving group is smaller (fluoride in diethyl phosphorofluoridate versus *p*-nitrophenol in paraoxon), the rate of phosphorylation is not affected by the F338A mutation (Ordentlich et al., 1996).

AChEs are characterized by a pronounced stereoselectivity toward various OP agents, such as soman and sarin (de Jong et al., 1988). To determine that stereoselectivity is not affected in the modified enzyme, active site titration analysis of PEG-F338A-AChE and PEG-WT-AChE was performed with soman. In parallel, active site titration analysis of both enzymes was performed with MEPQ, an OP inhibitor known to react with AChE at a 1:1 stoichiometric ratio (i.e., both the P_S and P_R enantiomers react equally with AChE). By comparing the intercept for zero enzymatic activity with soman to that obtained with MEPQ (Fig. 1A), we could determine a 1:2 stoichiometry for the interactions with soman of the PEGylated forms of the WT- and F338A-AChEs. Thus, the PEGylated F338A-mutated version of AChE retains its pronounced stereoselectivity toward the toxic soman diastereomers.

As mentioned, WT-AChE undergoes rapid aging in the presence of soman (Barak et al., 1997). In contrast, the F338A-AChE mutant exhibits a pronounced decrease in its rate of aging after soman inhibition (Shafferman et al., 1996). To verify whether the PEGylated version of F338A AChE retains this unique trait, the first-order rate constant (k_a) of aging was determined for the soman-inhibited PEGylated AChE enzymes (WT and F338A), by measuring the enzyme activity restored in the presence of the oxime HI-6, which is known to be a potent reactivator of soman adducts (de Jong and Kossen, 1985; Dawson, 1994). PEGylated WT-AChE was rapidly aged (Fig. 1B, inset), displaying an apparent k_a value of $95 \times 10^3 \text{ min}^{-1}$. In contrast, the aging rate of the soman-inhibited PEG-F338A-AChE was found to be ~ 170 -fold slower than that of PEG-WT enzyme (Fig. 1B), exhibiting an apparent k_a value of $0.55 \times 10^3 \text{ min}^{-1}$. These results are in good correlation with our previous data in which a 160-fold difference was found between the aging rates of the WT- and the F338A-AChE native forms (approx. k_a values of $130 \times$

10^3 min^{-1} and $0.8 \times 10^3 \text{ min}^{-1}$, respectively; Shafferman et al., 1996; Grosfeld et al., 1996).

The ability of oximes to effectively restore the enzymatic activity of OP-inhibited AChE depends not only on the rate of aging of the enzyme but also on its reactivation kinetics. To quantify the reactivability of OP-inhibited F338A AChEs, we monitored the increase in enzyme activity over time after the addition of HI-6, and determined the apparent reactivation rate constant, k_r , and the dephosphorylation rate constant, k_2 (Table 3). Chemical modification of the F338A AChE by PEG conjugation did not alter its ability to interact with the HI-6, resulting in similar reactivation rates for both for the native ($k_r = 300 \text{ M}^{-1} \text{ min}^{-1}$; $k_2 = 2.3 \times 10^2 \text{ min}^{-1}$) and the PEGylated ($k_r = 254 \text{ M}^{-1} \text{ min}^{-1}$; $k_2 = 1.6 \times 10^2 \text{ min}^{-1}$) forms of this enzyme. In the WT-AChEs, reactivation kinetics values could not be determined due to the rapid aging of its adducts with soman (see above). In contrast, the aging process of sarin and VX inhibited AChE is much slower (Barak et al., 1997; Worek et al., 2004); therefore, with these inhibitors, it was possible to determine and compare the reactivation rate constants of the F338A- and WT-AChEs.

Reactivation of sarin-inhibited WT- and F338A-AChEs




Fig. 1. Plots of active site titration of PEGylated WT- and F338A-AChEs and aging of their adducts with soman. A, active site titration of PEGylated WT- and F338A-AChEs with soman and MEPQ. In all titration experiments, the concentration of the catalytic subunit of PEGylated WT- and F338A-AChEs was 120 to 170 nM. The residual activity was determined after incubation at 27°C in the presence of various concentrations of the different compounds for 60 min. B, time course of aging of PEGylated WT- and F338A-AChEs with soman. PEG-WT-AChE or PEG-F338A-AChE (100 nM) was incubated with soman (200 nM) at 27°C. HI-6 (0.5 mM) was added to samples, removed at various time points, and enzymatic activity (E_t) was determined after 120 min. Plots present the ln values of the fraction of reactivatable enzyme (E_t/E_0) versus enzyme/soman incubation times. Inset, expansion of the scale of time for the fast aging of PEG-WT-AChE.

TABLE 2

Apparent bimolecular rate constant for phosphorylation (k_i) of WT- and F338A AChEs

Values represents mean of triplicate determination with S.D. not exceeding 20%.

AChE Species	Phosphorylation (k_i)			
	Soman	Sarin	VX	Paraoxon
	$\times 10^{-4} \text{ M}^{-1} \text{ min}^{-1}$			
WT ^a	8600	2350	11,400	100
F338A ^a	9000	1450	2700	7
PEG-WT	7900	3100	8200	180
PEG-F338A	13,000	2700	3000	8

^a Values compiled from our previous studies (Ordentlich et al., 1993, 1998) are presented here for comparison.

proceeded at similar rates, with less than 4-fold and 2-fold differences in the k_r and k_2 values, respectively (Table 3). Similar results were observed for the VX-inhibited enzymes (less than 3-fold and 2-fold differences between the k_r and k_2 values of the WT- and F338A-AChEs). In all cases, the reactivation kinetic values of OP-inhibited PEGylated WT or PEGylated F338A enzymes were similar to those measured for their non-PEGylated counterparts (Table 3), attesting to the fact that PEG conjugation does not influence the reactivation process. It should be noted that there is an overall 10-fold decrease in the reactivation rates (k_r and k_2) of soman-inhibited F338A mutant compared with sarin/VX-inhibited enzymes (Table 3). This considerable difference in the reactivation kinetics values may be attributed to the branched alkyl group of soman, which can interfere with the ability of HI-6 to remove the phosphyl residue from the active site of the enzyme (Barak et al., 1997).

Taken together, measurement of rate of hydrolysis, inhibition kinetics, phosphorylation rates, stereoselectivity, and rates of reactivation of OP adducts allowed us to determine that neither the introduction of the F338A mutation nor the chemical modification of the enzyme by PEG conjugation deleteriously affected the bioscavenging properties of the enzyme. It should be noted that in many instances, the conjugation of PEG moieties to proteins was accompanied by a concomitant loss of their biological activity (Harris and Chess, 2003). We note that elaborate studies were indeed required to determine the exact conditions for performing efficient PEG conjugation without compromising the biological activity of ChEs (Cohen et al., 2001, 2007).

The only discernible difference between F338A-AChE and WT-AChE was their differential rate of aging. Controlled PEG conjugation of F338A-AChE did not affect its rate of aging, so that both the non-PEGylated and PEGylated versions of F338A-AChE are characterized by significantly lower rates of aging than WT-AChE. This low rate of aging may provide the enzyme with an advantageous trait, in terms of its ability to serve for prophylactic treatment of OP intoxication as demonstrated in the next section.

Reactivated PEG-F338A-AChE Effectively Protects Mice against Repeated OP Exposure

We next examined whether the slower aging rate of the F338A mutant (native and PEGylated form) allowed it to be more efficient than the WT-AChE in providing long-term protection against repeated OP exposure in general, and soman in particular. To this end, we set out to perform a series of experiments in which we determined the pharmacokinetic profile of PEG-WT- and F338A-AChEs and their

ability to serve for prophylaxis treatment against exposure to various OP agents. Finally, we compared the protective potential of these AChEs against repeated soman exposures in mice.

Pharmacokinetics of PEG-F338A-AChE. PEG-F338A-AChE, as well as its non-PEGylated counterpart, were administered i.v. to mice, and their pharmacokinetic profiles were determined (Fig. 2). PEG-F338A-AChE displayed a mean residence time (MRT) value of ~2800 min (Table 4), at least 20-fold greater than the MRT value exhibited by the nonmodified form of the enzyme (120 min). Thus, in terms of its pharmacokinetic performance, the PEGylated version of F338A-AChE displays an extended circulatory lifetime, similar to that determined previously for the PEGylated version of WT-AChE (Cohen et al., 2001; also shown in Fig. 2 for comparison).

Prophylactic Protection of Mice against OP Challenge.

We next assessed the ability of PEG-F338-AChE to confer protection to mice against OP compound intoxication. Mice were administered either PEG-F338A-AChE or PEG-WT-AChE, and then exposed to lethal doses of soman, sarin, or VX. In the past, we demonstrated that 1.2 nmol of PEG-WT-AChE is required for the effective protection of mice against 2.5 LD₅₀ of soman (=3 nmol soman/30-g mouse; Cohen et al., 2006). Administration of the same amount of PEG-F338-AChE protected all of the mice challenged immediately after enzyme administration with 2.5 LD₅₀ soman (Table 5). Likewise, PEG-F338-AChE was as effective as PEG-WT-AChE in its ability to confer protection against sarin or VX intoxication. Thus, mice administered with 2.6 or 3.2 nmol of PEG-F338-AChE, respectively, were fully protected against challenge of 2.5 LD₅₀ sarin (=6.5 nmol sarin/30-g mouse) or VX (=4 nmol VX/30-g mouse). In comparison with the AChE doses required for effective protection against soman or sarin intoxication (molar ratio of AChE:soman/sarin = 0.4:1), full protection against 2.5 LD₅₀ VX was achieved only when twice as high molar amounts of AChE were administered (molar ratio of AChE:VX = 0.8:1). This is due to the fact that AChE displays a marked stereoselectivity toward half of the enantiomers comprising racemic soman and sarin compounds, yet it interacts with both enantiomers of VX (Cohen et al., 2006; also see Fig. 1A). Taken together, these findings demonstrate that the PEG-F338A version of AChE is indeed as effective as PEG-WT-AChE in its ability to protect mice against OP compounds.

HI-6 Mediated Reactivation of Phosphorylated AChE Derivatives. As stated earlier, we hypothesized that the slow aging rate of the soman-inhibited PEG-F338A-AChE will allow restoration of its activity in the bloodstream by oxime reactivation, whereas the soman-inhibited WT-AChE will remain re-

TABLE 3

HI-6 reactivation constants of OP-inhibited WT- and F338A-AChEs

Values represents mean of triplicate determination with S.D. not exceeding 20%. K_r values were calculated from the ratio k_2/k_r .

AChE Species	Soman			Sarin			VX			Paraoxon		
	k_r	$k_2 \times 10^2$	$K_r \times 10^5$	k_r	$k_2 \times 10^2$	$K_r \times 10^5$	k_r	$k_2 \times 10^2$	$K_r \times 10^5$	k_r	$k_2 \times 10^2$	$K_r \times 10^5$
	$M^{-1} \min^{-1}$	\min^{-1}	M	$M^{-1} \min^{-1}$	\min^{-1}	M	$M^{-1} \min^{-1}$	\min^{-1}	M	$M^{-1} \min^{-1}$	\min^{-1}	M
WT	N.D.			11,800	36	3.1	14,700	20	1.4	170 ^a	6 ^a	35 ^a
F338A	300	2.3	7.8	3400	15	4.4	5900	10	1.7	80 ^a	0.6 ^a	7 ^a
PEG-WT	N.D.			13,000	28	2.1	5100	13	3.1	100	2.3	24
PEG-F338A	260	1.6	6.2	4200	10	2.3	4300	13	2.6	200	0.7	3.5

N.D., not determined.

^a Values compiled from our previous studies (Grosfeld et al., 1996) are presented here for comparison.

fractive to this treatment. To examine this, we first performed an in vitro experiment in which we measured the reactivation of soman-inhibited PEG-F338A-AChE in blood samples. To this end, heparinized blood samples were removed from mice and spiked with either PEG-WT-AChE or PEG-F338A-AChE (0.8 μ M). Soman was then added to the samples (final concentration, 1.6 μ M), and the reaction mixtures were incubated at 37°C for 20 min. Under these conditions, 90% of both enzymes were inactivated, as judged by their diminished ability to hydrolyze ATC (Fig. 3). HI-6 (1.5 mM) was then added, and the catalytic




Fig. 2. Circulatory clearance profiles of non-PEGylated and PEGylated F338A-AChEs. Non-PEGylated, PEGylated F338A-AChEs, and PEGylated WT-AChE (100 U) were administered i.v. to three mice each at levels that are at least 30-fold higher than background level. Blood samples withdrawn at various time points were assayed for ChE activity, and values were corrected for background hydrolytic activity in the blood. Values are presented as percentage of input ChE determined 1 min after enzyme administration.

TABLE 4

Pharmacokinetics parameters of F338A-AChE enzymes in mouse blood. Mice were intravenously injected with 100 U, presented data is average \pm S.D. of at least three mice for each enzyme.

	F338A-AChE	PEG-F338A-AChE
Distribution phase		
$t_{1/2\alpha}$ (min)	8 ± 2	46 ± 16
Area under the curve (%) ^a	15	1
Elimination phase		
$t_{1/2\beta}$ (min)	75 ± 8	2130 ± 110
Area under the curve (%) ^a	85	99
MRT (min)	120	2870

^a The percentage each term contributes to the area under the curve.

TABLE 5

Survival rates of PEGylated WT- and F338A-AChE pretreated mice following exposure to different OP agents

Animals were administered with 3 nmol/mouse AChE for soman challenge, 2.6 nmol/mouse AChE for sarin challenge, and 3.2 nmol/mouse AChE for VX challenge. Survival is presented as percentage surviving animals from a total of five animals.

Administered OP Agent	Administered AChE		
	None	PEG-WT	PEG-F338A
		%	
2.5 LD ₅₀ Soman ^a	0	100	100
2.5 LD ₅₀ Sarin ^a	0	100	100
2.5 LD ₅₀ VX	0	100	100

^a Animals were pretreated with CBDP 1 h before enzyme administration.

activity of the two enzymes was monitored in samples removed at various time points. As expected, the soman-inhibited WT-AChE remained unaffected throughout the experiment. In contrast, the inhibited F338A-AChE enzyme displayed a gradual increase in enzymatic activity, reaching 60% of input activity after 120 min of incubation in the presence of the oxime (Fig. 3). Thus, unlike in the case of WT-AChE, soman adducts of F338A-AChE remain susceptible to oxime-mediated reactivation in the blood samples, for extended periods.

To examine the ability to reactivate soman-inhibited F338A-AChE in vivo, mice were administered either with PEG-WT-AChE or PEG-F338A-AChE (4 nmol/mouse), and 1 min later they were exposed to the highest dose of soman (6.5 nmol/mouse; 5.4 LD₅₀) that is compatible with complete survival of the enzyme-administered mice. Measurement of AChE activity in blood samples removed 10 min later allowed us to determine that 70% of both enzyme forms were inactivated. HI-6 at a dose that is well tolerated, 50 mg/kg, was then administered i.v. to the mice, and enzyme reactivation was monitored in blood samples removed at various time points. In the WT-AChE-administered mice, the enzyme remained inhibited throughout the experiment, whereas in the F338A-AChE-administered mice, the level of circulating active enzyme increased by 67% within 10 min after HI-6 administration, after which enzymatic activity remained constant (50% of input enzyme) up to 1 h (Fig. 4A), probably due to the relatively rapid clearance of HI-6 from the circulation (Maxwell et al., 1993). Due to the elimination process of the PEGylated enzymes from the circulation, the levels of circulating PEGylated enzyme at the time of HI-6 administration were approximately 70% of input enzyme (Fig. 2), and therefore actually 50% of the inhibited PEG-F338A-AChE present at this time point can be expected to undergo reactivation by HI-6, which is in good agreement with the results obtained for in vitro reactivation (Fig. 3). We thereby examined whether the increment in circulating active enzyme brought




Fig. 3. In vitro reactivation of soman-inhibited PEGylated WT- and F338A-AChEs. PEGylated AChEs and soman (0.8 and 1.6 μ M, respectively) were added to heparinized blood samples removed from mice and incubated for 20 min at 37°C, after which HI-6 (1.5 mM) was added. AChE activity determined at various time points after the addition of HI-6 is expressed as percentage of AChE activity measured in blood samples spiked with AChE (0.8 μ M) only.

about by HI-6 reactivation of PEG-F338A-AChE may protect the mice against an additional soman exposure.

Protection Of Mice against Repeated Soman Exposure. To assess the ability of PEG-F338A-AChE to confer protection against repeated soman exposure, mice were administered PEGylated and non-PEGylated versions of both WT-AChE and F338A-AChE as described above, and then exposed to 5.4 LD₅₀ of soman. The mice, all of which survived this soman challenge (Fig. 4B, soman A) and displayed no more than mild symptoms of OP intoxication, were adminis-

tered HI-6 20 min later. After an additional 40 min, they were re-exposed to a second soman challenge of 4 LD₅₀. As expected, mice pretreated with the circulatory short-lived non-PEGylated versions of AChE succumbed to the second soman challenge (Fig. 4B, soman B) regardless of whether the WT or F338A version of the enzyme was administered. The PEGylated WT-AChE protected only 16% of the mice, even though this enzyme form exhibits long-term circulatory residence, so that at the time of the second soman challenge, 80% of the input enzyme still resides in the circulation (Fig. 2). In contrast, 100% of the PEG-F338A-AChE-pretreated mice survived the second soman challenge (Fig. 4B). Because the pharmacokinetics of PEG-F338A-AChE and PEG-WT-AChE are very similar (Fig. 2), the marked difference between the survival rates of the mice administered with these two enzyme forms necessarily stems from the differential ability of these enzymes to undergo reactivation by in vivo oxime treatment (Fig. 4A). Based on these findings, it indeed seems that the in vivo reactivation of soman-inhibited PEG-F338A-AChE, can be harnessed to provide effective protection against multiple exposures to OP compounds.

Discussion

The use of recombinant ChEs as a source for OP bioscavengers can serve not only to generate the large quantities of enzyme required for stoichiometric neutralization but also provide the means to incorporate favorable characteristics into the enzyme fold by genetic engineering. For example, amino acids involved in the aging of OP-ChE adducts, may be favorably altered by means of genetic engineering, to decelerate the irreversible inactivation of the bioscavenger. Extensive studies carried out in the past indeed allowed us to identify various amino acid residues that are involved in the aging of human AChE adduct with OPs. These include the Trp86 residue, a key element in the anionic binding subsite of the enzyme, Glu202, Glu450 and Tyr133 which are members of the H-bond network in the active center of the enzyme (Ordentlich et al., 1993; Shafferman et al., 1996), as well as Phe338, which is located within the aromatic patch lining the active center gorge of the enzyme (Ariel et al., 1998). Although replacement of any of these amino acids reduces the rate of aging of the HuAChE-OP adducts, only substitution of Phe338 results in a significantly lower rate of aging without compromising reactivity of the enzyme toward a wide variety of OPs (Shafferman et al., 1996; Table 2). Therefore, from a practical point of view, replacement of Phe338, but not of the other residues involved in aging, can be used for designing an AChE-based bioscavenger of OP compounds, displaying improved biochemical traits. Indeed, here we demonstrate that incorporation of the F338A mutation into PEGylated HuAChE decelerates aging, so that the soman-inhibited enzyme can be reactivated by oximes in vivo to an extent that enables mice to withstand repeated exposure to high doses of soman (e.g., 5.4 LD₅₀ and then 4 LD₅₀). It should be noted in this context, that attempts to generate a recombinant BChE-based slow-aging enzyme by replacement of residue Phe329, which is analogous to Phe338 in HuAChE, were not successful and had a very limited effect on the rate of aging of BChE (Masson et al., 1997).

An additional limitation in the exploitation of recombinant BChE as a bioscavenger of OP compounds relates to the




Fig. 4. In vivo reactivation and protective potential against repeated soman exposure of WT- and F338A-AChEs. Mice were pretreated with non-PEGylated or PEGylated WT- and F338A-AChEs (2 nmol/mouse i.v.; T = 0), exposed to soman (4 nmol/mouse i.v.; T = 1 min.), treated with HI-6 (50 mg/kg i.v.; T = 21 min.), and re-exposed to soman (4 nmol/mouse i.v.; T = 61 min.). A, AChE activity was monitored in blood samples removed at various time points from mice pretreated with the PEGylated versions of AChE. Activity is expressed as percentage of input values, determined immediately after administration of the exogenous PEG-AChEs. Times of AChE administration, first soman exposure (1st soman), HI-6 treatment, and second soman exposure (2nd soman) are indicated by arrows. B, percentage of survival after the first and second soman exposures of mice ($n = 6$) pretreated with the different AChE enzyme forms.

ability to extend its circulatory longevity. Recombinant ChEs are retained in the circulation of experimental animals for much shorter periods than native serum-resident ChEs (Kronman et al., 1995). We have shown in the past (Chitlaru et al., 2001) that recombinant AChE can be converted into a circulatory long-lived enzyme by the combined optimization of post-translation-related factors (sialic acid occupancy, enzyme tetramerization, and glycan loading). However, obtaining a homogenous long-lived circulatory scavenger through such a biotechnological/biochemical route may be difficult. In parallel, we demonstrated that conjugation of PEG to lysine residues of rHuAChE or rHuBChE, can effectively extend their circulatory longevity; however, the generation of a bioactive PEGylated ChE is dependent on a delicate balance between efficient PEG appendage and enzyme activity preservation, and therefore needs to be carried out in a highly controlled manner. For recombinant AChE, conditions for high level PEGylation without compromising enzymatic activity were determined (Cohen et al., 2001), and the introduction of a set of defined lysine replacements was shown recently to give rise to a homogenous uniformly PEGylated enzyme product (Cohen et al., 2007). In contrast, PEGylation of rHuBChE resulted in the generation of a heterogeneous set of products, which unlike PEGylated rHuAChE, were eliminated more rapidly than native serum-derived BChE (Chilukuri et al., 2005). This limitation may stem from sub-optimal PEGylation, which in turn may be related to BChE containing a high number of lysine residues (37 lysines) that may serve as target sites for PEG conjugation (Cohen et al., 2007). Reducing the number of BChE lysines by mutagenesis, to allow for the generation of a more homogenous PEG-BChE preparation without seriously affecting enzymatic properties, might be a much more challenging task than that achieved in AChE.

Finally, comparative protection experiments allowed us to determine that on a molar basis, AChE conferred a higher level of protection than BChE against soman and sarin intoxication because the latter, as opposed to the highly stereoselective AChE, is partially consumed by reaction with the nontoxic P_R enantiomers of soman and sarin (Cohen et al., 2006). If indeed 200 mg of BChE is required to protect humans against 2 LD₅₀ of soman, it seems that only 78 mg of AChE will be needed to confer the same extent of protection (Cohen et al., 2006). The ability to reduce the quantities of exogenously administered protein when using an AChE-based bioscavenger, and the relative facility to generate homogenous preparations of PEGylated enzyme of AChE source, serve as additional motives for preferring AChE over BChE as a therapeutic bioscavenger.

Altogether, we have tailored a specified form of human AChE that combines optimized kinetic and pharmacokinetic traits required for effective bioscavenging of OP compounds. By means of chemical conjugation of PEG to the AChE molecule, the pharmacokinetic properties of the enzyme can be improved so that the enzyme will reside in the circulation for sufficiently long periods, and thereby could be used for prophylactic treatment. The kinetic traits of the enzyme can also be tailored so that the enzyme will be able to cope more efficiently with OP-mediated intoxication. Specifically, substitution of Phe338 by alanine within the active center of the AChE molecule can significantly reduce the rate of irreversible dealkylation (aging) of the AChE-based bioscavenger,

without affecting the reactivity of the enzyme toward OP agents. It thus seems that, to date, the PEG-F338A-HuAChE is the most optimal OP bioscavenger engineered for prophylactic treatment against OP poisoning.

Acknowledgments

We thank our colleagues Drs. Dov Barak and Baruch Velan for valuable suggestions and careful reading of the manuscript. We thank Sharon Ehrlich and Shirley Lazar for excellent technical assistance.

References

- Ariel N, Ordentlich A, Barak D, Bino T, Velan B, and Shafferman A (1998) The "aromatic patch" of three proximal residues in the human acetylcholinesterase active center allows for versatile interaction modes with inhibitors. *Biochem J* **335**:95–102.
- Ashani Y, Grauer E, Grunwald J, Allon N, and Raveh L (1998) Current capabilities in extrapolating from animal to human the capacity of human butyrylcholinesterase to detoxify organophosphates, in *Structure and Function of Cholinesterases and Related Proteins* (Doctor BP, Quinn DM, Rotundo RL, and Taylor P eds), pp 255–260, Plenum Press, New York.
- Barak D, Kronman C, Ordentlich A, Ariel N, Bromberg A, Marcus D, Lazar A, Velan B, and Shafferman A (1994) Acetylcholinesterase peripheral anionic site degeneracy conferred by amino acid arrays sharing a common core. *J Biol Chem* **269**:6296–6305.
- Barak D, Ordentlich A, Segall Y, Velan B, Benschop HP, de Jong LP, and Shafferman A (1997) Carbocation-mediated processes in biocatalysts. Contribution of aromatic moieties. *J Am Chem Soc* **119**:3157–3158.
- Chilukuri N, Parikh K, Sun W, Naik R, Tipparaju P, Doctor BP, and Saxena A (2005) Polyethylene glycosylation prolongs the circulatory stability of recombinant human butyrylcholinesterase. *Chem Biol Interact* **157**:158:115–121.
- Chitlaru T, Kronman C, Velan B, and Shafferman A (2001) Effect of human acetylcholinesterase subunit assembly on its circulatory residence. *Biochem J* **354**:613–625.
- Cohen O, Kronman C, Chitlaru T, Ordentlich A, Velan B, and Shafferman A (2001) Effect of chemical modification of recombinant human acetylcholinesterase by polyethylene glycol on its circulatory longevity. *Biochem J* **357**:795–802.
- Cohen O, Kronman C, Velan B, and Shafferman A (2004) Amino acid domains control the circulatory residence time of primate acetylcholinesterases in rhesus macaques. *Biochem J* **378**:117–128.
- Cohen O, Kronman C, Raveh L, Mazor O, Ordentlich A, and Shafferman A (2006) Comparison of polyethylene glycol-conjugated recombinant human acetylcholinesterase and serum human butyrylcholinesterase as bioscavengers of organophosphate compounds. *Mol Pharmacol* **70**:1121–1131.
- Cohen O, Kronman C, Lazar A, Velan B, and Shafferman A (2007) Controlled concealment of exposed clearance and immunogenic domains by site-specific polyethylene glycol attachment to acetylcholinesterase hypolysine mutants. *J Biol Chem* **282**:35491–35501.
- Dawson RM (1994) Review of oximes available for treatment of nerve agents poisoning. *J Appl Toxicol* **14**:317–331.
- de Jong LP and Kossen SP (1985) Stereospecific reactivation of human brain and erythrocyte acetylcholinesterase inhibited by 1,2,2-trimethylpropyl methylphosphonofluoridate (soman). *Biochim Biophys Acta* **830**:345–348.
- de Jong LP, van Dijk C, and Benschop HP (1988) Hydrolysis of the four stereoisomers of soman catalyzed by liver homogenate and plasma from rat, guinea pig and marmoset and by human plasma. *Biochem Pharmacol* **37**:2939–2948.
- Ellman GL, Courtney KD, Andres V Jr, and Feather-Stone RM (1961) A new and rapid colorimetric determination of acetylcholinesterase activity. *Biochem Pharmacol* **7**:88–95.
- Fischer M, Ittah A, Liefer I, and Gorecki M (1993) Expression and reconstitution of biologically active human acetylcholinesterase from *E. Coli*. *Cell Mol Neurobiol* **13**:25–38.
- Fleisher JH and Harris LW (1965) Dealkylation as a mechanism for aging of cholinesterase after poisoning with pinacolyl methylphosphonofluoridate. *Biochem Pharmacol* **14**:641–650.
- Grosfeld H, Barak D, Ordentlich A, Velan B, and Shafferman A (1996) Interaction of oxime reactivators with diethylphosphoryl adducts of human acetylcholinesterase and its mutant derivatives. *Mol Pharmacol* **50**:639–649.
- Harris JM and Chess RB (2003) Effect of PEGylation on pharmaceuticals. *Nat Rev Drug Discov* **2**:214–221.
- Kronman C, Velan B, Gozes Y, Leitner M, Flashner Y, Lazar A, Marcus D, Sery T, Papier Y, and Grosfeld H (1992) Production and secretion of high levels of recombinant human acetylcholinesterase in cultured cell lines: microheterogeneity of the catalytic subunit. *Gene* **121**:295–304.
- Kronman C, Velan B, Marcus D, Ordentlich A, Reuveny S, and Shafferman A (1995) Involvement of oligomerization, N-glycosylation and sialylation in the clearance of cholinesterases from circulation. *Biochem J* **311**:959–967.
- Kronman C, Cohen O, Raveh L, Mazor O, Ordentlich A, and Shafferman A (2007) Polyethylene-glycol conjugated recombinant human acetylcholinesterase serves as an efficacious bioscavenger against soman intoxication. *Toxicology* **233**:40–46.
- Lenz DE, Yeung D, Smith JR, Sweeney RE, Lumley LA, and Cerasoli DM (2007) Stoichiometric and catalytic scavengers as protection against nerve agent toxicity: a mini review. *Toxicology* **233**:31–39.
- Levy D and Ashani Y (1986) Synthesis and in vitro properties of a powerful quater-

- nary methylphosphonate inhibitor of acetylcholinesterase. A new marker in blood-brain barrier research. *Biochem Pharmacol* **35**:1079–1085.
- Masson P, Adkins S, Gouet P, and Lockridge O (1993) Recombinant human butyrylcholinesterase G390V, the fluoride-2 variant, expressed in Chinese hamster ovary cells, is a low affinity variant. *J Biol Chem* **268**:14329–14341.
- Masson P, Fortier PL, Albaret C, Froment MT, Bartels CF, and Lockridge O (1997) Aging of di-isopropyl-phosphorylated human butyrylcholinesterase. *Biochem J* **327**:601–607.
- Maxwell DM, Brecht KM, Doctor BP, and Wolfe AD (1993) Comparison of antidote protection against soman by pyridostigmine, HI-6 and acetylcholinesterase. *J Pharmacol Exp Ther* **264**:1085–1089.
- Millard CB, Koellner G, Ordentlich A, Shafferman A, Silman I, and Sussman JL (1999) Reaction products of acetylcholinesterase and VX reveal a mobile histidine in the catalytic triad. *J Am Chem Soc* **121**:9883–9884.
- Ordentlich A, Kronman C, Barak D, Stein D, Ariel N, Marcus D, Velan B, and Shafferman A (1993) Engineering resistance to “aging” of phosphorylated human acetylcholinesterase. Role of hydrogen bond network in the active center. *FEBS Lett* **334**:215–220.
- Ordentlich A, Barak D, Kronman C, Ariel N, Segall Y, Velan B, and Shafferman A (1996) The architecture of human acetylcholinesterase active center probed by interactions with selected organophosphate inhibitors. *J Biol Chem* **271**:11953–11962.
- Ordentlich A, Barak D, Kronman C, Ariel N, Segall Y, Velan B, and Shafferman A (1998) Functional characteristics of the oxyanion hole in human acetylcholinesterase. *J Biol Chem* **273**:19509–19517.
- Ordentlich A, Barak D, Kronman C, Benschop HP, De Jong LP, Ariel N, Barak R, Segall Y, Velan B, and Shafferman A (1999) Exploring the active center of human acetylcholinesterase with stereoisomers of an organophosphorous inhibitor with two chiral centers. *Biochemistry* **38**:3055–3066.
- Ordentlich A, Barak D, Sod-Moriah G, Kaplan D, Mizrahi D, Segall Y, Kronman C, Karton Y, Lazar A, Marcus D, et al. (2005) The role of AChE active site gorge in determining stereoselectivity of charged and noncharged VX enantiomers. *Chem Biol Interact* **157–158**:191–198.
- Radić Z, Pickering NA, Vellom DC, Camp S, and Taylor P (1993) Three distinct domains in the cholinesterase active center allows for versatile interaction modes with inhibitors. *Biochemistry* **32**:12074–12084.
- Raveh L, Grunwald J, Marcus D, Papier Y, Cohen E, and Ashani Y (1993) Human butyrylcholinesterase as a general prophylactic antidote for nerve agent toxicity. In vitro and in vivo quantitative characterization. *Biochem Pharmacol* **45**:2465–2474.
- Raveh L, Grauer E, Grunwald J, Cohen E, and Ashani Y (1997) The stoichiometry of protection against soman and VX toxicity in monkeys pretreated with human butyrylcholinesterase. *Toxicol Appl Pharmacol* **145**:43–53.
- Shafferman A, Ordentlich A, Barak D, Stein D, Ariel N, and Velan B (1996) Aging of phosphorylated human acetylcholinesterase: catalytic processes mediated by aromatic and polar residues of the active center. *Biochem J* **318**:833–840.
- Shafferman A, Barak D, Kaplan D, Ordentlich A, Kronman C, and Velan B (2005) Functional requirements for the optimal catalytic configuration of the AChE active center. *Chem Biol Interact* **157–158**:123–131.
- Sussman JL, Harel M, Frolow F, Oefner C, Goldman A, Tokor L, and Silman I (1991) Atomic structure of acetylcholinesterase from *Torpedo californica*: a prototypic acetylcholine-binding protein. *Science* **253**:872–879.
- Worek F, Thiermann H, Szinicz L, and Eyer P (2004) Kinetic analysis of interactions between human acetylcholinesterase, structurally different organophosphorus compounds and oximes. *Biochem Pharmacol* **68**:2237–2248.

Address correspondence to: Dr. Avigdor Shafferman, Israel Institute for Biological Research, Ness-Ziona, 74100, Israel. E-mail: avigdors@iibr.gov.il