

The Translesion Polymerase Rev3L in the Tolerance of Alkylating Anticancer Drugs

Wynand Paul Roos, Anastasia Tsaalbi-Shtylik, Roman Tsaryk, Fatma Güvercin, Niels de Wind, and Bernd Kaina

Institute of Toxicology, University Medicine Mainz, Mainz, Germany (W.P.R., R.T., F.G., B.K.); and Department of Toxicogenetics, Leiden University Medical Center, Leiden, Netherlands (A.T.S., N.d.W.)

Received May 27, 2009; accepted July 27, 2009

ABSTRACT

Temozolomide and fotemustine, representing methylating and chloroethylating agents, respectively, are used in the treatment of glioma and malignant melanoma. Because chemoresistance of these tumors is a common phenomenon, identification of the underlying mechanisms is needed. Here we show that Rev3L, the catalytic subunit of the translesion DNA polymerase ζ , mediates resistance to both temozolomide and fotemustine. Rev3L knockout cells are hypersensitive to both agents. It is remarkable that cells heterozygous for Rev3L showed an intermediate sensitivity. Rev3L is not involved in the tolerance of the toxic O^6 -methylguanine lesion. However, a possible role of Rev3L in the tolerance of O^6 -chloroethylguanine or the subsequently formed N1-guanine-N3-cytosine interstrand cross-link

is shown. Rev3L had no influence on base excision repair (BER) of the *N*-alkylation lesions but is very likely to be involved in the tolerance of *N*-alkylations or apurinic/apyrimidinic sites originating from them. We also show that Rev3L exerts its protective effect in replicating cells and that loss of Rev3L leads to a significant increase in DNA double-strand breaks after temozolomide and fotemustine treatment. These data show that Rev3L contributes to temozolomide and fotemustine resistance, thus acting in concert with O^6 -methylguanine-DNA methyltransferase, BER, mismatch repair, and double-strand break repair in defense against simple alkylating anticancer drugs.

Alkylating agents are widely used in cancer therapy, and chemotherapy often fails because of the development of a drug-resistant tumor phenotype. Because alkylating agents primarily target DNA, DNA repair and damage tolerance mechanisms are on the forefront in the development of drug resistance. The chemotherapeutics temozolomide and fotemustine belong to the groups of methylating and chloroethylating agents, respectively. They are used for the treatment of brain tumors (astrocytomas, glioblastoma multiforme) and malignant melanoma as well as brain metastasis of other tumors. Chloroethylating agents such as carmustine are also part of the combination therapy for the treatment of Hodgkin's lymphomas, lung cancer, and breast cancer.

Methylating and chloroethylating drugs induce a broad spectrum of DNA adducts. One of them, O^6 -alkylguanine, has

been identified to be a major killing lesion that has a high propensity in activating the apoptotic cell death pathway (Roos and Kaina, 2006). This damage is repaired by the suicide repair enzyme O^6 -methylguanine-DNA methyltransferase (MGMT), which attenuates or even completely abolishes the toxic effects provoked by O^6 -alkylguanine. Therefore, MGMT is considered to be a key node in the defense against O^6 -alkylating agents (Kaina et al., 2007).

Other targets of O^6 -alkylating agents are the *N*-residues of purines and pyrimidines in DNA. Most of the *N*-alkylpurines such as N7-alkylguanine, N3-alkyladenine, and N3-alkylguanine are repaired by base excision repair (BER). Defects in BER sensitize cells to the killing effects of O^6 -alkylating agents. Thus, cells lacking DNA polymerase β (Pol β) are hypersensitive to *N*-methyl-N'-nitro-N-nitrosoguanidine (MNNG), temozolomide, and chloroethylnitrosoureas (Sobol et al., 1996), and this hypersensitivity is executed by apoptosis (Ochs et al., 2002). Cells defective in poly(ADP-ribose)polymerase (de Murcia et al., 1997) or X-ray repair cross-complementing group 1 (XRCC1) (Zdzienicka et al., 1992), which coordinate the last steps of BER, are also hypersensitive, and pharmacological inhibition of BER

This work was supported by Deutsche Forschungsgemeinschaft [Grant DFG-Ka 724]; the German Cancer Foundation [Grant Ka-106748]; the Dutch Cancer Foundation [Grant UL 2001-2517]; and the European Union [Grant EU-IP FP6-512113].

Article, publication date, and citation information can be found at <http://molpharm.aspetjournals.org>.
doi:10.1124/mol.109.058131.

ABBREVIATIONS: MGMT, O^6 -methylguanine-DNA methyltransferase; BER, base excision repair; DSB, DNA double-strand break; ssb, single-strand break; TLS, translesion DNA synthesis; O^6 BG, O^6 -benzylguanine; MNNG, *N*-methyl-N'-nitro-N-nitrosoguanidine; Pol, polymerase; DMSO, dimethyl sulfoxide; AP, apurinic/apyrimidinic; PBS, phosphate-buffered saline.

by methoxyamine, which blocks the repair of apurinic/apyrimidinic (AP) sites, enhances the killing effects of simple alkylating agents (Taverna et al., 2001; Liu et al., 2003). All of this supports the view that deficiency or down-regulation of BER, or imbalance in BER enzymatic steps, contribute to alkylating drug sensitization.

Unrepaired DNA alkylation damage interferes with DNA replication and, thus, gives rise to critical secondary DNA damage. The primary lesions, however, can be tolerated by translesion DNA synthesis (TLS), which prevents cells from becoming blocked irreversibly in the S- or G₂-phase of the cell cycle. One of the key translesion polymerases is DNA polymerase ζ (Pol ζ), the catalytic subunit of which is Rev3L (Rattray and Strathern, 2003). Rev3L recently has been shown to operate along with other translesion DNA polymerases to bypass different DNA lesions, including AP sites (Shachar et al., 2009). Although Rev3L is expressed to similar levels in lung, gastric, colon, and renal tumors compared with normal tissue (Kawamura et al., 2001) it has recently been found to be overexpressed in human gliomas compared with normal brain tissue (Wang et al., 2009).

Here we studied the response of cells lacking Rev3L to the cytotoxic effects of temozolomide and fotemustine. We show that *Rev3L* knockout cells are remarkably sensitive to these agents, undergoing apoptosis at high level. We also show that BER and MGMT are not involved in this phenotype, which suggests that Rev3L is a new player in alkylating drug resistance and is a target for new potential anticancer drugs. In addition, we hypothesize that drug-induced overexpression of Rev3L may mediate alkylating drug resistance in human gliomas.

Materials and Methods

Drugs and Drug Treatment. Temozolomide (Schering-Plough, Kenilworth, NJ) stocks (35 mM) were prepared by dissolving the drug in dimethyl sulfoxide (DMSO) and diluting it with sterile distilled H₂O (2-fold dilution). The temozolomide stocks were aliquoted and stored at -80°C until use. Fotemustine [diethyl-1-[3-(2-chloroethyl)-3-nitrosourea-ido]ethylphosphate; Muphoran, Servier Research International, Neuilly sur Seine, France] was prepared fresh for each treatment at a concentration of 1 $\mu\text{g}/\mu\text{l}$ in ethanol. In all cases, exponentially growing cells were treated with fotemustine or temozolomide. Once added to the medium, temozolomide has a half-life of 1.81 h, whereas fotemustine has an even shorter half-life (Tapiero et al., 1989; Denny et al., 1994). Therefore the treatment of cells can be seen as a pulse treatment. Time points indicated in figures refer to the time after the addition of the drug to the medium unless stated otherwise. *N*-methyl-*N'*-nitro-*N*-nitrosoguanidine (MNNG; Sigma, Munich, Germany) stocks were prepared by dissolving the drug in DMSO and then diluting with sterile distilled H₂O to a 10 mM concentration (100-fold dilution). To deactivate the MGMT protein, 10 μM *O*⁶-benzylguanine (*O*⁶BG) was added to the cells 1 h before drug treatment.

Cells and Culture Conditions. *Rev3L* wild-type, *Rev3L* heterozygote, and *Rev3L* knockout mice were obtained as follows. *Rev3L*(+/-) p53(+/-) C57BL/6 mice (Van Sloun et al., 2002) were mated to *Rev3L*(+/-) 129/OLA mice. *Rev3L*(-/-) p53(+/-) embryos that were 13.5 days post coitum were trypsinized and cultured until spontaneous immortalization, at very low frequency, occurred. Immortalized clones were isolated and displayed loss of the wild-type p53 allele, as determined by allele-specific polymerase chain reaction. p53 is a necessary but not sufficient factor for immortalization of these cells (Zander and Bemark, 2004). It was also shown that loss of p53 does not rescue the embryonic lethality of *Rev3L*-deficient embryos (Van Sloun et al., 2002). Cells were routinely grown in

Dulbecco's modified Eagle's medium containing 10% fetal calf serum at 37°C and 5% CO₂.

Determination of Cell Survival. Cell viability experiments were performed using the WST assay according to the manufacturer's protocol (Roche Diagnostics, Indianapolis, IN). Cell viability in untreated controls was set to 100%. Colony assays were performed on monolayers, growing in log phase. Cells were seeded in duplicate at appropriate cell numbers in 60-mm Petri dishes to yield approximately 100 surviving colonies. After 16 h, when cells were attached, *O*⁶BG (10 μM) was added for 1 h, and then the cells were exposed to graded doses of temozolomide or fotemustine. After 10 to 14 days, cultures were fixed (in acetic acid/methanol/H₂O 1:1:8), stained (in 0.01% amido black), and colonies were counted. Results are from three independent experiments.

Determination of Apoptosis by Flow Cytometry. The amount of apoptosis induced by temozolomide or fotemustine was determined by quantifying the fraction of cells containing a subdiploid amount of DNA. Both adherent and detached cells were collected and suspended in 70% ethanol for fixation. Immediately before analysis, cells were treated with RNase (0.03 mg/ml) and subsequently stained with propidium iodide (16.5 mg/ml) in phosphate-buffered saline (PBS). Propidium iodide fluorescence was measured by flow cytometry (FACScalibur; BD Biosciences, Heidelberg, Germany). For each sample, 10,000 cells were analyzed (WinMDI Software; <http://facs.scripps.edu/software.html>).

Preparation of Protein Extracts. Cell pellets were suspended in buffer (10 mM HEPES-KOH, pH 7.4, 0.1 mM EDTA, 1 mM ethylene glycol-bis (b-aminoethyl ether), 250 mM sucrose, 1 μM Na₃VO₄, 0.5 mM phenylmethylsulfonyl fluoride, and 10 mM dithiothreitol). The cells were lysed by freeze/thaw/vortexing. The lysate was then centrifuged at 10,000 rpm for 10 min, and the supernatant containing the cytoplasmic proteins was used for Western blot analysis. Cell extracts for MGMT activity assay were prepared as follows. Cells were harvested and homogenized by sonication (Branson SONIFIER cell disruptor B15; Branson Ultrasonics Corporation, Danbury, CT) in buffer containing 20 mM Tris-HCl, pH 8.5, 1 mM EDTA, 1 mM β -mercaptoethanol, 5% glycerol, and the protease inhibitor phenylmethylsulfonyl fluoride (0.1 mM). The extract was centrifuged at 10,000 rpm (for 10 min) in the cold to remove debris, and the supernatant was snap-frozen in aliquots using liquid nitrogen and stored at -80°C until use. The protein concentration was determined by the Bradford method (Bradford, 1976).

Western Blot Analysis. The method used here is based on the method described previously (Renart et al., 1979). Protein (30 μg) of cytoplasmic extracts of temozolomide- or fotemustine-treated cells was separated in a 12% SDS polyacrylamide gel. Thereafter, proteins were transferred onto a nitrocellulose membrane (Protran; Schleicher & Schuell, Dassel, Germany). Membranes were blocked in 5% (w/v) fat-free milk powder in Tris-buffered saline containing 0.1% Tween 20, incubated with the primary antibody (1:1000 dilution), washed with 0.1% Tween 20 in Tris-buffered saline, and incubated with a horseradish peroxidase-coupled secondary antibody (1:3000) (GE Healthcare, München, Germany). Antibodies used were anti-extracellular signal-regulated kinase-2 (Santa Cruz Biotechnology Inc., Santa Cruz, CA), anti-caspase-7 (Cell Signaling Technology, Inc., Danvers, MA), and anti-caspase-3 (Cell Signaling). After final washing, blots were developed by using a chemiluminescence detection system (GE Healthcare).

MGMT Activity Assay. The MGMT activity in *Rev3Lwt*, *Rev3L*(+/-), and *Rev3L*(-/-) cells was determined using a method based on the radioactive assay in which tritium-labeled methyl groups are transferred from the *O*⁶-position of guanine to protein in the cell extract (Preuss et al., 1995). HeLa S3 cells expressing MGMT (588 \pm 86 fmol/mg protein) and HeLa MR cells deficient in MGMT served as positive and negative controls, respectively. The radioactivity of the protein was then measured. For each assay, cell extracts containing 200 μg of protein was incubated with [³H]methyl-nitrosourea-labeled calf thymus DNA containing *O*⁶MeG (total, 80,000

cpm/sample) in 700 mM HEPES-KOH, pH 7.8, 10 mM dithiothreitol, and 50 mM EDTA for 90 min. Data are expressed as femtomoles of radioactivity transferred from ^3H -labeled DNA to protein per milligram of protein within the sample. The MGMT activity data presented are results obtained from three independent experiments.

Single-Cell Gel Electrophoresis. DNA single-strand breaks were determined and quantified by the highly sensitive alkaline single-cell gel electrophoresis or comet assay as described previously (Olive et al., 1990). Agarose-imbedded MNNG pulse-treated cells were lysed (2.5 M NaCl, 100 mM EDTA, 10 mM Tris 1% sodium lauryl sarcosinate, pH 10, 1% Triton X-100, and 10% DMSO) and subjected to electrophoresis. Analysis of DNA migration was done by staining DNA with ethidium bromide and using the image analysis system Kinetic Imaging Komet 4.0.2 (OptiLas Systems, Evry, France). The mean tail moment (defined as the percentage of DNA in the tail \times tail length) of 50 cells per sample was determined.

Serum Starvation. *Rev3Lwt* and *Rev3L(-/-)* cells were plated in medium containing the indicated concentrations of serum. After 72 h of incubation, cells were trypsinized and counted. The number of cells obtained in 10% serum was set to 100%, and all other cell numbers obtained at lower serum concentrations were expressed as a fraction of this number.

Immunofluorescence Labeling and Microscopy. *Rev3Lwt* and *Rev3L(-/-)* cells were seeded on coverslips. After treatment with 0.5 mM temozolomide or 5 $\mu\text{g}/\text{ml}$ fotemustine and 24-h incubation for temozolomide-treated cells or 48-h incubation for fotemus-

tine-treated cells, the cells on the coverslips were fixed with 4% formaldehyde. A second fixation step was performed using 100% methanol (-20°C , 20 min). Cells and coverslips were then blocked in 5% bovine serum albumin PBS (0.3% Triton X-100). The antibodies used were anti- γH2AX (Millipore, Schwalbach, Germany) and Alexa Fluor 546 (Invitrogen, Karlsruhe, Germany). Just before mounting, DNA was stained with 100 nM 4,6-diamidino-2-phenylindole for 15 min. Between all steps, cells were washed in PBS (0.3% Triton X-100) for 5 min. Slides were mounted in antifade medium (glycerol/PBS 1:1, 2.5% 1,4-diazabicyclo[2.2.2]octane, pH 8.6, with HCl). For all time points, at least 40 nuclei were scored for foci. All experiments were repeated two times. Foci were not scored in apoptotic cells.

Results and Discussion

Rev3L Knockout Cells Are Hypersensitive to Temozolomide and Fotemustine. This study was aimed at elucidating whether Pol ζ is involved in the defense against DNA damage induced by temozolomide and fotemustine, which represent methylating and chloroethylating anticancer drugs, respectively. We used *Rev3L* knockout cells that lack the catalytic subunit of Pol ζ and compared isogenic wild-type (*Rev3Lwt*), heterozygous [*Rev3L(+/-)*], and homozygous [*Rev3L(-/-)*] knockout cells. Temozolomide (Fig. 1A) and fotemustine (Fig. 1B) caused a dose-dependent decrease in the viability of the




Fig. 1. Contribution of Rev3L to the protection of cells against temozolomide and fotemustine. Viability of temozolomide- (A) and fotemustine- (B) treated *Rev3Lwt*, *Rev3L(+/-)*, and *Rev3L(-/-)* cells. Cells were treated with the indicated doses, and viability was determined 72h later using the WST assay. Colony survival after temozolomide (C) and fotemustine (D) treatment in *Rev3Lwt*, *Rev3L(+/-)*, and *Rev3L(-/-)* cells. Cells were treated with the indicated concentrations, and colonies were scored after appearance 10 to 12 days later. For all experiments, to deplete MGMT, 10 μM O 6 BG was added to the medium 1 h before temozolomide or fotemustine treatment.

cells, and *Rev3L* deficiency caused a clear sensitization to these agents (Fig. 1, A and B). It is noteworthy that cells heterozygous for *Rev3L* showed an intermediate sensitivity toward temozolomide and fotemustine. Similar results were obtained with the more sensitive colony survival assay (Fig. 1, C and D).

Viability as measured by the WST assay and colony survival reflects both cell cycle arrest and cell death. To specifically address cell killing, the apoptotic response of these cells was determined after temozolomide and fotemustine treatment. Again, *Rev3L* knockout cells were hypersensitive to both temozolomide (Fig. 2A) and fotemustine (Fig. 2B). Similar to what was observed in the viability and colony assays, the *Rev3L* heterozygous cells showed an intermediate level of apoptosis compared with wild-type and *Rev3L*($-/-$) cells after treatment with both temozolomide and fotemustine (Fig. 2, A and B). To verify that what was observed by sub- G_1 flow cytometry was truly apoptosis, Western blot analyses of temozolomide- and fotemustine-treated cells were performed. Temozolomide and fotemustine caused the activation of both caspase-3 and caspase-7 in *Rev3L*($-/-$) cells, as revealed by the activated fragments (Fig. 2, C and D). Collectively, the data show that *Rev3L* protects cells against cytotoxicity induced by temozolomide and fotemustine. It is noteworthy that the intermediate phenotype of the *Rev3L*($+/-$) cells in the survival and apoptosis assays suggests that the expression level of *Rev3L* may be rate-limiting in the tolerance of temozolomide and fotemustine.

Rev3L Does Not Sensitize Cells to O^6 -Methylguanine. In previous work, it was shown that O^6 -methylguanine is the dominant apoptotic lesions induced by methylating drugs and that MGMT is therefore a key node in cell resistance to these agents (Kaina et al., 2007). O^6 -Methylguanine is repaired by MGMT in a one-step damage-reversal reaction that causes protection against the killing effect of this lesion. To test the hypothesis of whether *Rev3L*($-/-$) hypersensitiv-

ity is due to a lack of bypass of O^6 -methylguanine adducts, the influence of MGMT on the sensitivity of these cells was determined. The cell lines used display different levels of MGMT activity, from 5 up to 130 fmol/mg protein (Fig. 3A). The reason for the different expression levels is unknown, although it is well established that immortalized cells display variable levels of MGMT (Harris et al., 1996). To inactivate MGMT, the cell lines were treated with the specific inhibitor O^6 BG (Moschel et al., 1992). Thereafter, *Rev3Lwt* and *Rev3L*($-/-$) cells were treated with 0.5 mM temozolomide (in the presence or absence of 10 μ M O^6 BG), and the frequency of apoptosis was determined 72 h later (Fig. 3B). *Rev3L*($-/-$) cells were significantly more sensitive to temozolomide, and no difference in their apoptotic response was observed when they were pretreated with O^6 BG to inactivate MGMT (Fig. 3B).

The cytotoxicity of O^6 -methylguanine in the presence of O^6 BG depends on its miscoding properties during replication, followed by lethal processing by DNA mismatch repair (Karran and Bignami, 1992). Therefore, we anticipated that, if O^6 -methylguanine sensitivity is mediated by misincorporation opposite O^6 -methylguanine by *Rev3L*, *Msh2* deficiency would fully rescue the sensitivity of *Rev3L*-deficient mouse embryonic fibroblasts to temozolomide. However, sensitivity of *Rev3L* knockout mouse embryonic fibroblasts to the temozolomide analog MNNG was rescued only partially by concomitant *Msh2* deficiency (Fig. 3C). This result indicates that DNA mismatch repair is not involved in damage-processing provoked by a lack of *Rev3L*. Because the extent of rescue of MNNG sensitivity by *Msh2* deficiency was identical for *Rev3L* knockout cells as for wild-type cells (Fig. 3C) we also infer that O^6 -methylguanine is not subject of TLS by Pol ζ . The data rather suggest that Pol ζ is required for the tolerance of *N*-alkylation lesions or gaps that arise from spontaneous depurination of lesions such as N3-methyladenine or N3-methylguanine, or from BER intermediates like AP sites. In line with this is the observation that




Fig. 2. Activation of apoptosis and the executing caspases after temozolomide and fotemustine. Apoptosis induced by temozolomide (A) and fotemustine (B) in *Rev3Lwt*, *Rev3L*($+/-$), and *Rev3L*($-/-$) cells. Cells were treated with the indicated concentrations, and apoptosis was determined 72 h later by the sub- G_1 method using flow cytometry. The activated fragments of caspase-3 (C) and caspase-7 (D) were assayed in *Rev3Lwt* and *Rev3L*($-/-$) cells using immunoblotting at indicated times after 0.5 mM temozolomide or 5 μ M fotemustine treatment. For all experiments 10 μ M O^6 BG was added to the medium 1 h before temozolomide or fotemustine treatment.

Rev3L($-/-$) cells are also hypersensitive to methyl methanesulfonate (Okada et al., 2005; Takenaka et al., 2006; Wittschleben et al., 2006), which produces very low amounts of O^6 MeG (Beranek, 1990). Inhibition of BER (e.g., by methoxyamine) was shown to ameliorate the killing response of cells to temozolomide (Taverna et al., 2001), which was taken to dem-

onstrate that nonrepaired *N*-methylation lesions contribute to temozolomide-induced cytotoxicity, and BER protects against it. Here, we extend this finding showing that a translesion DNA polymerase, Pol ζ , contributes to resistance against temozolomide. Therefore, Pol ζ seems to be a potential target for anticancer chemotherapy.




Fig. 3. Influence of Rev3L on O^6 -alkylguanine triggered apoptosis and BER. A, MGMT repair activity in Rev3Lwt, Rev3L $+/+$, and Rev3L $-/-$ cells. B, apoptosis induced by temozolomide (0.5 mM) or fotemustine (10 μ g/ml) in the presence or absence of the MGMT inhibitor O⁶BG (10 μ M). Results obtained by sub-G1 determination using flow cytometry for Rev3Lwt and Rev3L $-/-$ are shown. Apoptosis frequency was determined 72 h after treatment. C, survival of Rev3Lwt, Msh2 $-/-$, Rev3L $-/-$, and Rev3L; Msh2 $-/-$ double-knockout cells after 250 μ M MNNG treatment. D, single-cell gel electrophoresis results obtained in Rev3Lwt and Rev3L $-/-$ cells treated for 1 h with indicated doses of MNNG. E, time course of ssb for Rev3Lwt and Rev3L $-/-$ cells pulse-treated for 1 h with 25 μ M MNNG; samples were stopped at the indicated time points.

Rev3L Sensitizes Cells to O^6 -Chloroethylguanine. Although O^6 BG did not affect on the killing response of temozolomide in *Rev3L*($-/-$) cells, it clearly ameliorated cell death by apoptosis upon treatment with fotemustine in *Rev3L*-lacking cells (Fig. 3B). This indicates that O^6 -chloro-

ethylguanine adducts are a substrate for Pol ζ -mediated TLS. It was shown recently that Pol ζ is involved in TLS of many different lesions, including mitomycin C, methyl methanesulfonate, benzo[*a*]pyrene adducts, AP sites, cisplatin-induced guanine-guanine cross-links, 4-hydroxyequilenin-cytosine,




Fig. 4. Replication dependence of apoptosis induction in *Rev3L*($-/-$) cells. A, relative growth in relation to serum concentration in *Rev3L*($-/-$) cells. Cells were grown in medium containing indicated concentrations of serum, and growth was determined by counting cell numbers after 72 h and plotting it as a relative number compared with cell number obtained in 10% serum. B, relative apoptotic response of *Rev3L*($-/-$) cells. Results were obtained by treating cells grown in the indicated concentrations of serum with 0.5 mM temozolomide or 5 μ g/ml fotemustine and then determining the apoptotic response 72 h later using sub-G1. The amount of apoptosis observed in 10% serum was set to 100%. C, γ H2AX foci formation in *Rev3L*wt cells treated with 0.5 mM temozolomide. D, quantification of γ H2AX foci formation in *Rev3L*wt and *Rev3L*($-/-$) cells after 0.5 mM temozolomide or 5 μ g/ml fotemustine treatment. For temozolomide-treated cells, foci were scored 24 h after treatment, and for fotemustine, 48 h after treatment. Foci were not scored in apoptotic cells. E, γ H2AX formation in *Rev3L*wt, *Msh2*($-/-$), *Rev3L*($-/-$), and *Rev3L**Msh2*($-/-$) double-knock-out cells after 100 μ M MNNG treatment at indicated time points.

and thymine-thymine 6-4 photoproducts (Shachar et al., 2009). Our data indicate that Polζ may also be involved in TLS across *O*⁶-chloroethylguanine. It is noteworthy that *O*⁶-chloroethylguanine is a rather unstable adduct, undergoing rearrangement to N1-*O*⁶-etheno-guanine and finally N1-guanine-N3-cytosine interstrand cross-links. Whether *O*⁶-chloroethylguanine or the etheno adduct derived from it is a subject of TLS by Polζ remains to be determined.

Rev3L Does Not Have an Influence on the Formation of Base Excision Repair Intermediates. Most of the DNA lesions induced by alkylating agents are repaired by BER, and BER-defective cells (e.g., Polβ knockout cells) are hypersensitive to methylating agents such as MNNG (Sobol et al., 1996). They also display a high level of DNA repair intermediates, which can be detected in the alkaline comet assay (Ochs et al., 2002). To elucidate the level of BER intermediates in *Rev3Lwt* and *Rev3L(-/-)* cells, cells were treated with a pulse of MNNG, and single-strand breaks (ssb) were determined. As shown in Fig. 3D, the frequency of ssb increased with the dose of MNNG. There was no difference in the tail moment between *Rev3Lwt* and *Rev3L(-/-)* cells. Figure 3E shows the decrease in the tail moment level with postincubation time. Again, no difference was observed between *Rev3Lwt* and *Rev3L(-/-)* cells. The data suggest that BER is not different in wild-type and *Rev3L* knockout cells, and Rev3L has no impact on the formation of ssb after treatment with methylating agents.

Rev3L Knockout Causes Replication-Dependent Toxicity and the Increased Formation of DNA Double-Strand Breaks. Temozolomide and fotemustine are used as chemotherapeutic agents in the treatment of cancer. Cancer cells are replicating and, therefore, it is important to know whether replicating cells will be targeted. To this end, *Rev3L(-/-)* cells were grown under conditions of different serum concentrations, and relative cell growth was determined. As the serum concentration increased, a significant increase in cell proliferation was observed after incubation for 72 h (Fig. 4A). When determining the influence of proliferation on apoptosis induced by either 0.5 mM temozolomide or 5 μg/ml fotemustine, an increase in apoptosis was observed with the increasing proliferation rate (Fig. 4B). The data clearly show that the killing effect of both temozolomide and fotemustine is strongly dependent on cell proliferation in *Rev3L(-/-)* cells. Because *Rev3Lwt* cells were quite resistant to the doses of temozolomide and fotemustine used, we can also infer from the data that Rev3L specifically protects proliferating cells from DNA damage induced by temozolomide and fotemustine.

If Rev3L is responsible for the bypass of temozolomide and fotemustine lesions during DNA synthesis in S-phase, then the lack of Rev3L may lead to more replication blocks that could collapse and form DNA double-strand breaks (DSBs) (Van Sloun et al., 2002). DSBs are effective triggers of apoptosis (Lips and Kaina, 2001). To determine whether *Rev3L(-/-)* cells show more DSBs than *Rev3Lwt* cells, both cell lines were treated with 0.5 mM temozolomide or 5 μg/ml fotemustine and the formation of γH2AX foci, a very good indicator of DSBs, was determined using fluorescence microscopy (for an example of γH2AX foci formation after chemotherapeutic treatment, see Fig. 4C). These foci were scored and plotted. In both temozolomide- and fotemustine-treated cells, *Rev3L(-/-)* showed significantly higher levels

of foci formation than the *Rev3Lwt* cells (Fig. 4D). Consistent with what was observed in the survival assays, loss of Msh2 was not able to correct for DSB formation in *Rev3L(-/-)* cells after MNNG treatment (Fig. 4E). This result confirms that Rev3L is involved in the TLS of methylated bases other than *O*⁶MeG.

In conclusion, the data obtained highlight a pivotal role for Polζ in the protection against two important groups of anticancer drugs (i.e., methylating and chloroethylating agents) that are used as first-line therapy of brain tumors and malignant melanomas. Because the heterozygous cells displayed a phenotype of intermediate sensitivity, the levels of Rev3L may determine drug tolerance. In this context, it is important to note that Rev3L is overexpressed in gliomas (Wang et al., 2009) and can presumably be up-regulated upon drug exposure (Wakana et al., 2000). Combined with our finding that Rev3L levels may be rate-limiting in the development of drug tolerance, Rev3L may be a primary determinant in the development of tolerance to alkylating chemotherapy. Therefore, determination of the expression levels of Rev3L in tumors, before and after treatment, is desirable.

Polζ was shown to cooperate with other DNA polymerases, notably Polη and Polκ, that insert nucleotides opposite the lesion. Polζ in turn performs the extension step until DNA replication can proceed in a normal manner (Shachar et al., 2009). The extension mediated by Polζ is required for TLS across many lesions, including AP sites, which explains why Polζ is an exceptionally important player in tolerating critical DNA damage. It has been shown that it is involved in the tolerance of cisplatin-guanine-guanine intrastrand adducts (Shachar et al., 2009) and interstrand cross-links formed by mitomycin C (Wittschieben et al., 2006). As expected, Polζ knockdown sensitized cells to cisplatin whereas stable overexpression caused resistance (Wang et al., 2009). Here, we extend this finding to *O*⁶-alkylating chemotherapeutics and show that for these agents Polζ is a marker of resistance, acting in concert with MGMT, BER, mismatch repair, and DSB repair in protecting against toxic alkylation damage. Similar to strategies of down-regulating or inhibiting MGMT and BER in tumor cells, it is pertinent to consider Polζ as a novel target in cancer therapy.

Acknowledgments

We acknowledge Georg Nagel for technical assistance.

References

- Beranek DT (1990) Distribution of methyl and ethyl adducts following alkylation with monofunctional alkylating agents. *Mutat Res* 231:11–30.
- Bradford MM (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 72:248–254.
- de Murcia JM, Niedergang C, Trucco C, Ricoul M, Dutrillaux B, Mark M, Oliver FJ, Masson M, Dierich A, LeMour M, et al. (1997) Requirement of poly(ADP-ribose) polymerase in recovery from DNA damage in mice and in cells. *Proc Natl Acad Sci U S A* 94:7303–7307.
- Denny BJ, Wheelhouse RT, Stevens MF, Tsang LL, and Slack JA (1994) NMR and molecular modeling investigation of the mechanism of activation of the antitumor drug temozolomide and its interaction with DNA. *Biochemistry* 33:9045–9051.
- Harris LC, von Wronski MA, Venable CC, Remack JS, Howell SR, and Brent TP (1996) Changes in *O*⁶-methylguanine-DNA methyltransferase expression during immortalization of cloned human fibroblasts. *Carcinogenesis* 17:219–224.
- Kaina B, Christmann M, Naumann S, and Roos WP (2007) MGMT: key node in the battle against genotoxicity, carcinogenicity and apoptosis induced by alkylating agents. *DNA Repair* 6:1079–1099.
- Karran P and Bignami M (1992) Self-destruction and tolerance in resistance of mammalian cells to alkylation damage. *Nucleic Acids Res* 20:2933–2940.
- Kawamura K, O-Wang J, Bahar R, Koshikawa N, Shishikura T, Nakagawara A, Sakiyama S, Kajiura K, Kimura M, and Tagawa M (2001) The error-prone DNA

- polymerase zeta catalytic subunit (Rev3) gene is ubiquitously expressed in normal and malignant human tissues. *Int J Oncol* **18**:97–103.
- Lips J and Kaina B (2001) DNA double-strand breaks trigger apoptosis in p53-deficient fibroblasts. *Carcinogenesis* **22**:579–585.
- Liu L, Yan L, Donze JR, and Gerson SL (2003) Blockage of abasic site repair enhances antitumor efficacy of 1,3-bis-(2-chloroethyl)-1-nitrosourea in colon tumor xenografts. *Mol Cancer Ther* **2**:1061–1066.
- Moschel RC, McDougall MG, Dolan ME, Stine L, and Pegg AE (1992) Structural features of substituted purine derivatives compatible with depletion of human O⁶-alkylguanine-DNA alkyltransferase. *J Med Chem* **35**:4486–4491.
- Ochs K, Lips J, Proffittlich S, and Kaina B (2002) Deficiency in DNA polymerase beta provokes replication-dependent apoptosis via DNA breakage, Bcl-2 decline and caspase-3/9 activation. *Cancer Res* **62**:1524–1530.
- Okada T, Sonoda E, Yoshimura M, Kawano Y, Saya H, Kohzaki M, and Takeda S (2005) Multiple roles of vertebrate REV genes in DNA repair and recombination. *Mol Cell Biol* **25**:6103–6111.
- Olive PL, Ban  th JP, and Durand RE (1990) Heterogeneity in radiation-induced DNA damage and repair in tumor and normal cells measured using the “comet” assay. *Radiat Res* **122**:86–94.
- Preuss I, Eberhagen I, Haas S, Eibl RH, Kaufmann M, von Minckwitz G, and Kaina B (1995) O⁶-Methylguanine-DNA methyltransferase activity in breast and brain tumors. *Int J Cancer* **61**:321–326.
- Rattray AJ and Strathern JN (2003) Error-prone DNA polymerases: when making a mistake is the only way to get ahead. *Annu Rev Genet* **37**:31–66.
- Renart J, Reiser J, and Stark GR (1979) Transfer of proteins from gels to diazobenzyloxymethyl-paper and detection with antisera: a method for studying antibody specificity and antigen structure. *Proc Natl Acad Sci U S A* **76**:3116–3120.
- Roos WP and Kaina B (2006) DNA damage-induced cell death by apoptosis. *Trends Mol Med* **12**:440–450.
- Shachar S, Ziv O, Avkin S, Adar S, Wittschieben J, Reissner T, Chaney S, Friedberg EC, Wang Z, Carell T, et al. (2009) Two-polymerase mechanisms dictate error-free and error-prone translesion DNA synthesis in mammals. *EMBO J* **28**:383–393.
- Sobel RW, Horton JK, K  hn R, Gu H, Singhal RK, Prasad R, Rajewsky K, and Wilson SH (1996) Requirement of mammalian DNA polymerase-beta in base-excision repair. *Nature* **379**:183–186.
- Takenaka K, Ogi T, Okada T, Sonoda E, Guo C, Friedberg EC, and Takeda S (2006)

- Involvement of vertebrate Pol   in translesion DNA synthesis across DNA mono-alkylation damage. *J Biol Chem* **281**:2000–2004.
- Tapiero H, Yin MB, Catalin J, Paraire M, Deloffre P, Rustum Y, Bizzari JP, and Tew KD (1989) Cytotoxicity and DNA damaging effects of a new nitrosourea, fotemustine, diethyl-1-(3-(2-chloroethyl)-3-nitrosoureido) ethylphosphonate-S10036. *Anti-cancer Res* **9**:1617–1622.
- Taverna P, Liu L, Hwang HS, Hanson AJ, Kinsella TJ, and Gerson SL (2001) Methoxyamine potentiates DNA single strand breaks and double strand breaks induced by temozolomide in colon cancer cells. *Mutat Res* **485**:269–281.
- Van Sloun PP, Varlet I, Sonneveld E, Boei JJ, Romeijn RJ, Eeken JC, and De Wind N (2002) Involvement of mouse Rev3 in tolerance of endogenous and exogenous DNA damage. *Mol Cell Biol* **22**:2159–2169.
- Wakana S, Sugaya E, Naramoto F, Yokote N, Maruyama C, Jin W, Ohguchi H, Tsuda T, Sugaya A, and Kajiwaru K (2000) Gene mapping of SEZ group genes and determination of pentylene-tetrazol susceptible quantitative trait loci in the mouse chromosome. *Brain Res* **857**:286–290.
- Wang HB, Zhang SY, Wang S, Lv J, Wu WT, Weng L, Chen D, Zhang Y, Lu ZP, Yang JM, et al. (2009) REV3L confers chemoresistance to cisplatin in human gliomas: the potential of its RNAi for synergistic therapy. *Neuro Oncol* doi: 10.1215/15228517-2009-015.
- Wittschieben JP, Reshmi SC, Gollin SM, and Wood RD (2006) Loss of DNA polymerase zeta causes chromosomal instability in mammalian cells. *Cancer Res* **66**:134–142.
- Zander L and Bemark M (2004) Immortalized mouse cell lines that lack a functional Rev3 gene are hypersensitive to UV irradiation and cisplatin treatment. *DNA Repair* **3**:743–752.
- Zdzienicka MZ, van der Schans GP, Natarajan AT, Thompson LH, Neuteboom I, and Simons JW (1992) A Chinese hamster ovary cell mutant (EM-C11) with sensitivity to simple alkylating agents and a very high level of sister chromatid exchanges. *Mutagenesis* **7**:265–269.

Address correspondence to: Dr. Bernd Kaina, Institute of Toxicology, University Medicine Mainz, Obere Zahlbacher Str. 67, D-55131 Mainz, Germany. E-mail: kaina@uni-mainz.de