

Nectandrin B Activates Endothelial Nitric-Oxide Synthase Phosphorylation in Endothelial Cells: Role of the AMP-Activated Protein Kinase/Estrogen Receptor α /Phosphatidylinositol 3-kinase/Akt Pathway

Tran Thi Hien, Won Keun Oh, Phi Hung Nguyen, Seok Jeong Oh, Moo Yeol Lee, and Keon Wook Kang

BK21 Project Team, College of Pharmacy, Chosun University, Gwangju, Republic of Korea (T.T.H., W.K.O., P.H.N.); College of Pharmacy, Dongguk University, Goyang, Republic of Korea (S.J.O., M.Y.L.); and College of Pharmacy and Research Institute of Pharmaceutical Sciences, Seoul National University, Seoul, Republic of Korea (T.T.H., K.W.K.)

Received May 9, 2011; accepted September 22, 2011

ABSTRACT

We revealed previously that nectandrin B isolated from *Myristica fragrans* (nutmeg, Myristicaceae) functions as a potent AMP-activated protein kinase (AMPK) activator and showed its antiobesity effect. In this study, we investigated whether nectandrin B affects phosphorylation of endothelial nitric-oxide synthase (eNOS) in human endothelial cells. Nectandrin B increased the phosphorylation of eNOS and nitric oxide (NO) production in a concentration-dependent manner and maximal effect was found at 10 μ g/ml. Nectandrin B activates AMPK, presumably via Ca^{2+} /calmodulin kinase II activation and nectandrin B-stimulated eNOS phosphorylation was reversed by AMPK inhibition. Both the enzyme activity of phosphatidyl-

inositol 3-kinase (PI3K) and the estrogen receptor (ER)-dependent reporter gene transcription were enhanced by nectandrin B. ER α inhibition by specific antagonist or small interfering siRNA (siRNA) suppressed nectandrin B-mediated eNOS phosphorylation. Moreover, AMPK inhibition significantly reversed the activation of ER-dependent transcription and PI3K activation in response to nectandrin B. Nectandrin B evoked endothelium-dependent relaxation in rat aortic rings, and this was blocked by inhibition of AMPK, ER, or PI3K. These results suggest that potent AMPK activator nectandrin B enhances NO production via eNOS phosphorylation in endothelial cells and ER α -dependent PI3K activity is required.

Introduction

Endothelial function is frequently impaired during atherosclerosis, hyperlipidemia, and diabetes, and this has been well correlated with a high risk of cardiovascular events (Gokce et al., 2003; Widlansky et al., 2003). In particular,

endothelial dysfunction is a key event in both clinical and experimental type II diabetes (Morcos et al., 2001; Schalkwijk and Stehouwer, 2005). Impaired endothelium-dependent vasodilation and increased adhesion of monocytes and platelets are frequently found in diabetes (Fatehi-Hasanabad et al., 2010).

This work was supported by the National Research Foundation of Korea, funded by the Korean government [Grant 2010-0001707]; and the Ministry of Education Science and Technology of the Korean government [Post 21st Frontier Research Program Grant 2010-0026355].

Article, publication date, and citation information can be found at <http://molpharm.aspetjournals.org>.
doi:10.1124/mol.111.073502.

The endothelial nitric-oxide synthase (eNOS) has important basal regulatory functions in the vasculature. Constitutively expressed eNOS in endothelial cells oxidizes L-arginine to generate L-citrulline and nitric oxide (NO) in response to diverse stimuli such as shear stress (Moncada and Higgs, 1993; Li and Förstermann, 2000). Because endothelial NO

ABBREVIATIONS: eNOS, endothelial nitric-oxide synthase; AICAR, 5-aminoimidazole-4-carboxamide-1- β -D-ribofuranoside; AMPK, AMP-activated protein kinase; HUVEC, human umbilical vein endothelial cell; Akt, protein kinase B; ERK, extracellular signal regulated kinase; JNK, c-Jun N-terminal kinase; ER, estrogen receptor; ACC, acetyl CoA carboxylase; CaMK, Ca^{2+} /calmodulin-dependent protein kinase; LY294002, 2-(4-morpholinyl)-8-phenyl-4H-1-benzopyran-4-one; PD98059, 2'-amino-3'-methoxyflavone; B203580, 4-(4-fluorophenyl)-2-(4-methylsulfinylphenyl)-5-(4-pyridyl)-1H-imidazole; SP600125, 1,9-pyrazoloanthrone; ICI 182780, fulvestrant; A3281, N-(6-aminoethyl)-5-chloro-1-naphthalenesulfonamide hydrochloride; PPT, 4,4',4''-(4-Propyl-[1H]-pyrazole-1,3,5-triyl)trisphenol; DPN, diethylpropionitrile; MPP, methyl-piperidino-pyrazole; THC, tetrahydrochrysene; siRNA, small interfering RNA; DN, dominant-negative; CA, constitutive active; MTT, 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl-tetrazolium bromide; DAF-2 DA, 4,5-diaminofluorescein diacetate; TE, Tris-EDTA; PI3K, phosphatidylinositol 3-kinase; LKB, liver kinase B1; compound C, 6-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-3-pyridin-4-yl-pyrazolo[1,5-a]-pyrimidine.




Fig. 1. A, structure of nectandrin B. B, cytotoxicity of nectandrin B in ECV 304 cells. Cells were seeded in a 48-well plate, and various concentrations of nectandrin B were incubated for 24 h. Cell viability were estimated by MTT assay. Each bar represents the mean \pm S.D. calculated from eight different samples (*, $p < 0.05$, significant compared with control).

production evokes a decrease in vascular tone and inhibits oxidation of low-density lipoprotein (Howes et al., 1997), safe compounds that activate eNOS may be beneficial for patients with chronic cardiovascular diseases.

A series of recent studies have revealed that activation of AMP-activated protein kinase (AMPK) has beneficial effects on endothelial dysfunction. AMPK activation inhibits oxidized low-density lipoprotein-triggered endoplasmic reticulum stress in endothelial cells (Dong et al., 2010) and metformin, a clinical AMPK activator, normalizes endothelial function by suppressing vasoconstrictor prostanooids in arteries from a type II diabetes rat model (Matsumoto et al., 2008). Moreover, AMPK activity is involved in eNOS activation. An experimental AMPK activator, 5-aminoimidazole-4-carboxamide-1- β -D-ribofuranoside (AICAR), increased eNOS activity via Ser1177 phosphorylation; infusion of AICAR markedly increased muscle microvascular blood volume (Bradley et al., 2010). Conversely, silencing AMPK α 1 in human umbilical vein endothelial cells (HUVECs) reduced eNOS content (Colombo and Moncada, 2009).

Plant-derived compounds are becoming of increasing interest as potential antiatherosclerosis therapeutics. It has been reported that some natural compounds in fruits, vegetables, oil seeds, and herbs have lipid-lowering effects and reduce atherosclerotic lesions (Afrose et al., 2009; Katsuda et al., 2009; Magnone et al., 2009). *M. fragrans* (nutmeg) has been used as a food and cosmetic as well as a traditional oriental medicine against dysentery, diarrhea, and pain (Grover et




Fig. 2a. Effect of nectandrin B on eNOS phosphorylation and expression. A, concentration-dependent eNOS phosphorylation by nectandrin B. ECV 304 cells were treated with nectandrin B (1–10 µg/ml) for 1 h, and the total cell lysates were subjected to immunoblotting with antibody against Ser1177 phosphorylated eNOS or total eNOS. Relative changes in the eNOS phosphorylation were assessed by scanning densitometry. Data represent the means \pm S.D. of 3 separate experiments (*, $p < 0.05$, significant compared with control; control level = 1). B, time course of eNOS phosphorylation by nectandrin B. ECV 304 cells were treated with nectandrin B (10 µg/ml) for the indicated time (5–60 min). Phosphorylated eNOS and total eNOS levels were detected by Western blot analyses under identical conditions. Relative changes in the eNOS phosphorylation were assessed by scanning densitometry. Data represent the means \pm S.D. of three separate experiments (*, $p < 0.05$, significant compared with control; control level = 1). C, no change of eNOS expression by nectandrin B. ECV 304 cells were treated with nectandrin B (10 µg/ml) for 3 to 24 h, and the total cell lysates were subjected to immunoblotting with eNOS and β -actin antibodies. Relative change in the eNOS protein expression was assessed by scanning densitometry. Data represent the means \pm S.D. of three separate experiments (control level = 1).




Fig. 2b. Concentration-(D) and time-dependent (E) NO production by nectandrin B. ECV 304 cells were serum-starved overnight and loaded with DAF2-DA as described under *Materials and Methods*. Cells were then stimulated with nectandrin B (0.5–10 µg/ml) for 60 min or nectandrin B (10 µg/ml) for the indicated time. After nectandrin B treatments, cells were fixed in 5% paraformaldehyde and visualized with an epifluorescent microscope. Emission of green fluorescence is indicative of NO production. FITC, fluorescein isothiocyanate.

al., 2002). Furthermore, its seed extracts possess antihyperlipidemic and antiatherosclerotic activities in vivo (Sharma et al., 1995). However, the mechanisms underlying these activities have not been clarified. To address this gap in our knowledge, we isolated seven 2,5-bis-aryl-3,4-dimethyltetrahydrofuran lignans from total extracts of *M. fragrans*, and nectandrin B had a strong AMPK stimulation effect at a concentration of 5 µM in differentiated C2C12 cells (Nguyen et al., 2010).

In the present study, we found that nectandrin B potently activates AMPK in both ECV 304 (a human endothelial cell line) and HUVECs in primary cultures. We determined whether nectandrin B affects phosphorylation and expression of eNOS and tried to identify cellular signaling pathways for the phosphorylation and expression of eNOS in response to nectandrin B.

Materials and Methods

Nectandrin B Isolation. The dried semens of *M. fragrans* (nutmeg) were purchased at a folk medicine market in Gwangju city, Republic of Korea. The sample was identified by Professor Y. H. Moon at Chosun University, and its specimen (0010) was deposited at the Department of Pharmacy, Chosun University. The EtOH extract of *M. fragrans* was subjected to an HP-20 column (10 × 60 cm), eluted with a gradient of EtOH in H₂O (60, 80, 90, and 100%, each 3 L), and finally washed by acetone (2 L) to give five fractions. Bioassay of the five fractions on the AMPK activity revealed that the 80% ethanol-eluted fraction was most active as AMPK activator. This fraction was further

chromatographed over silica gel (6 × 60 cm; 63–200 µm particle size) using a gradient of *n*-hexane/acetone (from 6:1 to 0:1), to yield five fractions (F.1–F.5) according to their profiles. Nectandrin B was purified from a part of fraction 2 by chromatography on a reversed-phase ODS-A column (20 × 150 mm, 4 µm particle size; YMC Co., Ltd., Kyoto, Japan) or HP-20 column (10 × 60 cm; Mitsubishi Chemical Co., Tokyo, Japan) eluted with MeOH/H₂O (1.5:1, to 2:1, each 3 L) and purity was confirmed by high performance liquid chromatography.

Materials. Antibodies against eNOS, phospho-eNOS, phospho-Akt, Akt, phospho-mitogen-activated protein kinase, p38, phospho-extracellular signal regulated kinase (ERK), ERK, phospho-c-Jun N-terminal kinase (JNK), JNK, ERα, phospho-AMPK, AMPK, phospho-acetyl CoA carboxylase (ACC), ACC, phospho-Ca²⁺/calmodulin-dependent protein kinase II (CaMK II), and horseradish peroxidase-conjugated anti-mouse and anti-rabbit IgG antibodies were purchased from Cell Signaling Technology (Danvers, MA). 2-(4-Morpholinyl)-8-phenyl-4*H*-1-benzopyran-4-one (LY294002), 2'-amino-3'-methoxyflavone (PD98059), 4-(4-fluorophenyl)-2-(4-methylsulfinylphenyl)-5-(4-pyridyl)-1*H*-imidazole (SB203580), 1,9-pyrazoloanthrone (SP600125), fulvestrant (ICI-182780), *N*-(6-aminohexyl)-5-chloro-1-naphthalenesulfonamide hydrochloride (A3281), compound C (6-[4-(2-piperidin-1-yl-ethoxy)-phenyl]-3-pyridin-4-yl-pyrazolo[1,5-*a*]-pyrimidine), and *N*^G-nitro-L-arginine methyl ester (NOS inhibitor) were purchased from Calbiochem (La Jolla, CA). 4,4',4''-(4-Propyl-[1*H*]-pyrazole-1,3,5-triyl)trisphenol (PPT), diarylpropionitrile (DPN), methyl-piperidino-pyrazole (MPP), and tetrahydrochrysenes (THC) were obtained from Tocris Biosciences (Ellisville, MO). ERβ antibody was obtained from Abcam (Cambridge, MA). siRNAs targeting ERα and ERβ were purchased from Santa Cruz Biotechnology

(Santa Cruz, CA). Actin antibody and other reagents used for molecular studies were obtained from Sigma-Aldrich (St. Louis, MO). Dominant-negative mutant (DN-AMPK) or constitutive active forms (CA-AMPK) of AMPK overexpression plasmids were kindly donated by Dr. J. H. Ha of Kyunghee University (Seoul, Korea).

Cell Culture. ECV 304 cells were obtained from the American Type Culture Collection (Manassas, VA). The cells were maintained at 37°C in an incubator with a humidified atmosphere of 5% CO₂ and cultured in Dulbecco's modified Eagle's medium containing 10% fetal bovine serum. Primary cultured HUVEC cells were purchased from Innopharmascreeen (Asan, Chungnam, Korea) and cultured in M199 medium containing 10 units/ml heparin, 20% fetal bovine serum, and 20 ng/ml fibroblast growth factor. Nectandrin B was dissolved in dimethyl sulfoxide, and the stock solutions were added directly to the culture media. Control cells were treated with dimethyl sulfoxide only. The final concentration of solvent was always <0.1%.

Cytotoxicity of Nectandrin B in ECV 304 Cells. Viable adherent cells were stained with 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl-tetrazolium bromide (MTT) (2 mg/ml) for 4 h. Media were then removed, and the formazan crystals produced were dissolved by adding 200 μ l of dimethyl sulfoxide. Absorbance was assayed at 570

nm using a microplate reader (TriStar LB 941; Berthold Technologies, Bad Wildbad, Germany), and cell viabilities were expressed as ratios versus untreated control cells.

Reporter Gene Analysis. A dual-luciferase reporter assay system (Promega, Madison, WI) was used to determine gene promoter activity. In brief, cells were plated in 12-well plates and transiently transfected with the reporter and phRL-SV plasmids (*Renilla reniformis* luciferase expression for normalization; Promega, Madison, WI) using Hillymax reagent (Dojindo Molecular Technologies, Kumamoto, Japan). The cells were then incubated in culture medium without serum for 18 h. Firefly and *R. reniformis* luciferase activities in the cell lysates were measured using a luminometer (LB 941; Berthold Technologies, Bad Wildbad, Germany). The relative luciferase activity was calculated by normalizing the promoter-driven firefly luciferase activity to the *R. reniformis* luciferase activity.

Western Blot Analysis. After treatment, cells were collected and washed with ice-cold phosphate-buffered saline. The harvested cells were then lysed on ice for 30 min in 100 μ l of lysis buffer (120 mM NaCl, 40 mM Tris, pH 8, 0.1% Nonidet P-40) and centrifuged at 12,000 rpm for 30 min. Supernatants were collected from the lysates, and protein concentrations were determined using the bicinchoninic acid protein assay kit (Thermo Fisher Scientific, Waltham, MA).




Fig. 3. Effect of nectandrin B on eNOS phosphorylation and NO production in HUVEC cells. A, nitrite amounts in culture medium. ECV 304 cells were serum-starved overnight, incubated with nectandrin B (1–10 μ g/ml) for additional 24 h and culture media were collected for nitrite determination. Data represent the means \pm S.D. of three separate experiments (*, $p < 0.05$, significant compared with control; control level = 1). B, concentration-dependent eNOS phosphorylation in HUVEC cells. HUVEC cells were treated with nectandrin B (1–10 μ g/ml) for 1 h, and the total cell lysates were subjected to immunoblotting with antibodies against Ser1177 phosphorylated eNOS, total eNOS, and β -actin. C, NO production by nectandrin B in HUVEC cells. HUVEC cells were serum-starved overnight and loaded with DAF2-DA as described under *Materials and Methods*.

Aliquots of the lysates (30 μ g of protein) were boiled for 5 min and electrophoresed on 10% SDS-polyacrylamide gels. Proteins in the gels were transferred onto nitrocellulose membranes, which were then incubated with primary antibodies or mouse monoclonal β -actin antibodies. The membranes were further incubated with secondary anti-mouse or anti-rabbit antibodies. Finally, protein bands were detected using an enhanced chemiluminescence Western blotting detection kit (Thermo Fisher Scientific).

Measurement of NO Production. Production of NO was assessed using the NO-specific fluorescent dye 4,5-diaminofluorescein diacetate (DAF-2 DA; Cayman Chemical, Ann Arbor, MI) as described previously (Formoso et al.,2006). In brief, ECV 304 cells were grown to 95% confluence in Lab-Tek chamber slides (Nalge Nunc International, Rochester, NY) and serum-starved overnight. Cells were then loaded with DAF-2 DA (final concentration, 2 μ M) for 30 min at 37°C, rinsed three times with Dulbecco’s modified Eagle’s medium, and kept in the dark. Cells were then treated without or with nectandrin B as indicated in the figure legends. The cells were fixed in 5% paraformaldehyde for 5 min at 4°C. Fixed cells were visualized using a fluorescence microscope (Axiovert 200M; Carl Zeiss, Jena, Germany) and an inverted epifluorescence microscope with an attached charge-coupled device camera using appropriate filters, with a peak excitation wavelength of 480 nm and a peak emission wavelength of 510 nm.

The amount of NO in the culture media was also determined using the Griess reagent (Enzo Life Sciences, Plymouth Meeting, PA). Culture media obtained 24 h after treatment with various concentrations of nectandrin B were incubated with nitrate reductase for 1 h at 37°C to reduce nitrate to nitrite. An equal volume of Griess reagent was then added, and the optical density of the samples were measured at a wavelength of 540 nm. Data were expressed as fold change over untreated controls.

Recombinant ER α Binding Assay. Vehicle or test chemicals were incubated with 1 nM tritiated estradiol ($[^3\text{H}]\text{E}_2$; PerkinElmer Life and Analytical Sciences, Waltham, MA) and 0.6 nM recombinant human ER α (MyBioSource, San Diego, CA) in TE buffer (10 mM Tris and 1 mM EDTA, pH 7.5) at 4°C overnight. Hydroxylapatite (60% in TE buffer) was added, mixed well, and incubated for 15 min at room temperature. The resulting slurry was washed three times by centrifugation with TE buffer. Bound ligand was extracted by incubation of the slurry with absolute ethanol at 30°C for 10 min. Tritium (^3H) decay (counts per minute) was measured by liquid scintillation in a β -counter (PerkinElmer Life and Analytical Sciences).

Organ Chamber Study. Male Sprague-Dawley rats (270–330 g) were sacrificed, and thoracic aortas were carefully removed and placed in a modified Krebs-Ringer bicarbonate solution containing 118.3 mM NaCl, 4.7 mM KCl, 1.2 mM MgSO_4 , 1.2 mM KH_2PO_4 , 2.5 mM CaCl_2 , 25.0 mM NaHCO_3 , 0.016 mM Ca^{2+} EDTA, and 11.1 mM glucose (control solution). The aortas were cleaned of loose connective tissue and then cut into rings (2 mm wide). The aortic rings were suspended horizontally between two stainless steel stirrups in the organ chambers filled with 5 ml of control solution (37°C, pH 7.4) and bubbled with 95% O_2 and 5% CO_2 . The change in tension was measured isometrically with Grass FT03 force transducers (Grass Instrument Co., Quincy, MA), and data were acquired and analyzed with a PowerLab 8/30 Data Acquisition System and LabChart pro software (AD Instruments, Colorado Springs, CO). The rings were stretched progressively to the optimal tension (2 g) before the addition of 90 mM KCl. Once the plateau of the contraction elicited by KCl was obtained, the aortic rings were rinsed three times with warm control solution, and 1 μ M acetylcholine-mediated relaxation was tested in the precontracted rings by 1 μ M phenylephrine to check endothelium-dependent relaxation responsiveness. In some experiments, rings were incubated for 30 min with LY294002 (10




Fig. 4. AMPK activation by nectandrin B. A, concentration-dependent phosphorylation of AMPK or ACC by nectandrin B in ECV 304 cells. The cells were incubated with nectandrin B (1–10 μ g/ml) for 1 h. AICAR (1 mM) was used as a representative AMPK activator. B, time-course phosphorylation of AMPK, ACC, or CaMK II by nectandrin B in ECV 304 cells. Nectandrin B (10 μ g/ml) was treated for the indicated time. C, concentration-dependent phosphorylation of AMPK or ACC by nectandrin B in HUVEC cells. The cells were incubated with nectandrin B (1–10 μ g/ml) for 1 h.

μM), ICI-182780 (100 nM), or compound C (20 μM). After 30-min incubation, 10 $\mu\text{g/ml}$ nectandrin B was added and the relaxation response was monitored in the precontracted aortic rings after addition of 1 μM phenylephrine.

Statistical Analysis. Unpaired Student's *t* test was used to determine the significance of differences between treatment groups. Statistical significance was accepted for *p* values < 0.05.

Results

Cytotoxicity of Nectandrin B in ECV 304 Cells. Initially, we determined the cytotoxicity of nectandrin B to ECV 304 cells by MTT assay. Figure 1B shows that nectandrin B at the tested concentrations did not cause cytotoxicity except at concentrations above 30 $\mu\text{g/ml}$. Thus, we treated cells with nectandrin B in the concentration range 1–10 $\mu\text{g/ml}$ during subsequent experiments.

Nectandrin B Increases Phosphorylation of eNOS and Production of NO in Endothelial Cells. NO produc-

tion from eNOS activation plays a protective physiological role in the vasculature (Li and Förstermann, 2000). We did Western blot analyses to detect changes in eNOS phosphorylation. When cells were treated with 1 to 10 $\mu\text{g/ml}$ nectandrin B for 1 h and 10 $\mu\text{g/ml}$ nectandrin B at the indicated time points (5–60 min), the levels of phosphorylated eNOS were increased in a concentration- and time-dependent manner (Fig. 2, A and B). However, nectandrin B incubation for 3 to 24 h did not affect the expression level of eNOS in ECV 304 cells (Fig. 2C). To examine whether eNOS phosphorylation by nectandrin B stimulates NO production, ECV 304 cells were loaded with DAF-2 DA, a dye that upon binding to an oxidized species of NO results in fluorescence. As shown in Fig. 2, D and E, green fluorescence (indicative of NO production) was increased by nectandrin B in a concentration- and time-dependent manner. We further determined nitrite levels in culture medium. Nitrite production was significantly enhanced by 3 or 10 $\mu\text{g/ml}$ nectandrin B (Fig. 3A). To confirm





Fig. 5a. Involvement of AMPK in nectandrin B-induced eNOS phosphorylation. A, NO production by AICAR. 1 mM AICAR was treated for 1 h in ECV 304 cells. B, effect of compound C on nectandrin B-induced eNOS phosphorylation. ECV cells were pretreated with 10 μM compound C (AMPK inhibitor) for 30 min and then incubated with 10 $\mu\text{g/ml}$ nectandrin B for an additional 60 min. Phosphorylated eNOS and ACC were detected by Western blot analyses. Relative changes in the eNOS phosphorylation were assessed by scanning densitometry. Data represent the means \pm S.D. of three separate experiments (*, *p* < 0.05, significant compared with control; control level = 1; #, *p* < 0.05, significant compared with nectandrin B-treated group). C, effect of DN-AMPK transfection on nectandrin B-induced eNOS phosphorylation. ECV 304 cells were transfected with DN-AMPK and pcDNA control for 24 h. The cells were then treated with 10 $\mu\text{g/ml}$ nectandrin B for 60 min. Phosphorylated eNOS and ACC were detected by Western blot analyses. Relative changes in the eNOS phosphorylation were assessed by scanning densitometry. Data represent the means \pm S.D. of three separate experiments (*, *p* < 0.05, significant compared with control; control level = 1; #, *p* < 0.05, significant compared with nectandrin B-treated group).

D



E




Fig. 5b. D, effect of CA-AMPK on the phosphorylation of eNOS and ACC. ECV 304 cells were transfected with CA-AMPK and pcDNA control for 24 h, the cells were then treated with or without 10 μg/ml nectandrin B for 60 min. Phosphorylated eNOS and ACC were detected by Western blot analyses. Relative changes in the eNOS phosphorylation were assessed by scanning densitometry. Data represent the means ± S.D. of three separate experiments (*, $p < 0.05$, significant compared with control; control level = 1). **E,** role of CaMK II in nectandrin B-mediated AMPK activation and eNOS phosphorylation. ECV 304 cells were pretreated with 10 μM A3281 (calmodulin inhibitor) for 30 min and then incubated with 10 μg/ml nectandrin B for additional 60 min. Phosphorylation intensity of eNOS, AMPK, or ACC was detected by Western blot analyses. Relative changes in the eNOS phosphorylation were assessed by scanning densitometry. Data represent the means ± S.D. of three separate experiments (*, $p < 0.05$, significant compared with control; control level = 1; #, $p < 0.05$, significant compared with nectandrin B-treated group). FITC, fluorescein isothiocyanate.

these results, HUVECs were treated with 1 to 10 μg/ml nectandrin B, and eNOS phosphorylation and NO production were detected. As expected, phosphorylated eNOS level and DAF-2 DA fluorescence were enhanced by nectandrin B treatment in HUVECs (Fig. 3, B and C).

Nectandrin B Increases eNOS Phosphorylation via AMPK through CaMKII. It is known that phosphorylation of Ser1177 in eNOS, which plays an important role in the regulation of eNOS activity, is induced by AMPK and results in NO production in endothelial cells (Chen et al., 1999; Morrow et al., 2003). In our previous study, we reported that nectandrin B potently activates AMPK in differentiated skeletal muscle cells (Nguyen et al., 2010). Hence, we examined the effect of nectandrin B on AMPK activity in human endothelial cells. Representative AMPK activation markers, phosphorylation of AMPK and ACC, were increased by nectandrin B in concentration- and time-dependent manners in ECV 304 cells (Fig. 4, A and B). The same results were also found in HUVECs (Fig. 4C).

LKB1 (also called STK11) is a mammalian kinase that activates AMPK by phosphorylating Thr172 on its catalytic (α) subunit (Carling et al., 2008). Recent studies have reported evidence that CaMKs can act upstream of AMPK, at least in some cell types (Hurley et al., 2005; Woods et al., 2005). ECV 304 cells were treated with nectandrin B for the indicated times and we determined the active phosphorylated form of CaMK II and LKB1. Figure 4B shows that the

levels of phospho-CaMK II were increased by 10 μg/ml nectandrin B, but this treatment did not affect LKB1 phosphorylation (data not shown), which suggests that nectandrin B-induced AMPK activation may be mediated through CaMK II activation.

We then tested whether AMPK is necessary for eNOS phosphorylation induced by nectandrin B. AMPK activation by 1 mM AICAR treatment increased NO production in ECV 304 cells (Fig. 5A). ECV 304 cells were pretreated with compound C, an AMPK inhibitor, and we examined nectandrin B-dependent eNOS phosphorylation. Nectandrin B-induced phosphorylation of ACC was impaired in compound C-pretreated ECV 304 cells, and eNOS phosphorylation was attenuated (Fig. 5B), which suggests that AMPK is required for eNOS activation. To confirm these results, we used overexpression vector for DN-AMPK or CA-AMPK of AMPK. Transfection of DN-AMPK before nectandrin B treatment reduced nectandrin B-stimulated ACC and eNOS phosphorylation (Fig. 5C). Conversely, CA-AMPK transfection markedly increased phosphorylation levels of ACC and eNOS (Fig. 5D). To investigate whether the CaMK II pathway is involved in the process by which nectandrin B causes AMPK and eNOS phosphorylation, ECV 304 cells were pretreated with A3281, a CaMK II inhibitor (10 μM), and exposed to nectandrin B. As shown in Fig. 5E, A3281 blocked nectandrin B-induced phosphorylation of eNOS, ACC, and AMPK. These results

suggest that nectandrin B-mediated eNOS phosphorylation is dependent on AMPK signaling through CaMK II.

Role of the PI3K/Akt Pathway in eNOS Phosphorylation by Nectandrin B. Diverse kinases such as PI3K/Akt, p38 kinase, ERK, and JNK have been shown to be involved in cellular signaling for vascular relaxation and NO production (Merla et al., 2007; Grossini et al., 2008). To further elucidate the upstream signaling pathways involved in nectandrin B-mediated eNOS phosphorylation and subsequent NO production, we examined the activity of PI3K, ERK, p38 kinase, and JNK in nectandrin B-treated ECV 304 cells. Western blot analyses using phospho-specific antibodies showed that nectandrin B incubation caused a sustained phosphorylation of Akt or JNK but not ERK and p38 (Fig. 6A). To address the role of PI3K or JNK activation in eNOS phosphorylation by nectandrin B, the effects of specific kinase inhibitors were investigated. LY294002, a PI3K inhibitor, significantly reduced nectandrin B-induced eNOS phosphorylation (Fig. 6B). However, SP600125, an inhibitor of the JNK pathway, had no effect (Fig. 6C).

Involvement of the ER α -Dependent PI3K/Akt Pathway in Phosphorylation of eNOS by Nectandrin B. A recent study suggests that AMPK activation restores estrogen responsiveness in human endothelial cells (Chakrabarti and Davide, 2009). It has also been shown that nordihy-

droguaiaretic acid, a lignan, has estrogenic activity (Fujimoto et al., 2004). Until now, no evidence has been reported of a correlation between ER and eNOS phosphorylation induced by nectandrin B. When we determined ER-dependent transcription by using an ERE reporter, nectandrin B significantly increased the reporter activities of ERE in a concentration-dependent manner, but the increase was marginal compared with the effects of a full ER agonist, 17- β -estradiol (Fig. 7A). It has been shown that ER α directly interacts with PI3K and modulates its activity in human vascular endothelial cells (Simoncini et al., 2000). To further test the possible role of ER activation in PI3K-dependent eNOS phosphorylation in nectandrin B-exposed endothelial cells, we examined whether ICI-182780, an ER-specific inhibitor, affects the eNOS phosphorylation that occurs in response to nectandrin B. ICI-182780 potentially suppressed eNOS phosphorylation as well as Akt phosphorylation (Fig. 7B). These results indicate that nectandrin B-stimulated eNOS phosphorylation is linked with ER-dependent PI3K/Akt pathways. We then confirmed the effects of an ER antagonist and a PI3K inhibitor on NO production induced by nectandrin B. As shown in Fig. 7C, nectandrin B-mediated NO production (DAF-2 fluorescence) was suppressed by ICI-182780 or LY294002 pretreatment. These data suggest that ER activation is critical for PI3K/Akt-mediated eNOS phosphorylation by nectandrin B.




Fig. 6. Role of PI3K/Akt pathway in nectandrin B-stimulated eNOS phosphorylation. **A**, effect of nectandrin B on the activities of PI3K, ERK, JNK, and p38 kinase. ECV 304 Cells were treated with 10 µg/ml nectandrin B for the indicated times and then immunoblotted with phosphorylation-specific antibodies that recognize phospho-Akt (p-Akt), phospho-ERK (p-ERK), phospho-p38 kinase (p-p38), and phospho-JNK (p-JNK). Parallel immunoblots were analyzed for total kinase levels with anti-Akt, ERK, p38, and JNK antibodies. **B**, effect of PI3K inhibitor on nectandrin B-stimulated eNOS phosphorylation. ECV 304 cells were preincubated with 10 µM LY294002 (LY) and then cells were incubated with 10 µg/ml nectandrin B for 60 min. Cell lysates were subjected to Western blotting with antibodies against phosphorylated eNOS, eNOS, phosphorylated Akt, or Akt. **C**, effects of JNK inhibitor on nectandrin B-stimulated eNOS phosphorylation. ECV 304 cells were preincubated with 10 µM SP600125 (SP), and then cells were incubated with 10 µg/ml nectandrin B for 60 min.




Fig. 7. Role of estrogen receptor signaling in Akt-dependent eNOS phosphorylation. **A**, nectandrin B-induced estrogen response element (ERE) reporter activation. ECV 304 cells were transiently transfected with ERE-Luc plasmid. After transfection, cells were treated with nectandrin B (1–10 μ g/ml) or 17- β -estradiol (E2; 100 nM) for 24 h before lysis and measurement of ERE reporter activity. Data represent the means \pm S.D. of four separate samples (*, $p < 0.05$, significant compared with control). **B**, effect of ER antagonist on nectandrin B-induced phosphorylation of Akt and eNOS. ECV 304 cells were preincubated with 100 nM ICI-182,780 (ER antagonist). The cells were then incubated with 10 μ g/ml nectandrin B for 60 min. Cell lysates were subjected to Western blotting using antibodies against phosphorylated eNOS, eNOS, phosphorylated Akt or Akt. **C**, effects of PI3K inhibitor or ER antagonist on nectandrin B-mediated NO production. Under identical conditions, the cells were serum-starved overnight and loaded with DAF2-DA as described under *Materials and Methods*. Cells were preincubated with inhibitors for 30 min and then incubated with 10 μ g/ml nectandrin B. L- N^G -nitro-L-arginine methyl ester was used as a NOS inhibitor. FITC, fluorescein isothiocyanate.

It has been shown that both ER α and ER β are involved in eNOS phosphorylation in endothelial cells (Mineo and Shaul, 2006). Immunoblot analyses showed that both the receptor types were expressed in ECV 304 cells (Fig. 8A, left). Either DPN (ER α -selective agonist) or PPT (ER β -selective agonist) increased eNOS phosphorylation in ECV 304 cells, demonstrating that both ER subtypes are coupled with eNOS phosphorylation process in this cell type (Fig. 8A, right). We further determined the effects of specific antagonists targeting ER α and ER β on eNOS phosphorylation by nectandrin B. MPP, a selective ER α antagonist, potently suppressed nectandrin B-mediated eNOS phosphorylation, whereas THC, a selective ER β antagonist, did not (Fig. 8B). Moreover, ER α siRNA potently suppressed nectandrin B-mediated eNOS phosphorylation, but ER β siRNA marginally affected eNOS phosphorylation in response to nectandrin B (Fig. 8C). To assess whether nectandrin B directly binds to ER α , we

further performed ER α ligand binding assay using human recombinant ER α . Nectandrin B (3 and 10 μ g/ml) significantly inhibited tritiated estradiol binding to human ER α , but the inhibition intensity is lower than 10 nM 17- β -estradiol (Fig. 8D). These data suggest that nectandrin B acts as a relatively selective agonist on ER α .

ER-dependent eNOS phosphorylation is coupled with diverse signaling molecules. It has been shown that Src kinase mediates PI3K/Akt-dependent rapid eNOS activation in endothelial cells (Haynes et al., 2003). We found that the Src-specific inhibitor 4-amino-5-(4-chlorophenyl)-7-(*t*-butyl)pyrazolo[3,4-*d*]pyrimidine suppressed eNOS phosphorylation by nectandrin B (Fig. 8E). Coimmunoprecipitation studies of plasma membranes from COS-7 cells transfected with ER α and G α proteins demonstrated estrogen-stimulated selective interaction between ER α and G α . Moreover, G α inhibitor pertussis toxin blocked estrogen-dependent eNOS activation




Fig. 8. Crucial role of ER α /Src in nectandrin B-mediated eNOS phosphorylation. **A**, eNOS activation effects of ER α and ER β agonists in ECV 304 cells. Left, ER α and ER β expression was detected by immunoblotting using specific antibodies. Right, DPN (ER α agonist, 100 nM) and PPT (ER β agonist, 100 nM) were exposed to ECV 304 cells for 60 min, and eNOS phosphorylation was determined. **B**, effects of ER α (MPP) and ER β (THC) antagonists on eNOS phosphorylation induced by nectandrin B. ECV 304 cells were preincubated with 10 μ M MPP or 10 μ M THC and then cells were incubated with 10 μ g/ml nectandrin B for 60 min. **C**, effects of siRNAs for ER α (left) and ER β (right) on eNOS phosphorylation induced by nectandrin B. ECV 304 cells were preincubated with 60 pmol of control siRNA, ER α siRNA, or ER β siRNA, and then cells were incubated with 10 μ g/ml nectandrin B for 60 min. **D**, ER α binding activity of nectandrin B. 17- β -estradiol (E2; 10 nM) or 1–10 μ g/ml nectandrin B were incubated with 1 nM tritiated estradiol and 0.6 nM recombinant human ER α protein. Data represent the means \pm S.D. of three separate samples (*, $p < 0.05$; **, $p < 0.01$, significant compared with control). **E**, effects of G α_i inhibitor [pertussis toxin (PTX), 10 nM] and Src inhibitor 4-amino-5-(4-chlorophenyl)-7-(*t*-butyl)pyrazolo[3,4-*d*]pyrimidine (PP2; 10 μ M) on eNOS phosphorylation induced by nectandrin B. ECV 304 cells were preincubated with 10 nM PTX or 10 μ M PP2, and then cells were incubated with 10 μ g/ml nectandrin B for 60 min.

(Mineo and Shaul, 2006). However, pertussis toxin did not abrogate nectandrin B-stimulated eNOS phosphorylation in ECV 304 cells (Fig. 8E). Hence, Src tyrosine kinase, but not G proteins, may be involved in the ER α and subsequent eNOS activation process by nectandrin B.

PI3K/Akt and AMPK pathways are both required for eNOS phosphorylation by nectandrin B. Hence, we studied possible cross-talk between the two kinase pathways. PI3K inhibition by LY294002 pretreatment failed to inhibit the phosphorylation of ACC in response to nectandrin B (Fig. 9A). However, AMPK inhibition by compound C impaired nectandrin B-induced Akt phosphorylation (Fig. 9B). In addition, compound C attenuated ER-dependent reporter activity induced by nectandrin B (Fig. 9C). These data support the conclusion that AMPK/ER α /PI3K/Akt regulates eNOS phosphorylation in response to nectandrin B.

Nectandrin B-Mediated Endothelium-Dependent Relaxation Is Reversed by Inhibition of ER, PI3K, or AMPK.

We did organ chamber studies using endothelium-intact aortic rings. Nectandrin B (10 μ g/ml) relaxed phenylephrine-precontracted aortic rings, and LY294002 (PI3K inhibitor) and ICI-182780 (ER antagonist) completely blocked nectandrin B-mediated vasorelaxation (Fig. 10A). Although compound C (AMPK inhibitor) significantly reversed nectandrin B's relaxation effect, the inhibition intensity was weaker than that of LY294002 or ICI-182780 (Fig. 10A). Immunoblot analysis using the homogenates of aortic rings showed that nectandrin B increased the level of phosphorylated eNOS at 15 to 30 min after 10 μ g/ml nectandrin B exposure to aortic rings (Fig. 10B), which suggests that eNOS phosphorylation and subsequent NO production by nectandrin B is directly coupled with vascular relaxation.




Fig. 9. AMPK is required for the activation of ER/Akt pathway by nectandrin B. **A**, effect of PI3K inhibitor on nectandrin B-stimulated AMPK activation. ECV cells were pretreated with 10 µM LY294002 (LY; PI3K inhibitor) for 30 min and then incubated with 10 µg/ml nectandrin B for an additional 60 min. Phosphorylated AMPK was detected by Western blot analysis. **B**, effect of AMPK inhibitor on nectandrin B-stimulated Akt activation. ECV cells were pretreated with 10 µM compound C (AMPK inhibitor) for 30 min and then incubated with 10 µg/ml nectandrin B for an additional 60 min. Phosphorylated Akt was detected by Western blot analysis. **C**, effect of compound C on nectandrin B-induced increase in ERE activity. ECV 304 cells were transiently transfected with ERE-Luc plasmid. After transfection, cells were incubated with compound C (10 µM) and 10 µg/ml nectandrin B for 24 h before lysis and measurement of ERE reporter activity. Data represent the means \pm S.D. of four separate samples (*, $p < 0.05$, significant compared with control; #, $p < 0.05$, significant compared with nectandrin B-treated group).

Discussion

The species *M. fragrans* has been used traditionally for spices and various medicinal purposes—as a stomachic, carminative, tonic, aphrodisiac, and nervous system stimulant (Nguyen et al., 2010). Several studies have shown that lignan compounds from *M. fragrans* have beneficial anti-inflammatory, anti-diabetes and antioxidant effects (Han et al., 2008a,b; Kwon et al., 2008; Ma et al., 2009; Anggakusuma et al., 2010). AMPK is activated either by an increase in the AMP/ATP ratio during metabolic stress or by activation of upstream kinases such as LKB1 and CaMK (Hardie, 2007). Because AMPK activation suppresses ATP-consuming anabolic pathways and conversely activates ATP-generating catabolic pathways, a selective AMPK activator could function as an antidiabetes and antiobesity agent (Misra, 2008). In this sense, nectandrin B is an attractive natural compound, because the lignan is a potent AMPK activator at relatively low concentrations compared with other phytochemicals.

Here, we demonstrated that nectandrin B stimulates eNOS phosphorylation in human endothelial cells. Although it has been reported that eNOS levels are reduced in cells being silenced by AMPK (Colombo and Moncada, 2009), protein expression of eNOS was not affected by nectandrin B. The PI3K/Akt pathway was reported to be essential for Ser1177 or Ser1179 eNOS phosphorylation (Dimmeler et al., 1999; Fulton et al., 1999) and therefore functions as an essential kinase in regulating eNOS activity and NO production in various circumstances (Thomas et al., 2002; Cai et al., 2003). The current experiments show that nectandrin B strongly activates PI3K, and that eNOS phosphorylation and NO production in response to nectandrin B are attenuated by PI3K inhibition. A recent study showed that ER α directly interacts with PI3K and modulates its activity in human vascular endothelial cells (Simoncini et al., 2000). Furthermore, PI3K is included in ER-dependent signaling (Campbell et al., 2001). In our study, we found that nectandrin B has weak agonistic activities on the ER α , whereas nectandrin B-induced eNOS and Akt phosphorylations are suppressed by ER antagonists. Furthermore, nectandrin B-stimulated eNOS phosphorylation was inhibited by ER α antagonist or ER α siRNA, but not by ER β blocking. These data suggest that ER α activation is critical for PI3K/Akt-mediated eNOS phosphorylation by nectandrin B.

In addition to the PI3K/Akt pathway, MAP kinase pathways have been reported to be involved in eNOS regulation. The activity of ERK and p38 kinase was related with eNOS activation by several vasodilators (Kan et al., 2008; Grossini et al., 2009), and the JNK pathway is a downstream target kinase after eNOS activation (Go et al., 2001). Here, we found that nectandrin B activates JNK, but not ERK or p38, and confirmed that these pathways are not involved in nectandrin B-induced eNOS phosphorylation.

Although it is obvious that PI3K/Akt is a key kinase in regulating eNOS phosphorylation (Shiojima and Walsh, 2002), regulation of eNOS phosphorylation can be under the control of other kinases as well, including AMPK (Chen et al., 1999; Morrow et al., 2003), protein kinase A (Namkoong et al., 2009), and protein kinase C (Michell et al., 2001). AMPK is a serine/threonine protein kinase that is a critical mediator of energy metabolism (Hardie and Hawley, 2001; Carling, 2004). We showed that overexpression of the constitutively active form of AMPK alone was enough to increase eNOS phosphorylation in endothelial cells, indicating that AMPK also functions as an eNOS activator in our system. Moreover, nectandrin B potently activated AMPK, and this activation is closely associated with eNOS phosphorylation, which was shown using chemical inhibitors or a dominant-negative mutant of AMPK. Previous studies demonstrated that CaMKs and LKB1 act upstream of the AMPK pathway (Woods et al., 2005) and regulate eNOS phosphorylation (Mount et al., 2008; Bair et al., 2009). In our study, CaMK II phosphorylation was increased by nectandrin B, and, furthermore, CaMK II inhibitors significantly inhibit nectandrin B-induced phosphorylation of eNOS as well as AMPK. These results indicate that nectandrin B stimulates eNOS phosphorylation via the AMPK pathway, presumably through CaMKII activation. Levine et al. (2007) showed that AMPK lies upstream of Akt in the pathway leading from receptor activation to eNOS stimulation (Dimmeler et al., 1999). Because inhibition of either PI3K or AMPK simultaneously blocks nectandrin B-




Fig. 10. Role of ER/PI3K and AMPK pathways in nectandrin B-mediated vasorelaxation. **A**, effects of inhibitors targeting ER, PI3K, and AMPK on nectandrin B-mediated vasorelaxation. Endothelium-intact aortic rings were anchored to the organ chamber and preincubated with or without LY294002 (LY; 10 μ M), ICI-182780 (ICI; 100 nM), or compound C (20 μ M) for 30 min, and 10 μ g/ml nectandrin B-mediated vascular relaxation was monitored in the precontracted aortic rings by 1 μ M phenylephrine. Data are expressed as relaxation percentage relative to acetylcholine (ACh; 1 μ M)-mediated relaxation and represent the means \pm S.D. of four separate experiments (*, $p < 0.05$, significant compared with nectandrin B-treated sample; control level = 1). **B**, effect of nectandrin B on eNOS phosphorylation in aortic rings. Rat aortic rings were incubated with 10 μ M nectandrin B for the indicated time (5–90 min), and the homogenates of aortic rings were subjected to phosphorylated eNOS immunoblotting. L-NAME, N^G -nitro-L-arginine methyl ester.

stimulated eNOS phosphorylation, we further tested whether cross-talk between PI3K/Akt and AMPK is associated with eNOS phosphorylation by nectandrin B. We found that AKT-phosphorylation induced by nectandrin B was attenuated by compound C, but that LY294002 did not affect nectandrin B-mediated AMPK phosphorylation. In addition, compound C decreased the ER reporter activity stimulated by nectandrin B. Thus, we can conclude that nectandrin B first activates AMPK and that this may be responsible for further serial activation of the ER α /PI3K/Akt pathway and subsequent eNOS phosphorylation.

In an organ chamber study, we found that 10 μ g/ml nectandrin B potentially relaxed rat aortic rings, and inhibitors against ER or PI3K completely suppressed nectandrin B-stimulated vascular relaxation. Although compound C also results in significant inhibition, the inhibition intensity was marginal compared with LY294002 and ICI-182780. The discrepancy between the effectiveness of compound C in endothelial cell culture and that in organ chamber system may be due to the incomplete tissue uptake of compound C in the rat aortic rings.

In a previous study, 200 mg/kg tetrahydrofuran mixture of *M. fragrans*, which contains seven lignan compounds including nectandrin B, was daily administered in C57/BL6 mice for 6 weeks. The mixture showed antiobesity effects against high-fat diet, but it caused no toxic response in mice (Nguyen et al., 2010). We also found that oral administration of 20 mg/kg nectandrin B in C57/BL6 mice for 6 weeks made no significant change in the organs of mice (data not shown). Considering relatively long treatment schedules in these studies, we believe 1 to 10 μ g/ml nectandrin B seems to be a

clinically relevant concentration range in vivo. In summary, the present study shows that nectandrin B activates eNOS via eNOS phosphorylation and that the ER/PI3K/Akt pathway under the control of AMPK plays a critical role in nectandrin B-mediated eNOS phosphorylation. Nectandrin B would be applicable to prevent cardiovascular diseases and our observations have important implications for the elucidation of the pharmacological mechanism of nectandrin B.

Authorship Contributions

Participated in research design: Lee and Kang.

Conducted experiments: Hien, Hung, and S. J. Oh.

Contributed new reagents or analytic tools: W. K. Oh.

Performed data analysis: Hien and Lee.

Wrote or contributed to the writing of the manuscript: Hien and Kang.

References

- Afrose S, Hossain MS, Maki T, and Tsujii H (2009) Karaya root saponin exerts a hypocholesterolemic response in rats fed a high-cholesterol diet. *Nutr Res* **29**:350–354.
- Anggakusuma, Yanti, and Hwang JK (2010) Effects of macelignan isolated from *Myristica fragrans* Houtt. on UVB-induced matrix metalloproteinase-9 and cyclooxygenase-2 in HaCaT cells. *J Dermatol Sci* **57**:114–122.
- Bair AM, Thippagowda PB, Freichel M, Cheng N, Ye RD, Vogel SM, Yu Y, Flockerzi V, Malik AB, and Tiruppathi C (2009) Ca^{2+} entry via TRPC channels is necessary for thrombin-induced NF- κ B activation in endothelial cells through AMP-activated protein kinase and protein kinase C δ . *J Biol Chem* **284**:563–574.
- Bradley EA, Eringa EC, Stehouwer CD, Korstjens I, van Nieuw Amerongen GP, Musters R, Sipkema P, Clark MG, and Rattigan S (2010) Activation of AMP-activated protein kinase by 5-aminoimidazole-4-carboxamide-1-beta-D-ribofuranoside in the muscle microcirculation increases nitric oxide synthesis and microvascular perfusion. *Arterioscler Thromb Vasc Biol* **30**:1137–1142.
- Cai H, Li Z, Davis ME, Kanner W, Harrison DG, and Dudley SC Jr (2003) Akt-dependent phosphorylation of serine 1179 and mitogen-activated protein kinase/extracellular signal-regulated kinase 1/2 cooperatively mediate activation

- of the endothelial nitric-oxide synthase by hydrogen peroxide. *Mol Pharmacol* **63**:325–331.
- Campbell RA, Bhat-Nakshatri P, Patel NM, Constantinidou D, Ali S, and Nakshatri H (2001) Phosphatidylinositol 3-kinase/AKT-mediated activation of estrogen receptor alpha: a new model for anti-estrogen resistance. *J Biol Chem* **276**:9817–9824.
- Carling D (2004) The AMP-activated protein kinase cascade—a unifying system for energy control. *Trends Biochem Sci* **29**:18–24.
- Carling D, Sanders MJ, and Woods A (2008) The regulation of AMP-activated protein kinase by upstream kinases. *Int J Obes (Lond)* **32** (Suppl 4):S55–S59.
- Chakrabarti S and Davidge ST (2009) High glucose-induced oxidative stress alters estrogen effects on ERalpha and ERbeta in human endothelial cells: reversal by AMPK activator. *J Steroid Biochem Mol Biol* **117**:99–106.
- Chen ZP, Mitchellhill KI, Michell BJ, Stapleton D, Rodriguez-Crespo I, Witters LA, Power DA, Ortiz de Montellano PR, and Kemp BE (1999) AMP-activated protein kinase phosphorylation of endothelial NO synthase. *FEBS Lett* **443**:285–289.
- Colombo SL and Moncada S (2009) AMPKalpha1 regulates the antioxidant status of vascular endothelial cells. *Biochem J* **421**:163–169.
- Dimmeler S, Fleming I, Fisslthaler B, Hermann C, Busse R, and Zeiher AM (1999) Activation of nitric oxide synthase in endothelial cells by Akt-dependent phosphorylation. *Nature* **399**:601–605.
- Dong Y, Zhang M, Wang S, Liang B, Zhao Z, Liu C, Wu M, Choi HC, Lyons TJ, and Zou MH (2010) Activation of AMP-activated protein kinase inhibits oxidized LDL-triggered endoplasmic reticulum stress in vivo. *Diabetes* **59**:1386–1396.
- Fatehi-Hassanabad Z, Chan CB, and Furman BL (2010) Reactive oxygen species and endothelial function in diabetes. *Eur J Pharmacol* **636**:8–17.
- Formoso G, Chen H, Kim JA, Montagnani M, Consoli A, and Quon MJ (2006) Dehydroepiandrosterone mimics acute actions of insulin to stimulate production of both nitric oxide and endothelin 1 via distinct phosphatidylinositol 3-kinase- and mitogen-activated protein kinase-dependent pathways in vascular endothelium. *Mol Endocrinol* **20**:1153–1163.
- Fujimoto N, Kohta R, and Kitamura S (2004) Estrogenic activity of an antioxidant, nordihydroguaiaretic acid (NDGA). *Life Sci* **74**:1417–1425.
- Fulton DA, Gratton JP, McCabe TJ, Fontana J, Fujio Y, Walsh K, Franke TF, Papapetropoulos A, and Sessa WC (1999) Regulation of endothelium-derived nitric oxide production by the protein kinase Akt. *Nature* **399**:597–601.
- Go YM, Levonen AL, Moellering D, Ramachandran A, Patel RP, Jo H, and Darley-Usmar VM (2001) Endothelial NOS-dependent activation of c-Jun NH(2)-terminal kinase by oxidized low-density lipoprotein. *Am J Physiol Heart Circ Physiol* **281**:H2705–H2713.
- Gokce N, Keaney JF Jr, Hunter LM, Watkins MT, Nedeljkovic ZS, Menzoian JO, and Vita JA (2003) Predictive value of noninvasively determined endothelial dysfunction for long-term cardiovascular events in patients with peripheral vascular disease. *J Am Coll Cardiol* **41**:1769–1775.
- Grossini E, Molinari C, Caimmi PP, Uberti F, and Vacca G (2009) Levosimendan induces NO production through p38 MAPK, ERK and Akt in porcine coronary endothelial cells: role for mitochondrial K(ATP) channel. *Br J Pharmacol* **156**:250–261.
- Grossini E, Molinari C, Mary DA, Uberti F, Caimmi PP, Surico N, and Vacca G (2008) Intracoronary genistein acutely increases coronary blood flow in anesthetized pigs through beta-adrenergic mediated nitric oxide release and estrogenic receptors. *Endocrinology* **149**:2678–2687.
- Grover JK, Khandkar S, Vats V, Dhunoo Y, and Das D (2002) Pharmacological studies on *Myristica fragrans*—antidiarrheal, hypnotic, analgesic and hemodynamic (blood pressure) parameters. *Methods Find Exp Clin Pharmacol* **24**:675–680.
- Han CY, Cho KB, Choi HS, Han HK, and Kang KW (2008a) Role of FoxO1 activation in MDR1 expression in adriamycin-resistant breast cancer cells. *Carcinogenesis* **29**:1837–1844.
- Han KL, Choi JS, Lee JY, Song J, Joe MK, Jung MH, and Hwang JK (2008b) Therapeutic potential of peroxisome proliferators-activated receptor-alpha/gamma dual agonist with alleviation of endoplasmic reticulum stress for the treatment of diabetes. *Diabetes* **57**:737–745.
- Hardie DG (2007) AMP-activated protein kinase as a drug target. *Annu Rev Pharmacol Toxicol* **47**:185–210.
- Hardie DG and Hawley SA (2001) AMP-activated protein kinase: the energy charge hypothesis revisited. *Bioessays* **23**:1112–1119.
- Haynes MP, Li L, Sinha D, Russell KS, Hisamoto K, Baron R, Collinge M, Sessa WC, and Bender JR (2003) Src kinase mediates phosphatidylinositol 3-kinase/Akt-dependent rapid endothelial nitric-oxide synthase activation by estrogen. *J Biol Chem* **278**:2118–2123.
- Howes LG, Abbott D, and Straznicki NE (1997) Lipoproteins and cardiovascular reactivity. *Br J Clin Pharmacol* **44**:319–324.
- Hurley RL, Anderson KA, Franzoni JM, Kemp BE, Means AR, and Witters LA (2005) The Ca²⁺/calmodulin-dependent protein kinase kinases are AMP-activated protein kinase kinases. *J Biol Chem* **280**:29060–29066.
- Kan WH, Hsu JT, Ba ZF, Schwacha MG, Chen J, Choudhry MA, Bland KI, and Chaudry IH (2008) p38 MAPK-dependent eNOS upregulation is critical for 17beta-estradiol-mediated cardioprotection following trauma-hemorrhage. *Am J Physiol Heart Circ Physiol* **294**:H2627–H2636.
- Katsuda S, Suzuki K, Koyama N, Takahashi M, Miyake M, Hazama A, and Takazawa K (2009) Safflower seed polyphenols (N-(p-coumaroyl)serotonin and N-feruloylserotonin) ameliorate atherosclerosis and distensibility of the aortic wall in Kurosawa and Kusanagi-hypercholesterolemic (KHC) rabbits. *Hypertens Res* **32**:944–949.
- Kwon HS, Kim MJ, Jeong HJ, Yang MS, Park KH, Jeong TS, and Lee WS (2008) Low-density lipoprotein (LDL)-antioxidant lignans from *Myristica fragrans* seeds. *Bioorg Med Chem Lett* **18**:194–198.
- Levine YC, Li GK, and Michel T (2007) Agonist-modulated regulation of AMP-activated protein kinase (AMPK) in endothelial cells. Evidence for an AMPK → Rac1 → Akt → endothelial nitric-oxide synthase pathway. *J Biol Chem* **282**:20351–20364.
- Li H and Förstermann U (2000) Nitric oxide in the pathogenesis of vascular disease. *J Pathol* **190**:244–254.
- Ma J, Hwang YK, Cho WH, Han SH, Hwang JK, and Han JS (2009) Macelignan attenuates activations of mitogen-activated protein kinases and nuclear factor kappa B induced by lipopolysaccharide in microglial cells. *Biol Pharm Bull* **32**:1085–1090.
- Magnone M, Bruzzone S, Guida L, Damonte G, Millo E, Scarfi Si C, Sturla L, Palombo D, De Flora A, and Zocchi E (2009) Abscissic acid released by human monocytes activates monocytes and vascular smooth muscle cell responses involved in atherogenesis. *J Biol Chem* **284**:17808–17818.
- Matsumoto T, Noguchi E, Ishida K, Kobayashi T, Yamada N, and Kamata K (2008) Metformin normalizes endothelial function by suppressing vasoconstrictor prostanooids in mesenteric arteries from OLETF rats, a model of type 2 diabetes. *Am J Physiol Heart Circ Physiol* **295**:H1165–H1176.
- Merla R, Ye Y, Lin Y, Manickavasagam S, Huang MH, Perez-Polo RJ, Uretsky BF, and Birnbaum Y (2007) The central role of adenosine in statin-induced ERK1/2, Akt, and eNOS phosphorylation. *Am J Physiol Heart Circ Physiol* **293**:H1918–H1928.
- Michell BJ, Chen Zp, Tiganis T, Stapleton D, Katsis F, Power DA, Sim AT, and Kemp BE (2001) Coordinated control of endothelial nitric-oxide synthase phosphorylation by protein kinase C and the cAMP-dependent protein kinase. *J Biol Chem* **276**:17625–17628.
- Mineo C and Shaul PW (2006) Circulating cardiovascular disease risk factors and signaling in endothelial cell caveolae. *Cardiovasc Res* **70**:31–41.
- Misra P (2008) AMP activated protein kinase: a next generation target for total metabolic control. *Expert Opin Ther Targets* **12**:91–100.
- Moncada S and Higgs A (1993) The L-arginine-nitric oxide pathway. *N Engl J Med* **329**:2002–2012.
- Morcos M, Borcea V, Isermann B, Gehrke S, Ehret T, Henkels M, Schiekofer S, Hofmann M, Amiraj J, Tritschler H, et al. (2001) Effect of alpha-lipoic acid on the progression of endothelial cell damage and albuminuria in patients with diabetes mellitus: an exploratory study. *Diabetes Res Clin Pract* **52**:175–183.
- Morrow VA, Foufelle F, Connell JM, Petrie JR, Gould GW, and Salt IP (2003) Direct activation of AMP-activated protein kinase stimulates nitric-oxide synthesis in human aortic endothelial cells. *J Biol Chem* **278**:31629–31639.
- Mount PF, Lane N, Venkatesan S, Steinberg GR, Fraser SA, Kemp BE, and Power DA (2008) Bradykinin stimulates endothelial cell fatty acid oxidation by CaMKK-dependent activation of AMPK. *Atherosclerosis* **200**:28–36.
- Namkoong S, Kim CK, Cho YL, Kim JH, Lee H, Ha KS, Choe J, Kim PH, Won MH, Kwon YG, et al. (2009) Forskolin increases angiogenesis through the coordinated cross-talk of PKA-dependent VEGF expression and Epac-mediated PI3K/Akt/eNOS signaling. *Cell Signal* **21**:906–915.
- Nguyen PH, Le TV, Kang HW, Chae J, Kim SK, Kwon KI, Seo DB, Lee SJ, and Oh WK (2010) AMP-activated protein kinase (AMPK) activators from *Myristica fragrans* (nutmeg) and their anti-obesity effect. *Bioorg Med Chem Lett* **20**:4128–4131.
- Schalkwijk CG and Stehouwer CD (2005) Vascular complications in diabetes mellitus: the role of endothelial dysfunction. *Clin Sci* **109**:143–159.
- Sharma A, Mathur R, and Dixit VP (1995) Prevention of hypercholesterolemia and atherosclerosis in rabbits after supplementation of *Myristica fragrans* seed extract. *Indian J Physiol Pharmacol* **39**:407–410.
- Shiojima I and Walsh K (2002) Role of Akt signaling in vascular homeostasis and angiogenesis. *Circ Res* **90**:1243–1250.
- Simoncini T, Hafezi-Moghadam A, Brazil DP, Ley K, Chin WW, and Liao JK (2000) Interaction of oestrogen receptor with the regulatory subunit of phosphatidylinositol-3-OH kinase. *Nature* **407**:538–541.
- Thomas SR, Chen K, and Keaney JF Jr (2002) Hydrogen peroxide activates endothelial nitric-oxide synthase through coordinated phosphorylation and dephosphorylation via a phosphoinositide 3-kinase-dependent signaling pathway. *J Biol Chem* **277**:6017–6024.
- Widlansky ME, Gokce N, Keaney JF Jr, and Vita JA (2003) The clinical implications of endothelial dysfunction. *J Am Coll Cardiol* **42**:1149–1160.
- Woods A, Dickerson K, Heath R, Hong SP, Momcilovic M, Johnstone SR, Carlson M, and Carling D (2005) Ca²⁺/calmodulin-dependent protein kinase kinase-beta acts upstream of AMP-activated protein kinase in mammalian cells. *Cell Metab* **2**:21–33.

Address correspondence to: Dr. Keon Wook Kang, College of Pharmacy and Research Institute of Pharmaceutical Sciences, Seoul National University, Seoul 151-742, Republic of Korea. E-mail: kwkang@snu.ac.kr