

## PERSPECTIVE

# Casting a Wider Net: Whole-Cell Assays to Capture Varied and Biased Signaling

Terry Kenakin

Department of Pharmacology University of North Carolina School of Medicine, Chapel Hill, North Carolina

Received July 9, 2012; accepted July 24, 2012

### ABSTRACT

The observation of complex receptor behaviors has shown how ligands can have multiple efficacies and can also differentially stimulate certain cellular signaling pathways over others (i.e., biased signaling). Conventional pharmacological assays (usually proximal to the receptor) will detect ligands that produce the signal defined by the assay (i.e.,  $\text{Ca}^{2+}$ , cAMP, and others) but otherwise may miss biased ligands that produce little activation of pathways not measured by the assay. In theory, this is less of a hazard for generic whole-cell assays, which may be sensitive to multiple signaling inputs. Whole-cell assays have the advantage of de-

tecting effects induced by a variety of receptor interactions with cytosolic proteins, including those that may be previously unknown. These ideas are discussed within the context of the high-throughput flow cytometry measurement of receptor internalization described by Wu et al. in the current issue of the journal (p. 645). The internalization of receptors can be a useful therapeutic modality and the article by Wu et al. illustrates how this new assay, targeted to downstream cellular effects, can uncover unique ligand efficacies linked to receptor internalization.

### Introduction

The two characteristic properties of all drugs are *affinity* for a biological target (this gets them to the target) and *efficacy* to change the behavior of that target (what they do when they get there). The concept of drug efficacy has changed dramatically over the past 15 to 20 years, and a major reason for this revolution is the increasing availability of multiple receptor and cellular assays that allow observation of receptor behavior through numerous vantage points. The pharmacologists' window into drug efficacy is the assay; the type of assay used actually defines the efficacy that is observed. With this increase in the number of ways to monitor receptors has come an appreciation of the diverse behaviors these proteins have in their repertoire.

### How Assay Technology Changed Pharmacology

The new views of receptor behavior afforded by new assay technology have introduced two important ideas to pharmacology. The first is that drugs can have many and diverse efficacies; this behavior has been termed *pluridimensional efficacy* (Galandrin and Bouvier, 2006). For example, the  $\beta$ -adrenoceptor ligand propranolol produces inverse agonism for cAMP but positive agonism for ERK1/2 activation (Azzi et al., 2003; Baker et al., 2003). The cannabinoid ligand desacetyllevonantradole is a positive agonist for  $G_{i1}$  and  $G_{i2}$  but an inverse agonist for  $G_{i3}$  (Mukhopadhyay and Howlett, 2005). A diverse range of positive and negative agonism has been found for  $\beta$ -adrenoceptor ligands tested in adenylate cyclase versus mitogen-activated protein kinase assays (Galandrin and Bouvier, 2006).

The second receptor behavior relevant to drug discovery is the observation that ligands need not produce identical effects on receptors but rather can *bias stimulus* through receptors to different signaling pathways in the cell (for a review, see Kenakin, 2010a). This has clearly been shown for a number of

Article, publication date, and citation information can be found at <http://molpharm.aspetjournals.org>.  
<http://dx.doi.org/10.1124/mol.112.081117>.  
Please see the related article on page 645–657.

**ABBREVIATIONS:** ERK1/2, extracellular signal-regulated protein kinases 1 and 2; 7TM, seven-transmembrane; PTH, parathyroid hormone; RANTES, regulated on activation normal T cell expressed and secreted; TAK652, (S)-8-[4-(2-butoxyethoxy)phenyl]-1-isobutyl-N-(4-[(1-propyl-1H-imidazol-5-yl)-methyl]sulfinyl)phenyl)-1,2,3,4-tetrahydro-1-benzazocine-5-carboxamide monomethanesulfonate.

seven-transmembrane (7TM) receptor functions, notably G protein signaling versus  $\beta$ -arrestin recruitment to the receptor. For example, this has been reported for biased ligands such as [Trp<sup>1</sup>]PTHrp-(1–36) (G protein-biased) versus PTH-1A ([D-Trp<sup>12</sup>, Tyr<sup>34</sup>]PTH-(7–34);  $\beta$ -arrestin-biased) for the PTH receptor (Gesty-Palmer et al., 2006), the substance P analog SpD ([D-Arg<sup>1</sup>, D-Phe<sup>5</sup>, D-Trp<sup>7,9</sup>, Leu<sup>11</sup>]substance P; G protein-dependent response) versus bombesin (G protein-independent  $\beta$ -arrestin activity) (MacKinnon et al., 2001), and SII (Sar<sup>1</sup>, Ile<sup>4</sup>, Ile<sup>8</sup>-AngII) (Ahn et al., 2004) [biased toward producing activation of the  $\beta$ -arrestin pathway versus angiotensin II (Wei et al., 2003)].

It is difficult to separate the concepts of *pluridimensional efficacy* and *biased signaling* because both often involve the stabilization of ligand-specific receptor conformations (see below). In fact, the discovery of different efficacies for known molecules is directly linked to the testing of those molecules in different assays. This idea leads to the general concept that the discovery of new efficacies and the reduction of the likelihood of missing efficacy through biased signaling may be linked to the capability of the assay to detect signaling sequelae that result from varied receptor conformational active states. The number of unique types of molecules detected in different assays can easily exceed the number of assays used for screening. For example, it would be predicted that the testing of molecules in two assays such as a signaling assay and receptor internalization assay can detect up to seven different types of molecular efficacy (Kenakin, 2005). It is not the number of assays per se that is relevant but rather the range of detectors of molecular events that matters. From this point of view, cells furnish a complete set of interrogators of receptor conformation in the form of cytosolic signaling proteins (Magalhaes et al., 2012) (Fig. 1), thus, in turn, optimizing the probability of detecting new and useful efficacies for molecules.


These ideas suggest that if a specific assay is chosen for detection of ligands, it will see a subset of molecules that happen to have the particular efficacy monitored by the assay or ligands biased toward that particular signaling pathway. The corollary to this idea is that a whole-cell assay that monitors the outcome of multiple signaling pathways may have the advantage of detecting a range of efficacies with less

influence of bias. This idea has been effectively discussed in terms of using the phosphorylation of P42/44 mitogen-activated protein kinase (ERK1/2) as a means of detecting the receptor activation of a range of G proteins in cells (Osmond et al., 2005). In particular, a single assay of ERK1/2 activity in COS cells has been shown to detect receptor activation of  $G_{\alpha s}$ ,  $G_{\alpha q}$ , and  $G_{\beta\gamma}$  activation (Osmond et al., 2005). The measurement of cell surface receptors through fluorogen-activating protein-tagged  $\beta$ -adrenoceptors described in this issue of the journal (Wu et al., 2012) offers another approach to the use of whole-cell assays for detection of ligands with diverse and previously unknown efficacies.

## Receptor Internalization and Cell Function

Receptor systems are dynamic in that receptors are synthesized and transported to the cell surface, have a lifetime at the cell surface where they transmit information from the extracellular space to the cytosol, and then become internalized within the cytosol either to be degraded or recycled to the surface again. Before dedicated assays to view these processes became available, the desensitization and internalization of receptors were often deduced from the disappearance of agonist response. Under these circumstances, internalization necessarily was tied to the level of agonist response and the two processes were tacitly assumed to be directly linked. However, the ability to track receptor internalization as a response separate from physiological pathway stimulation showed that some antagonist ligands produce no observable physiological activation but nevertheless do actively internalize 7TM receptors (Roettger et al., 1997; Willins et al., 1998; Gray and Roth, 2001). For example, the chemokine RANTES(9–68) produces no observable CCR5 receptor activation (as seen by an absence of calcium mobilization or chemotaxis) (Gong et al., 1996), yet actively internalizes CCR5 receptors (Amara et al., 1997).

There are numerous therapeutic indications for drugs that reduce or eliminate endogenous receptor signaling. Reduction in signaling can be produced by antagonist binding to the endogenous agonist binding site (orthosteric inhibition) or allosteric binding to reduce endogenous agonist binding and/or function. Another approach is to eliminate the recep-



**Fig 1.** Receptor signaling systems. Assays can be centered on direct coupling of the receptor to various G proteins or  $\beta$ -arrestin to reveal selected efficacies or to the whole-cell response to detect varied receptor behaviors in response to ligand binding.

tor from the cell membrane through internalization; i.e., ligand-induced internalization can be a useful therapeutic modality. There are certain cases in which receptor internalization actually may be a more effective means of antagonism. Blockade of receptor signaling at the cell surface requires an appropriate target occupancy; i.e., if the antagonist diffuses away from the receptor, then endogenous agonism can resume and therefore antagonist offset kinetics is a complicating factor in the *in vivo* use of antagonists (Copeland et al., 2006; Vauquelin and Charlton, 2010). In contrast, internalization introduces a kinetic buffer into the removal of signaling in that the resumption of signaling then depends either on the rate of receptor recycling or synthesis. In particular, it would be predicted that the reemergence or synthesis of new receptors on the cell surface would be a slower process than the simple dissociation of an antagonist on the cell surface.

Direct antagonist-induced internalization of receptors provides an absolute cessation of the extracellular ligand-receptor interaction that can be useful. For example, the ability of 5-HT<sub>2A</sub> receptor antagonists to down-regulate cortical 5-HT<sub>2A</sub> receptors may play a role in the therapeutic activity of some atypical antipsychotic drugs (e.g., clozapine and olanzapine) (Yadav et al., 2011). Likewise, antagonist-induced down-regulation of 5-HT<sub>2A</sub> receptors may also play a role in protection against human polyomavirus virus infections (Elphick et al., 2004; O'Connor and Roth, 2005). Another example of therapeutically relevant receptor internalization is found in the treatment of AIDS (Mack et al., 1998). To be specific, HIV-1 uses the CCR5 receptor to infect cells, and the absence of this receptor from the cell surface through a genetic mutation that prevents expression ( $\Delta 32$  allele) leads to subsequent immunity from HIV-1 infection (Dean et al., 1996; Huang et al., 1996). This result suggests a viable option for prevention of HIV-1 infection, namely the direct internalization of CCR5 as discussed previously for RANTES(9–68).

### Whole-Cell Assays and Ligand Detection

High-throughput screens designed to detect allosteric effects can yield molecules that otherwise would not be seen in conventional screening formats (Kenakin, 2010b; Burford et al., 2011). The basic difference in these assays is the inclusion of low concentrations of endogenous agonists to determine the possible effects of cobinding synthetic ligands either to block, potentiate, or change the quality of the endogenous signal. Cases in which receptor internalization may be involved in allosteric function have been seen with antagonists that create bias in the endogenous system. For example, it has been shown that high gene copy numbers for CCL3L1 can lead to delayed onset of AIDS in HIV-1-infected patients (Gonzalez et al., 2005). Activation by chemokines leads to prevention of HIV-1 cell entry through internalization of receptors; thus, the implication for the CCL3L1 gene copy data is that high levels of CCL3L1 protect against HIV progression through CCR5 internalization. In fact, chemokine-induced CCR5 internalization is a strategy for the use of PSC-RANTES as a treatment against HIV-1 infection (Hartley et al., 2004). Therefore, a biased antagonist that blocks HIV-1 interaction with the CCR5 receptor, but allows chemokines, such as CCL3L1, to internalize receptors, theoretically offers a therapeutic advantage over nonbiased antagonists. An example of this type of ligand is (*S*)-8-[4-(2-butoxyethoxy)phenyl]-1-isobutyl-*N*-(4-[(1-propyl-1*H*-imidazol-

5-yl)methyl]sulfinyl)phenyl)-1,2,3,4-tetrahydro-1-benzazocine-5-carboxamide monomethanesulfonate (TAK652), a biased antagonist that selectively blocks HIV-1 while preserving chemokine function [this antagonist is 11–12 times more potent at blocking HIV-1 entry than blockade of chemokine-induced CCR5 internalization (Muniz-Medina et al., 2009)].

The simplest mechanism to account for biased agonism and antagonism is ligand-induced stabilization of unique receptor conformational active states (Kenakin, 1995). Within this theoretical framework, ligands would stabilize conformations that are predisposed to cellular internalization; a possible mechanism for this could be ligand-directed control of receptor phosphorylation. This has been described as a phosphorylation-mediated “barcoding” of receptors that is dependent on receptor conformation. The fact that receptors are barcoded through unique intracellular patterns of phosphorylation leads to long-term programming of the receptor's behavior (Tobin, 2008; Tobin et al., 2008; Butcher et al., 2011; Nobles et al., 2011). An example of this effect is seen with the  $\mu$ -opioid receptor agonists [D-Ala<sup>2</sup>,*N*-Me-Phe<sup>4</sup>,Gly<sup>5</sup>-ol]-enkephalin and etonitazene, which stimulate the phosphorylation of both Thr370 and Ser375; in contrast, morphine leads to phosphorylation of only Thr370 (Doll et al., 2011). One of the most prominent outcomes of these effects is receptor internalization and subsequent cytosolic reactions (i.e., degradation or recycling to the cell surface).

In general, the more an assay depends on cell complexity the more it will differentiate subtle ligand effects. Downstream assays such as cell sorting are an effective way of detecting the overall outcome of ligand-directed conformational stabilization (i.e., texture in efficacy). An assay such as that reported by Wu et al. (2012) is a way to observe the end-organ response of these interactions without the need to define the mechanism. The importance of these types of discovered textures in efficacy is underscored by the reports of previously unknown cytosolic protein interactants with 7TM receptors that may lead to alteration of receptor function and disposition (Magalhaes et al., 2012). For example, desipramine binds to  $\alpha_2$ -adrenoceptors but produces no overt  $\alpha_2$ -adrenoceptor response. However, although conventional signaling assays show no observable effect of desipramine, recently it has been shown that this ligand causes  $\beta$ -arrestin-3 recruitment to the receptor (with no  $\alpha_2$ -adrenoceptor signaling) and that this effect is linked to internalization of  $\alpha_2$ -adrenoceptors. This efficacy may be associated with a useful clinical phenotypic profile in depressive disorders (Cottingham et al., 2011). As new assays become available, it would seem reasonable to subject known drugs to new interrogations to possibly link interesting *in vivo* drug phenotypes with measurable *in vitro* activities (O'Connor and Roth, 2005).

Consideration of the multiple conformational nature of receptor systems through molecular dynamics (Frauenfelder et al., 1988, 1991; Woodward, 1993; Hilser and Freire, 1997; Kenakin, 2002; Hilser et al., 2006) predicts that signaling bias and texture in efficacy should be a common, not rare, effect. In particular, uniformity in efficacy would require that ligands have identical affinities for numerous receptor conformations; this would be an unlikely thermodynamic prediction. Therefore, some kind of ligand bias would be predicted for most, if not all, synthetic ligands. The detection of such bias is directly related to the assays used to observe drug

effect, a situation analogous to the discovery and progression of ideas related to inverse agonism. When first reported by Costa and Herz (1989), inverse agonism was thought to be rare. The “rarity” of inverse agonism seems to have been related to the lack of availability of constitutive receptor assays, because in the years subsequent to the discovery, the reports of increasingly common inverse agonism have paralleled the availability of constitutive receptor assays (Kenakin, 2004). It will be interesting to see whether biased agonism and antagonism follow the same progression; i.e., as more experience is gained with high-throughput assays capable of detecting complex behaviors, will bias signaling in ligands be seen to be a more common phenomenon?

#### Authorship Contributions

Wrote or contributed to the writing of the manuscript: Kenakin.

#### References

- Ahn S, Shenoy SK, Wei H, and Lefkowitz RJ (2004) Differential kinetic and spatial patterns of  $\beta$ -arrestin and G-protein-mediated ERK activation by the angiotensin II receptor. *J Biol Chem* **279**:35518–35525.
- Amara A, Gall SL, Schwartz O, Salamero J, Montes M, Loetscher P, Baggiolini M, Virelizier JL, and Arenzana-Seisdedos F (1997) HIV coreceptor downregulation as antiviral principle: SDF-1 $\alpha$ -dependent internalization of the chemokine receptor CXCR4 contributes to inhibition of HIV replication. *J Exp Med* **186**:139–146.
- Azzi M, Charest PG, Angers S, Rousseau G, Kohout T, Bouvier M, and Piñeyro G (2003)  $\beta$ -Arrestin-mediated activation of MAPK by inverse agonists reveals distinct active conformations for G-protein-coupled receptors. *Proc Natl Acad Sci USA* **100**:11406–11411.
- Baker JG, Hall IP, and Hill SJ (2003) Agonist and inverse agonist actions of  $\beta$ -blockers at the human  $\beta_2$ -adrenoceptor provide evidence for agonist-directed signaling. *Mol Pharmacol* **64**:1357–1369.
- Burford NT, Watson J, Bertekap R, and Alt A (2011) Strategies for the identification of allosteric modulators of G-protein-coupled receptors. *Biochem Pharmacol* **81**: 691–702.
- Butcher AJ, Prihandoko R, Kong KC, McWilliams P, Edwards JM, Bottrill A, Mistry S, and Tobin AB (2011) Differential G-protein-coupled receptor phosphorylation provides evidence for a signaling bar code. *J Biol Chem* **286**:11506–11518.
- Copeland RA, Pompliano DL, and Meek TD (2006) Drug-target residence time and its implications for lead optimization. *Nat Rev Drug Discov* **5**:730–739.
- Costa T and Herz A (1989) Antagonists with negative intrinsic activity at  $\delta$ -opioid receptors coupled to GTP-binding proteins. *Proc Natl Acad Sci USA* **86**:7321–7325.
- Cottingham C, Chen Y, Jiao K, and Wang Q (2011) The antidepressant desipramine is an arrestin-biased ligand at the  $\alpha_2A$ -adrenergic receptor driving receptor down-regulation in vitro and in vivo. *J Biol Chem* **286**:36063–36075.
- Dean M, Carrington M, Winkler C, Huttley GA, Smith MW, Allikmets R, Goedert JJ, Buchbinder SP, Vittinghoff E, Gomperts E, Donfield S, Vlahov D, Kaslow R, Saah A, Rinaldo C, Detels R, and O'Brien SJ (1996) Genetic restriction of HIV-1 infection and progression to AIDS by a deletion allele of the CCR5 structural gene. Hemophilia Growth and Development Study, Multicenter AIDS Cohort Study, Multicenter Hemophilia Cohort Study, San Francisco City Cohort, ALIVE Study [published erratum appears in *Science* **274**:1069, 1997]. *Science* **273**:1856–1862.
- Doll C, Konietzko J, Pöhl F, Koch T, Höllt V, and Schulz S (2011) Agonist-selective patterns of  $\mu$ -opioid receptor phosphorylation revealed by phosphosite-specific antibodies. *Br J Pharmacol* **164**:298–307.
- Elphick GF, Querbes W, Jordan JA, Gee GV, Eash S, Manley K, Dugan A, Stanifer M, Bhatnagar A, Kroeze WK, Roth BL, and Atwood WJ (2004) The human polyomavirus, JCV, uses serotonin receptors to infect cells. *Science* **306**:1380–1383.
- Frauenfelder H, Parak F, and Young RD (1988) Conformational substates in proteins. *Annu Rev Biophys Biophys Chem* **17**:451–479.
- Frauenfelder H, Sligar SG, and Wolynes PG (1991) The energy landscapes and motions of proteins. *Science* **254**:1598–1603.
- Galandrin S and Bouvier M (2006) Distinct signaling profiles of  $\beta_1$  and  $\beta_2$  adrenergic receptor ligands toward adenylyl cyclase and mitogen-activated protein kinase reveals the pluridimensionality of efficacy. *Mol Pharmacol* **70**:1575–1584.
- Gesty-Palmer D, Chen M, Reiter E, Ahn S, Nelson CD, Wang S, Eckhardt AE, Cowan CL, Spurney RF, Luttrell LM, and Lefkowitz RJ (2006) Distinct  $\beta$ -arrestin- and G protein-dependent pathways for parathyroid hormone receptor-stimulated ERK1/2 activation. *J Biol Chem* **281**:10856–10864.
- Gong JH, Ugucioni M, Dewald B, Baggiolini M, and Clark-Lewis I (1996) RANTES and MCP-3 antagonists bind multiple chemokine receptors. *J Biol Chem* **271**: 10521–10527.
- Gonzalez E, Kulkarni H, Bolivar H, Mangano A, Sanchez R, Mangano A, Sanchez R, Catano G, Nibbs RJ, Freedman BI, et al. (2005) The influence of CCL3L1 gene-containing segmental duplications on HIV-1/AIDS susceptibility. *Science* **307**: 14334–14340.
- Gray JA and Roth BL (2001) Paradoxical trafficking and regulation of 5-HT<sub>2A</sub> receptors by agonists and antagonists. *Brain Res Bull* **56**:441–451.
- Hartley O, Gaertner H, Wilken J, Thompson D, Fish R, Ramos A, Pastore C, Dufour B, Cerini F, Melotti A, Heveker N, Picard L, Alizon M, Mosier D, Kent S, and Offord R (2004) Medicinal chemistry applied to a synthetic protein: development of highly potent HIV entry inhibitors. *Proc Natl Acad Sci USA* **101**:16460–16465.
- Hilser VJ and Freire E (1997) Predicting the equilibrium protein folding pathway: structure-based analysis of staphylococcal nuclease. *Proteins* **27**:171–183.
- Hilser VJ, Garcia-Moreno E B, Oas TG, Kapp G, and Whitten ST (2006) A statistical thermodynamic model of the protein ensembles. *Chem Rev* **106**:1545–1558.
- Huang Y, Paxton WA, Wolinsky SM, Neumann AU, Zhang L, He T, Kang S, Ceradini D, Jin Z, Yazdanbakhsh K, Kunstman K, Erickson D, Dragon E, Landau NR, Phair J, Ho DD, and Koup RA (1996) The role of a mutant CCR5 allele in HIV-1 transmission and disease progression. *Nat Med* **2**:1240–1243.
- Kenakin T (1995) Agonist-receptor efficacy. II. Agonist-trafficking of receptor signals. *Trends Pharmacol Sci* **16**:232–238.
- Kenakin TP (2002) Efficacy at G protein coupled receptors. *Annu Rev Pharmacol Toxicol* **42**:349–379.
- Kenakin T (2004) Efficacy as a vector: the relative prevalence and paucity of inverse agonism. *Mol Pharmacol* **65**:2–11.
- Kenakin T (2005) New concepts in drug discovery: collateral efficacy and permissive antagonism. *Nat Rev Drug Discov* **4**:919–927.
- Kenakin T (2010a) Functional selectivity and biased receptor signaling. *J Pharmacol Exp Ther* **336**:296–302.
- Kenakin TP (2010b) Ligand detection in the allosteric world. *J Biomol Screen* **15**:119–130.
- Mack M, Luckow N, Nelson PJ, Cihak J, Simmons G, Clapham PR, Signoret N, Marsh M, Stangassinger M, Borlat F, Wells TN, Schlöndorff D, and Proudfoot AE (1998) Aminooxypentane-RANTES induces CCR5 internalization but inhibits recycling: a novel inhibitory mechanism of HIV infectivity. *J Exp Med* **187**:1215–1224.
- MacKinnon AC, Waters C, Jodrell D, Haslett C, and Sethi T (2001) Bombesin and substance-P analogues differentially regulate G-protein coupling to the bombesin receptor. Direct evidence for biased agonism. *J Biol Chem* **276**:28083–28091.
- Magalhaes AC, Dunn H, and Ferguson SS (2012) Regulation of GPCR activity, trafficking and localization by GPCR-interacting proteins. *Br J Pharmacol* **165**: 1717–1736.
- Mukhopadhyay S and Howlett AC (2005) Chemically distinct ligands promote differential CB1 cannabinoid receptor-G<sub>i</sub> protein interactions. *Mol Pharmacol* **67**: 2016–2024.
- Muniz-Medina VM, Jones S, Maglich JM, Galardi C, Hollingsworth RE, Kazmierski WM, Ferris RG, Edelstein MP, Chiswell KE, and Kenakin TP (2009) The relative activity of “function sparing” HIV-1 entry inhibitors on viral entry and CCR5 internalization: is allosteric functional selectivity a valuable therapeutic property? *Mol Pharmacol* **75**:490–501.
- Nobles KN, Xiao K, Ahn S, Shukla AK, Lam CM, Rajagopal S, Strachan RT, Huang TY, Bressler EA, Hara MR, Shenoy SK, Gygi SP, and Lefkowitz RJ (2011) Distinct phosphorylation sites on the  $\beta_2$ -adrenergic receptor establish a barcode that encodes differential functions of  $\beta$ -arrestin. *Sci Signal* **4**:ra51.
- O'Connor KA and Roth BL (2005) Finding new tricks for old drugs: an efficient route for public-sector drug discovery. *Nat Rev Drug Discov* **4**:1005–1014.
- Osmond RI, Sheehan A, Borowicz R, Barnett E, Harvey G, Turner C, Brown A, Crouch MF, and Dyer AR (2005) GPCR screening via ERK 1/2: a novel platform for screening G protein-coupled receptors. *J Biomol Screen* **10**:730–737.
- Roettger BF, Ghanekar D, Rao R, Toledo C, Yingling J, Pinon D, and Miller LJ (1997) Antagonist-stimulated internalization of the G protein-coupled cholecystokinin receptor. *Mol Pharmacol* **51**:357–362.
- Tobin AB (2008) G-protein-coupled receptor phosphorylation: where, when and by whom. *Br J Pharmacol* **153**:S167–S176.
- Tobin AB, Butcher AJ, and Kong KC (2008) Location, location, location...site-specific GPCR phosphorylation offers a mechanism for cell-type-specific signalling. *Trends Pharmacol Sci* **29**:413–420.
- Vauquelin G and Charlton SJ (2010) Long-lasting target binding and rebinding as mechanisms to prolong in vivo drug action. *Br J Pharmacol* **161**:488–508.
- Wei H, Ahn S, Shenoy SK, Karnik SS, Hunyady L, Luttrell LM, and Lefkowitz RJ (2003) Independent  $\beta$ -arrestin 2 and G protein-mediated pathways for angiotensin II activation of extracellular signal-regulated kinases 1 and 2. *Proc Natl Acad Sci USA* **100**:10782–10787.
- Willins DL, Alsayegh L, Berry SA, Backstrom JR, Sanders-Bush E, Friedman L, Khan N, and Roth BL (1998) Serotonergic antagonist effects on trafficking of serotonin 5-HT<sub>2A</sub> receptors in vitro and in vivo. *Ann NY Acad Sci* **861**:121–127.
- Woodward C (1993) Is the slow-exchange core the protein folding core? *Trends Biochem Sci* **18**:359–360.
- Wu Y, Tapia PH, Fisher GW, Simons PC, Strouse JJ, Foutz T, Waggoner AS, Jarvik J, and Sklar LA (2012) Discovery of regulators of receptor internalization by high throughput flow cytometry. *Mol Pharmacol* **82**:645–657.
- Yadav PN, Kroeze WK, Farrell MS, and Roth BL (2011) Antagonist functional selectivity: 5-HT<sub>2A</sub> serotonin receptor antagonists differentially regulate 5-HT<sub>2A</sub> receptor protein level in vivo. *J Pharmacol Exp Ther* **339**:99–105.

**Address correspondence to:** Dr. Terry P. Kenakin, Department of Pharmacology, University of North Carolina School of Medicine, 120 Mason Farm Rd., Room 4042 Genetic Medicine Building, CB#7365, Chapel Hill, NC 27599-7365. E-mail: kenakin@email.unc.edu