

Simple and efficient synthesis of 3-aminopropenones and 3-aminopropenoates catalyzed by copper(II) nitrate trihydrate under solvent-free conditions

Geng-Chen Li

School of Material Science and Engineering, Shijiazhuang Railway Institute, Shijiazhuang, China

Received 16 October 2007; Accepted 21 October 2007; Published online 9 June 2008

© Springer-Verlag 2008

Abstract A simple, efficient and environmentally benign method has been developed for the synthesis of 3-aminopropenones and 3-aminopropenoates through the reaction of 1,3-dicarbonyl compounds with amines in the presence of $\text{Cu}(\text{NO}_3)_2 \cdot 3\text{H}_2\text{O}$ under solvent-free conditions.

Keywords 3-Aminopropenones; 3-Aminopropenoates; 1,3-Dicarbonyl compounds; Amines; Copper(II) nitrate trihydrate.


Introduction

3-Aminopropenones and 3-aminopropenoates, also known as β -enaminones and β -enamino esters are versatile intermediates for the synthesis of various heterocycles [1], natural products [2], α - and β -amino acid compounds [3], alkaloids [4], and peptides [5]. Some synthetic approaches have been reported but the direct condensation of amines with 1,3-dicarbonyl compounds is one of most preferred synthesis approaches, which is usually catalyzed by protic acid [6], *Lewis* acid such as $\text{Zn}(\text{ClO}_4)_2 \cdot 6\text{H}_2\text{O}$ [7], $\text{CeCl}_3 \cdot 7\text{H}_2\text{O}$ [8], NaAuCl_4 [9], $\text{Bi}(\text{OTf})_3$ [10], InBr_3 [11], $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ [12], *CAN* [13], $\text{Yb}(\text{OTf})_3$ [14], $\text{SnCl}_4 \cdot 5\text{H}_2\text{O}$ [15], ZrCl_4 [16], $\text{ZrOCl}_2 \cdot 8\text{H}_2\text{O}$ [17], $\text{Zn}(\text{Ac})_2 \cdot 2\text{H}_2\text{O}$ [18], $\text{Sc}(\text{OTf})_3$ [19], I_2 [20], and solid acids such as montmorillonite K10 [21],

silica chloride [22], silica gel [23], natural clays [24], and sulfated zirconia [25]. Recently, $[\text{EtNH}_3]\text{-NO}_3$ [26] and $\text{HClO}_4 \cdot \text{SiO}_2$ [27] have also been used to promote this transformation. However, some of these methodologies have not been entirely satisfactory with disadvantage, such as low yields of products, longer reaction times, harsh reaction conditions, use of harmful organic solvents, and requirement of excess of catalysts and special apparatus. Hence, there is a need to develop an efficient, practically potential, and environmentally benign method for the synthesis of 3-aminopropenones and 3-aminopropenoates under mild and economically cheaper conditions.

Recently, copper salts have emerged as efficient *Lewis* acids in organic synthesis because these compounds are relatively non-toxic, easy to handle, of low cost, and good stability [28]. Among these, copper(II) nitrate has gained special attention because this compound not only is commercially available and inexpensive, but also of high stability. Copper(II) nitrate has been reported to be an efficient reagent for many important organic reactions, such as deprotection of 1,3-dithianes and 1,3-dithiolanes [29], chemoselective synthesis of 1,1-diacetates [30], aromatization of *Hantzsch* 1,4-dihydropyridines [31], synthesis of bis(indolyl)methanes [32], *etc.* Inspired by reports of continuation of interest in catalytic applications of $\text{Cu}(\text{NO}_3)_2 \cdot 3\text{H}_2\text{O}$ for various organic transformations, we report herein a simple and convenient solvent-free method for the synthesis of 3-aminopro-

Correspondence: Geng-Chen Li, School of Material Science and Engineering, Shijiazhuang Railway Institute, Shijiazhuang 050043, China. E-mail: gengchenli@sina.com.cn



Scheme 1

penones and 3-aminopropenoates in the presence of a catalytic amount of $\text{Cu}(\text{NO}_3)_2 \cdot 3\text{H}_2\text{O}$ (Scheme 1).

Results and discussion

Initially, a model reaction of ethyl acetoacetate and aniline was performed under solvent-free conditions in the presence of a catalytic of $\text{Cu}(\text{NO}_3)_2 \cdot 3\text{H}_2\text{O}$ (10 mol%) at room temperature. To our delight, we observed the formation of ethyl 3-(phenylamino)but-2-enoate (**3i**) in 95% isolated yield after 50 min. In the absence of the catalyst, only 30% of the product was obtained even after 24 h stirring at room temperature. Increasing the amount of the catalyst did not

show significant influence on the rate of the reaction as well as yield. Lower catalyst loading can be used with only a marginal drop in reaction rate. Various solvents such as *EtOH*, *MeCN*, CH_2Cl_2 , *THF*, *DMF*, and *EtOAc* were also screened for this reaction. However, the best results were observed under solvent-free conditions.

To explore the generality and scope of this method, a wide variety of amines including primary, benzylic, and aromatic amines were reacted with various 1,3-dicarbonyl compounds and the results are summarized in Table 1. In general, primary and benzylic amines reacted with a broad range of structurally diverse 1,3-dicarbonyl compounds to afford the cor-

Table 1 Synthesis of 3-aminopropenones and 3-aminopropenoates catalyzed by $\text{Cu}(\text{NO}_3)_2 \cdot 3\text{H}_2\text{O}$

Entry	R^1	R^2	R^3	Time/min	Yield ^a /%	Ref.
a	<i>Me</i>	<i>OEt</i>	$\text{CH}_3(\text{CH}_2)_3$	15	92	[25]
b	<i>Me</i>	<i>OBn</i>	$\text{CH}_3(\text{CH}_2)_3$	18	90	[34]
c	<i>Me</i>	<i>OMe</i>	$\text{CH}_2=\text{CHCH}_2$	15	95	[12]
d	<i>Me</i>	<i>OMe</i>	C_6H_{11}	15	93	[14b]
e	<i>Me</i>	<i>OMe</i>	<i>PhCH</i> ₂	12	93	[25]
f	<i>Me</i>	<i>OBn</i>	<i>PhCH</i> ₂	18	90	[34]
g	<i>Me</i>	<i>OMe</i>	<i>PhCH</i> ₂ CH_2	15	92	[11]
h	<i>Me</i>	<i>OMe</i>	(<i>R</i>)- <i>PhCH</i> (<i>Me</i>)	15	95	[12]
i	<i>Me</i>	<i>OEt</i>	<i>Ph</i>	50	95	[19]
j	<i>Me</i>	<i>OBn</i>	<i>Ph</i>	60	90	[34]
k	<i>Me</i>	<i>OEt</i>	4- <i>Me</i> - C_6H_4	40	94	[12]
l	<i>Me</i>	<i>OMe</i>	2- <i>Me</i> - C_6H_4	90	91	[12]
m	<i>Me</i>	<i>OMe</i>	2,6- <i>Et</i> ₂ - C_6H_3	360	75	[11]
n	<i>Me</i>	<i>OEt</i>	4- <i>OMe</i> - C_6H_4	40	95	[25]
o	<i>Me</i>	<i>OEt</i>	4- <i>Cl</i> - C_6H_4	480	80	[12]
p	<i>Me</i>	<i>Me</i>	$\text{CH}_3(\text{CH}_2)_3$	20	92	[12]
q	<i>Me</i>	<i>Me</i>	$\text{H}_2\text{NCH}_2\text{CH}_2\text{CH}_2$	15	95 ^b	[11]
r	<i>Me</i>	<i>Me</i>	$\text{CH}_2=\text{CHCH}_2$	15	96	[15]
s	<i>Me</i>	<i>Me</i>	<i>Ph</i>	10	92	[15]
t	<i>Me</i>	<i>Me</i>	4- <i>Me</i> - C_6H_4	10	95	[25]
u	<i>Me</i>	<i>Me</i>	2- <i>Me</i> - C_6H_4	15	90	[25]
v	<i>Me</i>	<i>Me</i>	4- <i>OMe</i> - C_6H_4	10	91	[15]
w	<i>Me</i>	<i>Me</i>	2- <i>OEt</i> - C_6H_4	45	90	[11]
x	<i>Me</i>	<i>Me</i>	4- <i>Cl</i> - C_6H_4	300	80	[25]
y	<i>Me</i>	<i>Ph</i>	<i>Ph</i>	50	85	[15]
z	<i>Me</i>	<i>Ph</i>	2- <i>OEt</i> - C_6H_4	80	80	[25]

^a Yield refer to pure, isolated products; ^b Propane-1,3-diamine:methyl acetoacetate molar ratio = 1:2

responding β -enaminones or β -enamino esters in high yields in short time. We have also observed delicate electron effects for aromatic amine. Aniline with electron-rich groups (Table 1, entries **2k**, **2n**, **2t**, and **2v**) reacted rapidly, while substitution involving electron-withdrawing groups (Table 1, entries **2o** and **2x**) on the benzene ring decreased the reactivity requiring longer reaction times. It was also found that the substituted groups (**2l**, **2m**, **2w**, and **2z**) on the ortho position in the anilines influenced the reaction rates. Moreover, the optically active (*R*)-(+)- α -methyl benzyl amine (**2h**) was converted into the corresponding 3-aminopropenoates without any racemization or inversion confirmed by measuring its optical rotation. When 1,3-diaminopropane (**2q**) was used as an amine, two equivalents of acetylacetone were used and the product was formed with two enaminone groups. It should be pointed out that 1,3-diketones with two different substituents, such as 1-benzoylacetone, reacted with amines in a regioselective amination of the aliphatic carbonyl group (**2y** and **2z**).

The products with (*Z*)-configuration were favored over the (*E*)-isomers. It is postulated that the formation of intramolecular hydrogen bonding could be responsible for the observed shift towards the (*Z*)-isomer [33].

In conclusion, $\text{Cu}(\text{NO}_3)_2 \cdot 3\text{H}_2\text{O}$ was found to be an efficient catalyst in the reaction of 1,3-dicarbonylic compounds and amines to afford 3-aminopropenones and 3-aminopropenoates. The main advantages of this method are mild, clean, and solvent-free reaction conditions, good to excellent yields, short reaction times, and an environmentally benign reagent.

Experimental

Melting points were measured on X-4 apparatus. IR spectra were obtained on a Perkin Elmer 793 instrument. ^1H NMR spectra were recorded as CDCl_3 solutions on a Bruker spectrometer at 300 MHz using *TMS* as internal standard. Elemental analyses were performed on a PE 2400 CHNS/O Analyzer and their results agreed favourably with the calculated values.

General procedure for the synthesis of 3-aminopropenones and 3-aminopropenoates (3)

To the mixture of 10 mmol 1,3-dicarbonylic compound and 10 mmol amine, 1 mmol $\text{Cu}(\text{NO}_3)_2 \cdot 3\text{H}_2\text{O}$ was added and stir-

red vigorously at room temperature until the disappearance of the starting amine. When the reaction was complete, 20 cm^3 of ethyl acetate was added and washed with brine ($2 \times 10\text{ cm}^3$). The combined organic layer was dried (MgSO_4) and concentrated under vacuum. The residue was subjected to column chromatography over alumina using ethyl acetate (10%) in *n*-hexane to obtain the pure product. All compounds prepared proved to be identical with those already reported (Table 1) with respect to their spectroscopic properties.

Acknowledgement

The financial support to this work from the Science and Technology Research and Development Program in Hebei Province (06213507D-2) is gratefully acknowledged.

References

1. a) Dannhardt G, Bauer A, Nowe U (1998) *J Prakt Chem* 340:256; b) Elassar A-ZA, El-Khair AA (2003) *Tetrahedron* 59:8463; c) Ferraz HMC, de Oliveira EO, Payret-Arrua ME, Brandt CA (1995) *J Org Chem* 60:7357
2. Michael JP, de Koning CB, Gravestock D, Hosken GD, Howard AS, Jungmann CM, Krause RWM, Parsons AS, Pelly SC, Stanbury TV (1999) *Pure Appl Chem* 71: 979
3. a) Jerphagnon T, Renaud J-L, Demonchaux P, Ferreira A, Bruneau C (2004) *Adv Synth Catal* 346:33; b) Dai Q, Yang WR, Zhang XM (2005) *Org Lett* 7:5343
4. Michael JP, Parsons AS (1999) *Tetrahedron* 55:10915
5. Beholz LG, Benovsky P, Ward DL, Barta NS, Stille JR (1997) *J Org Chem* 62:1033
6. Martin DF, Janusonis GA, Martin BB (1961) *J Am Chem Soc* 83:73
7. Bartoli G, Bosco M, Locatelli M, Marcantoni E, Melchiorre P, Sambri L (2004) *Synlett*:239
8. Khodaei MM, Khosropour AR, Kookhazadeh M (2004) *Synlett*:1980
9. Arcadi A, Bianchi G, Di Giuseppe S, Marinelli F (2003) *Green Chem* 5:64
10. Khosropour AR, Khodaei MM, Kookhazadeh M (2004) *Tetrahedron Lett* 45:1725
11. Zhang Z-H, Yin L, Wang Y-M (2006) *Adv Synth Catal* 348:184
12. Zhang Z-H, Hu J-Y (2006) *J Braz Chem Soc* 17:1447
13. a) Mo L-P, Liu S-F, Li W-Z (2007) *J Chin Chem Soc* 54:879; b) Sridharan V, Avendaño C, Menéndez JC (2007) *Synlett*:881
14. a) Epifano F, Genovese S, Curini M (2007) *Tetrahedron Lett* 48:2717; b) Varala R, Nuvula S, Adapa SR (2006) *Aust J Chem* 59:921
15. Zhang Z-H, Ma Z-C, Mo L-P (2007) *Indian J Chem* 46B:535
16. Lin J, Zhang L-F (2007) *Monatsh Chem* 138:77
17. Zhang Z-H, Li T-S, Li J-J (2007) *Catal Commun* 8: 1615
18. Vohra RK, Renaud J-L, Bruneau C (2005) *Collect Czech Chem Commun* 70:1943

19. Yadav JS, Kumar VN, Rao RS, Priyadarshini AD, Rao PP, Reddy BVS, Nagaiah K (2006) *J Mol Catal A Chem* 256:234
20. Gogoi S, Bhuyan R, Barua NC (2005) *Synth Commun* 35:2811
21. Braibante MEF, Braibante HTS, Morel AF, Costa CC, Lima MG (2006) *J Braz Chem Soc* 17:184
22. Gholap AR, Chakor NS, Daniel T, Lahoti RJ, Srinivasa KV (2006) *J Mol Catal A: Chem* 245:37
23. Gao Y-H, Zhang Q-H, Xu J-X (2004) *Synth Commun* 34:909
24. Silva FC, De Souza MCBV, Ferreira VF, Sabino SJ, Antunes OAC (2004) *Catal Commun* 5:151
25. Zhang Z-H, Song L-M (2005) *J Chem Res*:817
26. Bhosale RS, Suryawanshi PA, Ingle SA, Lokhande MN, More SP, Mane SB, Bhosale SV, Pawar RP (2006) *Synlett*:933
27. Das B, Venkateswarlu K, Majhi A, Reddy MR, Reddy KN, Rao YK, Ravikumar K, Sridhar B (2006) *J Mol Catal A Chem* 246:276
28. a) Heravi MM, Taheri S, Bakhtiari K, Oskooie HA (2006) *Monatsh Chem* 137:1075; b) Mo L-P, Ma Z-C, Zhang Z-H (2005) *Synth Commun* 35:1997; c) Wang L, Yan JC, Li PH, Wang M, Su CN (2005) *J Chem Res*:112; d) Khan AT, Choudhury LH, Ghosh S (2004) *Tetrahedron Lett* 45:7891; e) Akhlaghinia B, Tavakoli S (2005) *Synthesis*:1775; f) Jiang H-F, Tang J-Y, Wang A-Z, Deng G-H, Yang S-R (2006) *Synthesis*:1155; g) Bhatt S, Nayak SK (2007) *Synth Commun* 37:1381; h) Nayak SK (2006) *Synth Commun* 36:1307; i) Semwal A, Nayak SK (2006) *Synth Commun* 36:227; j) Liu C, Han JY, Wang J (2007) *Synlett* 643; k) Asadollah K, Heravi MM (2007) *Monatsh Chem* 138:867
29. a) Oksdath-Mansilla G, Peñeñory AB (2007) *Tetrahedron Lett* 48:6150; b) Lee JG, Hwang JP (1995) *Chem Lett* 24:507
30. Heravi MM, Bakhtiari K, Tehrani MH, Oskooie HA (2007) *Synth Commun* 37:1121
31. Maquestiau A, Mayence A, Eynde J-JV (1991) *Tetrahedron Lett* 32:3839
32. Nasreen A, Varala R, Adapa SR (2007) *J Heterocyclic Chem* 44:983
33. Zhou J-C (1997) *Magn Reson Chem* 35:311
34. Lenin R, Raju RM (2007) *Arkivoc* Xiii:204