

Further Evidence that Behavioral Tests and Neuropeptide mRNA and Tissue Level Alterations Can Differentiate between Typical and Atypical Antipsychotic Drugs

Richard Bauer, M.D., Adelcy Mayr, M.D., Wolfgang Lederer, M.D., Patricia L. Needham, B.Sc., Ian C. Kilpatrick, Ph.D., Wolfgang W. Fleischhacker, M.D., and Josef Marksteiner, M.D.

This study was designed to compare some behavioral and biochemical effects of chronic treatment with a range of antipsychotic drugs. Gene expression of enkephalin, chromogranin A, chromogranin B, and secretogranin II and their respective peptide products were studied with in situ hybridization and radioimmunoassays after daily oral administration of haloperidol, clozapine, risperidone, or zotepine for 21 days. In behavioral tests, significant catalepsy was induced by haloperidol only. All four antipsychotic drugs increased hind paw retraction time but only haloperidol also increased forelimb retraction time. In the caudate putamen, haloperidol increased both enkephalin mRNA expression and enkephalin tissue levels. Neither of these parameters was altered by the other three drugs. In the prefrontal cortex, antipsychotic drugs generated a distinct

pattern of gene expression in two regards. First, the dopamine D₂ receptor antagonist, haloperidol, did not significantly alter synaptic protein levels or their encoding mRNAs. Secondly, there was a differential change in tissue levels and mRNA expression since secretogranin II was not affected by any tested antipsychotic drug. This study shows that different types of antipsychotic drug induce distinct behavioural effects as well as differential changes in the biosynthesis of synaptic proteins and their encoding mRNAs. The data reinforce the notion that haloperidol can be classed as a typical antipsychotic drug whilst clozapine, zotepine, and risperidone reflect their atypical classification. [Neuropsychopharmacology 23:46–55, 2000]

© 2000 American College of Neuropsychopharmacology.
Published by Elsevier Science Inc. All rights reserved

KEY WORDS: Antipsychotic drugs; Neuropeptides; Chromogranins; Gene expression

Antipsychotic drugs are sometimes referred to as “atypical” if they demonstrate the ability to produce an antipsychotic action in most patients but with signifi-

cantly less extrapyramidal side effects than classical (typical) antipsychotics (Fleischhacker and Hummer 1997). Lately, this distinction between “typical” and “atypical” antipsychotics has become the focus for much research aimed at establishing the pharmacological characteristics that serve to distinguish atypical antipsychotic drugs from the typical ones (Arnt and Skarsfeldt 1998). For example, in animals, acute and chronic treatment with these compounds causes distinct behavioral changes and alterations in the expression of different genes.

Alterations in the biosynthesis of synaptic proteins and their mRNAs can be used to investigate changes in neuronal activity following different neuronal stimuli

From the Department of Pharmacology (RB), Department of Psychiatry (AM, WWF, JM), Department of Anaesthesiology and Intensive Care (WL), University of Innsbruck, Innsbruck, Austria; and Knoll Pharmaceuticals Research & Development (PLN, ICK), Nottingham, England.

Address correspondence to: Josef Marksteiner, M.D., Department of Psychiatry, Anichstrasse 35, A-6020 Innsbruck, Austria. E-mail: j.marksteiner@uibk.ac.at

Received July 15, 1999; revised December 8, 1999; accepted December 27, 1999.

(MacArthur and Eiden 1996). Haloperidol increases tissue levels of enkephalin and the expression of mRNA (Normand et al. 1987; Romano et al. 1987) in medium spiny striatal neurons which are associated with the dopamine 2 receptor (Le Moine et al. 1991) whereas clozapine, the prototypical "atypical" antipsychotic, does not induce catalepsy and does not lead to an increased enkephalin gene expression in the striatum (Mercugliano and Chesselet 1992). Consequently, antipsychotics which do not induce catalepsy appear unlikely to increase enkephalin biosynthesis (Augood et al. 1993; Mijster et al. 1998). Therefore, both behavioral changes and a distinct pattern of gene expression may be used to predict an atypical profile for an antipsychotic compound.

The prefrontal cortex is an important brain area in schizophrenia research (Knable and Weinberger 1997; O'Donnell and Grace 1998). In animals, antipsychotic treatment causes an induction of immediate early genes in various brain areas including the prefrontal cortex. In this region, acute application of haloperidol or clozapine induces a differential expression of immediate early genes (Nguyen et al. 1992; Robertson and Fibiger 1992; Deutch and Duman 1996). Therefore, a different mechanism of action was postulated for haloperidol or clozapine treatment. However, most of these experiments investigated the induction of early genes after acute treatment.

Chromogranin A, chromogranin B, and secretogranin II belong to the chromogranin family which are large protein molecules found in large dense core vesicles. They are endoproteolytically processed to smaller peptides, and are released after neuronal stimulation (Huttner et al. 1991). The distribution of chromogranin B (Mahata et al. 1991; Kroesen et al. 1996), secretogranin II mRNA, and their immunoreactivity (Marksteiner et al. 1993a) have been studied in rat brains. The biosynthesis of chromogranins is regulated by different stimuli (Shen and Gundlach 1996). For example, ten days of treatment with clozapine or haloperidol led to significant changes in the biosynthesis of chromogranin B and secretogranin II mRNA (Kroesen et al. 1995).

In the present study, we have investigated behavioral and biochemical effects after chronic application of haloperidol, clozapine, risperidone, and zotepine. Due to their receptor binding profiles and their clinical effects, clozapine, risperidone, and zotepine are referred to as novel antipsychotics. All novel antipsychotics show a higher affinity for 5HT_{2A} than for dopamine D₂ receptors (Schotte et al. 1996). The aim of this study was three-fold. First, we were interested whether any of the test drugs could be distinguished by behavioural tests. Secondly, we investigated whether the expression pattern of chromogranins and enkephalin is suitable to predict an atypical antipsychotic profile. Lastly, we at-

tempted to correlate neuronal mRNA expression to tissue concentrations of the respective peptides.

MATERIALS AND METHODS

Adult male Sprague-Dawley rats (Versuchstierzuchtanstalt, Himberg, Austria; body weight 225–250 g) were used for all experiments. Rats were maintained on a 12:12 h light/dark cycle with food and water available *ad libitum*. All procedures were done in accordance with the Austrian Guide for the Care and Use of Laboratory Animals and were approved by the local animal care committee.

Drug Administration

Animals ($n = 16$ per treatment group and control group) received daily oral application for 21 days of vehicle (10 μ l of N glacial acetic acid per 1000 μ l of saline), haloperidol (1 mg/kg), clozapine (20 mg/kg), risperidone (1mg/kg), or zotepine (5 mg/kg). Drug doses were chosen to reflect comparable multiples of (approximately $6 \times$ ED₅₀s) for inhibition of amphetamine-induced hyperlocomotion (Leysen et al. 1993; Needham et al. 1996). The drugs were administered between 8.00 and 10.00, during the light cycle.

Behavioral Testing

All behavioral testing was performed between 8.00 and 10.00, during the light cycle. In six out of sixteen animals, a behavioral testing was performed on days 5, 10, and 20 for the catalepsy. The paw test was applied on days 4, 9, and 19. The animals which were behaviorally tested were not used for biochemical analysis.

Catalepsy. Catalepsy was tested using the standard horizontal bar method (Ezrin and Seeman 1977) at 90 minutes post-drug application. The animals were placed with the front paws on a horizontal metal bar which was suspended 10 cm above the benchtop. Time was recorded until one of the animal's paws touched the benchtop. The maximum immobility duration was set at 200 seconds.

Paw Test. Thirty and fifty minutes after the administration of the compound, each animal was submitted to the paw test. The paw test was performed as described by Ellenbroek and Cools (1988). Briefly, it was performed on a platform measuring 30×30 cm with a height of 20 cm. The platform contained four holes; two for the forelimbs and two for the hindlimbs. The rat was placed in the box and two dependent variables were

scored; (i) forelimb retraction time (FRT); and (ii) hindlimb retraction time (HRT).

Antisera

The generation of the PE-11 antiserum was described previously (Kroesen et al. 1996). Briefly, it was generated against a synthetic peptide (PE-11) corresponding to rat chromogranin B 552–562 (Forss et al. 1989) which was identical to the human sequence. The antiserum only reacts with the free C-terminal part of PE-11 since an elongated peptide (chromogranin B 552–574) only reacts in the RIA when it is first subjected to trypsin digestion. No crossreactivity was found with peptides derived from chromogranin A and secretogranin II. It did also not cross-react with the following neuropeptides: galanin, substance P, neuropeptide Y, neurotensin, and calcitonin gene-related peptide.

The secretoneurin (rat secretogranin II 154–186 (Gerdes et al. 1988) antiserum was produced in rabbit. The production and specificity of the antiserum has been described previously (Kirchmair et al. 1993; Marksteiner et al. 1993a). It did not cross-react with chromogranin B or chromogranin A in Western blots. Synthetic peptides corresponding to the N- and C-terminus of secretogranin II did not cross-react in a specific radioimmunoassay for secretoneurin. Using radioimmunoassay, no significant cross-reactivity was detected with other peptides co-stored in large dense core vesicles, i.e., peptides derived from chromogranin A (LE-40; WE-14), substance P, neuropeptide Y, and calcitonin gene related peptide.

Tissue Extraction and Radioimmunoassay

Twenty-four hours after the last drug treatment, five animals from each treatment group were decapitated and their brains were removed. The dorsolateral caudate putamen and prefrontal cortex were dissected according to the Atlas of Paxinos: (Paxinos and Watson 1986). The extraction procedure was performed as previously described (Marksteiner and Sperk 1988).

Specific radioimmunoassays for enkephalin, chromogranin B (PE-11) (Kroesen et al. 1996), and secretogranin II (secretoneurin) were performed as previously described (Marksteiner et al. 1993a) with slight modifications. Briefly, samples and standards were incubated with antiserum for 24 hours. Tracer was then added and probes were incubated for an additional 24 hours.

In Situ Hybridization

Five animals of each treatment group were killed by decapitation 12 hours after the final injection. The brains

were rapidly removed, frozen in isopentane at -65°C and stored at -70°C until processing. Coronal cryostat-cut sections (20 μm) were thaw-mounted onto gelatin-coated slides and stored at -70°C until the day of the experiment. *In situ* hybridization was performed as described previously (Marksteiner et al. 1993b). Briefly, a synthetic oligonucleotide complementary to the bases 388–435 (NEP-502; New England Nuclear, Boston, MA) of prepro-enkephalin-A mRNA (Comb et al. 1982) was used. For chromogranin A, a 51-residue oligonucleotide corresponding to amino acids 1–17 (Iacangelo et al. 1988), for chromogranin B, a 54-mer based on amino acids 1–18 (Forss et al. 1989), and for secretogranin II, a 48-residue oligonucleotide corresponding to amino acids 25–40 (Gerdes et al. 1988) was constructed. Oligonucleotides were labelled with terminal deoxynucleotidyl transferase and ^{35}S - α -thio dATP (1300 Ci/mmol; NEN, Boston, MA). After hybridization, tissue sections and membrane standards were apposed to Hyperfilm- βmax autoradiographic film (Amersham, The Netherlands). Films were developed in Kodak 19 developer. After the quantification of the autoradiographic data, sections were dipped in Kodak NTB-2 photosensitive emulsion and exposed for seven weeks. After developing, the sections were counterstained with cresyl violet. In sections pretreated with ribonuclease A (100 $\mu\text{g}/\text{ml}$), only background labelling was observed.

Quantification

Quantification was performed on emulsion-coated sections. For each region, four sections were evaluated for each animal on emulsion-coated sections. Background grain density was measured over white matter. The number of labelled neurons per section within an area of 2.25 mm^2 and the number of grains per labelled cell were counted with a grid at a magnification $\times 100$ or $\times 400$. The values were averaged for each animal. Five animals were included in each group. The mean values of untreated animals were compared to those of treated animals. Comparisons between control and treated groups were performed using one-way ANOVA followed by Dunnett's test to determine significant differences between these groups.

RESULTS

Effects of Haloperidol, Clozapine, Risperidone, and Zotepine on Behavior

As anticipated, treatment with haloperidol caused robust catalepsy in most animals (Table 1) although one animal did not demonstrate significant catalepsy. In contrast, none of the comparator antipsychotics induced catalepsy when animals were tested on days 4, 9,

Table 1. Catalepsy Scores after Antipsychotic Treatment

	4 days	19 days
Control	4.3 ± 3.7	3.5 ± 1.5
Haloperidol	35.3 ± 11.1**	46.9 ± 12.7**
Clozapine	6.1 ± 2.9	5.3 ± 3.2
Risperidone	8.4 ± 5.2	12.4 ± 5.7
Zotepine	5.1 ± 3.4	4.5 ± 2.8

Values represent seconds ± S.E.M. Significant changes are indicated by asterisks: ** $p < .01$.

and 19. In the paw test, there was a significant difference between haloperidol and the effects of clozapine, risperidone and zotepine on each test day (Table 2). These latter three agents increased HRT without altering FRT, thereby increasing the HRT:FRT ratio. On the other hand, haloperidol increased both the FRT and HRT to similar extents without significantly altering the HRT:FRT ratio. No significant differences were found between the different days of testing.

Enkephalin, Chromogranin A, Chromogranin B and Secretogranin II mRNA Expression and Respective Peptide Tissue Levels in the Caudate Putamen

In the dorsolateral caudate-putamen, haloperidol treatment significantly increased both enkephalin-A mRNA expression and tissue levels (Figure 1; Tables 3 and 4). Neither mRNA expression nor peptide levels for enkephalin, chromogranin A, chromogranin B, or secretogranin II were significantly changed after clozapine, risperidone or zotepine treatment (Tables 3 and 4).


Enkephalin, Chromogranin A, Chromogranin B and Secretogranin II mRNA Expression and Respective Peptide/Protein Tissue Levels in the Prefrontal Cortex

Enkephalin-A mRNA could not be detected above the lower assay limits within the prefrontal cortex. Chro-

Table 2. The Effects of Antipsychotic Drugs on the Forelimb Retraction Time (FRT) and Hindlimb Retraction Time (HRT) as Measured with the Paw Test (see Materials and Methods)

	FRT	HRT
Control	1 ± 0.9	2 ± 1.4
Haloperidol	13 ± 4.3**	14 ± 5**
Clozapine	1 ± 1.4	17 ± 6.5**
Risperidone	3 ± 2.0	13 ± 4.9**
Zotepine	2 ± 1.5	21 ± 7.2**

Data are represented as mean seconds ± S.E.M. obtained at day 20. ** $p < .01$ as compared to controls.

**Figure 1.** Autoradiograms for enkephalin-A mRNA in controls (a) and after haloperidol treatment at the level of rostral caudate putamen (b). Scale bar represents 1mm.

mogranin A mRNA containing neurons were preferentially found in cortical layers III and IV (Figure 2a). Chromogranin B was distributed throughout all layers with the highest number in layers III and IV (Figure 2c). Secretogranin II was found preferentially in layer II with scattered neurons in all other layers (not shown). Cellular chromogranin A mRNA expression was significantly reduced by chronic risperidone (Figures 2a and 2b; Table 3) and by clozapine treatment (Table 3). Tissue levels of chromogranin A were not determined. Risperidone (Figures 2c and 2d; Table 3) and zotepine treatment (Table 3) caused a significant decrease in the expression of chromogranin B mRNA.

Although neither clozapine nor haloperidol significantly reduced chromogranin B mRNA in prefrontal cortex, the PE-11-like immunoreactivity was significantly lowered by clozapine, risperidone, and zotepine and a strong trend to a decreased peptide level was also noted after haloperidol treatment ($p = .09$) (Table 4). Neither secretogranin II mRNA transcript levels nor secretoneurin tissue levels were significantly altered by any of the drug treatments (Tables 3 and 4). The regions

Table 3. Enkephalin, Chromogranin A, Chromogranin B and Secretogranin II mRNA Expression after Antipsychotic Treatment for 21 Days

		Enkephalin	Chr A	Chr B	Sg II
Caudate putamen	Control	100 ± 6	100 ± 9	100 ± 4	100 ± 8
	Haloperidol	127 ± 5*	107 ± 6	98 ± 2	91 ± 1
	Clozapine	102 ± 9	106 ± 10	85 ± 11	114 ± 7
	Risperidone	113 ± 2	96 ± 7	84 ± 6	99 ± 2
	Zotepine	100 ± 3	103 ± 2	98 ± 2	109 ± 7
Prefrontal cortex	Control	n.d.	100 ± 5	100 ± 4	100 ± 12
	Haloperidol	n.d.	101 ± 7	86 ± 3	77 ± 12
	Clozapine	n.d.	75 ± 2**	88 ± 5	81 ± 11
	Risperidone	n.d.	50 ± 8**	70 ± 6**	111 ± 8
	Zotepine	n.d.	85 ± 11	77 ± 4**	107 ± 13

Values represent silver grain densities (see Materials and Methods) and are expressed as percentage of control values (control ± S.E.M.).

n.d., not detected; ** $p < .01$.

analyzed by in situ hybridization and radioimmunoassay are schematically shown in Figures 3a and 3b.

DISCUSSION

The present study reinforces the contention that typical and atypical antipsychotic drugs can be distinguished by behavioural and biochemical profiles. In the present study, these have included measures of catalepsy, paw retraction times, and distinct alterations in gene expression and synaptic peptide/protein production.

Behavioural Measures

In contrast to haloperidol, none of the reportedly atypical antipsychotic drugs induced catalepsy at any time point, consistent with the reduced extrapyramidal side-

effects observed in the clinic with these drugs (Fleischhacker and Hummer 1997). In further contrast to haloperidol, the atypical drugs increased the ratio between forelimb and hindlimb retraction times, a feature that has been proposed as indicator of atypicality of these drug types (Ellenbroek and Cools 1988; Ellenbroek et al. 1994).

Neurochemical Measures

In the caudate putamen, the expression of both enkephalin mRNA and its tissue levels were increased after haloperidol but not after risperidone, clozapine or zotepine administration. These results are consistent with findings from many other groups who have demonstrated that typical—but not atypical—antipsychotics increase enkephalin content and mRNA levels in the caudate-putamen (Herman et al. 1991; Angulo 1992;

Table 4. Enkephalin-, PE-11 (Chromogranin B) and Secretoneurin (Secretogranin II) Tissue Levels after Antipsychotic Treatment

		Enkephalin	PE-11	Secretoneurin
Caudate putamen	Control	100 ± 9	100 ± 12	100 ± 15
	Haloperidol	147 ± 5**	91 ± 10	83 ± 13
	Clozapine	98 ± 7	97 ± 6	91 ± 9
	Risperidone	107 ± 6	88 ± 8	82 ± 11
	Zotepine	105 ± 9	102 ± 14	78 ± 9
Prefrontal cortex	Control	100 ± 1	100 ± 9	100 ± 14
	Haloperidol	97 ± 2	75 ± 9	98 ± 13
	Clozapine	99 ± 2	63 ± 5**	92 ± 8
	Risperidone	95 ± 1	55 ± 7**	70 ± 9
	Zotepine	101 ± 3	56 ± 7**	93 ± 10

Tissue levels (fmol/per mg wet weight) were determined 24 hours after the drug or vehicle application. In the prefrontal cortex, absolute values were for enkephalin (7 ± 1), for secretoneurin (32 ± 4.5), and for PE-11 (42 ± 3.8). In the caudate-putamen of controls, absolute values were for enkephalin (164 ± 15.3), for secretoneurin (49 ± 7.4), and for PE-11 (21 ± 2.5). Data are shown as percent of the respective control group ± S.E.M.

** $p < .01$.




Figure 2. Darkfield photomicrographs of an aspect of the medial prefrontal cortex in controls (**a, c, e**) and after risperidone treatment (**b, d, f**). Pictures are taken from a region of the prefrontal cortex shown in Figure 3. The expression of chromogranin A shown in controls (**a**) is decreased after risperidone treatment (**b**). Chromogranin B containing neurons are shown in (**c**). After risperidone treatment a decrease in mRNA expression is observed (**d**). Secretogranin II expression in controls (**e**) is not significantly reduced after antipsychotic treatment with risperidone (**f**). Scale bar represents 75 μm .

Marksteiner et al. 1993b; Mijster et al. 1996; Marin et al. 1996). However, haloperidol was the only typical antipsychotic used in this study. Therefore, future experiments testing several typical antipsychotics are necessary to further discriminate between typical and atypical antipsychotics and to characterize their effects on gene expression. For zotepine, this is the first report

to show that this compound does not induce neuronal enkephalin mRNA expression or increased enkephalin tissue levels in the caudate putamen. The present findings therefore reaffirm the suggestion (Augood et al. 1993; Mijster et al. 1998) that enkephalin gene expression in the caudate-putamen is either a marker for or plays a role in the development of extrapyramidal side-


a) *in situ* hybridization**b) radioimmunoassay**

Figure 3. A schematic drawing according to the Atlas of Paxinos: (Paxinos and Watson 1986) indicates the region of prefrontal cortex analyzed by *in situ* hybridization (a) and radioimmunoassay (b). Four sections were sampled for *in situ* hybridization with a distance of about 400 μ m between each section. The schematic drawing represents the level of the third section. Scale bar represents 1mm.

effects associated with typical antipsychotic drugs. The lack of action of zotepine and clozapine on this parameter in the present study concurs with the atypical profile of these drugs both in animal models and in the clinic.

In the prefrontal cortex, antipsychotic drugs showed a distinct pattern on gene expression in two regards. First, haloperidol did not lead to any significant changes in synaptic proteins levels and in their encoding mRNAs. Secondly, in contrast to secretogranin II which was not affected by any antipsychotic drug, there was a change in tissue levels and mRNA expression of chromogranin A and B. In general, there was a good correspondence between decreased tissue levels and decreased expression of respective mRNAs. This indicates that decreased tissue levels are caused by a decreased expression of respective mRNAs in prefrontal cortex neurons rather than by an increased release of chromogranin A or chromogranin B.

The distinct effects of classical and novel antipsychotic drugs on synaptic proteins are in line with a differential induction of immediate early genes in the prefrontal cortex (Guo et al. 1998). Novel antipsychotic drugs, such as clozapine and olanzapine, caused an induction of these genes in acute experiments (Robertson et al. 1994; Deutch and Duman 1996; Robertson and Fibiger 1996). Repeated administration of typical versus atypical antipsychotic drugs leads to a region specific induction of deltaFosB (Atkins et al. 1999). In caudate putamen, typical antipsychotics cause a greater induction of deltaFosB than atypical antipsychotics (Hiroi and Graybiel 1996, Atkins et al. 1999).

What might be the reasons for changes in synaptic proteins? Clozapine, risperidone, and zotepine are more potent serotonin 2A receptor antagonists than dopamine D2-like receptor antagonists. Haloperidol lacks 5-serotonin 2A properties. Therefore, the changes

can not only be driven by dopamine effects. In contrast to haloperidol, clozapine as well as risperidone increase extracellular serotonin levels in medial prefrontal cortex (Ichikawa et al. 1998) which could influence the biosynthesis of chromogranins (Kroesen et al. 1995). What might be the consequences of changed chromogranin levels? Chromogranins are a family of regulatory secretory proteins. Their aggregation is promoted by calcium ions and low pH, two parameters in the trans-Golgi network. There is now increasing evidence that chromogranins play an important role for sorting to secretory granules (Thiele and Huttner 1998). In the case of chromogranin B, reduction of the disulfide bond is sufficient to cause complete missorting of constitutive secretory vesicles (Chanat et al. 1993). Therefore, our experiments showing changes in chromogranin tissue levels may be of importance for the sorting of vesicles. Another important aspect is the capability of chromogranin deriving peptides to influence release of other vesicle constituents. It was recently shown that chromogranin A desensitizes catecholamine release (Mahata et al. 1999).

Alterations in synaptic proteins and their encoding mRNAs are reported in several brain areas including the prefrontal cortex of schizophrenic patients in the absence of gross structural abnormalities (Selemon and Goldman-Rakic 1999). The emerging picture of neuropathology in schizophrenia is one of subtle changes in cellular architecture and brain circuitry (Lieberman et al. 1997) that nonetheless may have a devastating impact on cortical function. Antipsychotic drugs cause postsynaptic adaptations including regulation of neuronal gene expression leading to changes in synaptic protein content. Whether these results are of therapeutic relevance may depend on the neuronal system targeted by a particular drug (Hyman and Nestler 1996).

CONCLUSIONS

The present study shows that increases in enkephalin and its mRNA in the caudate-putamen can differentiate between typical and atypical antipsychotic drugs since they are not observed with atypical antipsychotic drugs. Conversely, similar increases in limbic regions occur irrespective of the side-effect profile of these antipsychotics and so may be linked to the therapeutic actions of these drugs. There is also evidence that chromogranins are sensitive and suitable markers to detect biochemical changes caused by chronic antipsychotic drugs. Expression patterns of some vesicle markers are distinctly altered depending on treatment with haloperidol or novel antipsychotic drugs.

ACKNOWLEDGMENTS

This work was supported by BASF Pharma/Knoll Pharmaceuticals Austrian Science Foundation (SFB F00206) and the Austrian National Bank (P 7150). The excellent technical assistance of I. Berger is gratefully acknowledged.

REFERENCES

- Angulo JA (1992): Involvement of dopamine D1 and D2 receptors in the regulation of proenkephalin mRNA abundance in the striatum and accumbens of the rat brain. *J Neurochem* 58:110–1109
- Arnt J, Skarsfeldt T (1998): Do novel antipsychotics have similar pharmacological characteristics? A review of the evidence. *Neuropsychopharmacology* 18:63–101
- Atkins JB, Chlan-Fourney J, Nye HE, Hiroi N, Carlezon WA, Nestler EJ (1999): Region-specific induction of deltaFosB by repeated administration of typical versus atypical antipsychotic drugs. *Synapse* 33:118–128
- Augood SJ, Westmore K, Faull RL, Emson PC (1993): Neuroleptics and striatal neuropeptide gene expression. *Prog Brain Res* 99:181–199
- Chanat E, Weiss U, Huttner WB, Tooze SA (1993): Reduction of the disulfide bond of chromogranin B (secretogranin I) in the trans-Golgi network causes its missorting to the constitutive secretory pathways. *EMBO J* 12:2159–2168
- Comb M, Seeburg PH, Adelman J, Eiden L, Herbert E (1982): Primary structure of the human met- and leu-enkephalin precursor of its mRNA. *Nature* 295:663–666
- Deutch AY, Duman RS (1996): The effects of antipsychotic drugs on Fos protein expression in the prefrontal cortex: Cellular localization and pharmacological characterization. *Neuroscience* 70:377–389
- Ellenbroek B, Cools AR (1988): The Paw Test: An animal model for neuroleptic drugs which fulfils the criteria for pharmacological isomorphism. *Life Sci* 42:1205–1213
- Ellenbroek BA, Prinssen EP, Cools AR (1994): The role of serotonin receptor subtypes in the behavioural effects of neuroleptic drugs. A paw test study in rats. *Eur J Neurosci* 6:1–8
- Ezrin WC, Seeman P (1977): Tolerance of haloperidol catalepsy. *Eur J Pharmacol* 41:321–327
- Fleischhacker WW, Hummer M (1997): Drug treatment of schizophrenia in the 1990s. Achievements and future possibilities in optimising outcomes. *Drugs* 53:915–929
- Forss PS, Danielson P, Battenberg E, Bloom F, Sutcliffe JG (1989): Nucleotide sequence and cellular distribution of rat chromogranin B (secretogranin I) mRNA in the neuroendocrine system. *J Mol Neurosci* 1:63–75
- Gerdes HH, Phillips E, Huttner WB (1988): The primary structure of rat secretogranin II deduced from a cDNA sequence. *Nucl Ac Res* 16:11811
- Guo N, Vincent SR, Fibiger HC (1998): Phenotypic characterization of neuroleptic-sensitive neurons in the fore-

- brain: Contrasting targets of haloperidol and clozapine. *Neuropsychopharmacology* 19:133–145
- Herman ZS, Huzarska M, Kmiecik KK, Kowalski J (1991): Chronic treatment with chlorpromazine, thioridazine or haloperidol increases striatal enkephalins and their release from rat brain. *Psychopharmacology (Berl)* 104:106–112
- Hiroi N, Graybiel AM (1996): Atypical and typical neuroleptic treatments induce distinct programs of transcription factor expression in the striatum. *J Comp Neurol* 374:70–83
- Huttner WB, Gerdes HH, Rosa P (1991): The granin (chromogranin/secretogranin) family. *TIBS* 16:27–30
- Hyman SE, Nestler EJ (1996): Initiation and adaptation: A paradigm for understanding psychotropic drug action. *Am J Psychiatry* 153:151–162
- Iacangelo A, Okayama H, Eiden LE (1988): Primary structure of rat chromogranin A and distribution of its mRNA. *FEBS Lett* 227:115–121
- Ichikawa J, Kuroki T, Dai J, Meltzer HY (1998): Effect of antipsychotic drugs on extracellular serotonin levels in rat medial prefrontal cortex and nucleus accumbens. *Eur J Pharmacol* 351:163–171
- Kirschmair R, Hogue-Angeletti R, Gutierrez J, Fischer-Colbrie R, Winkler H (1993): Secretoneurin—a neuropeptide generated in brain, adrenal medulla and other endocrine tissues by proteolytic processing of secretogranin II (chromogranin C). *Neuroscience* 53:359–365
- Knable MB, Weinberger DR (1997): Dopamine, the prefrontal cortex and schizophrenia. *J Psychopharmacol* 11:123–131
- Kroesen S, Marksteiner J, Leitner B, Hogue Angeletti R, Fischer-Colbrie R, Winkler H (1996): Rat brain: distribution of immunoreactivity of PE-11, a peptide derived from chromogranin B. *Eur J Neurosci* 8:2679–2689
- Kroesen S, Marksteiner J, Mahata SK, Mahata M, Fischer-Colbrie R, Saria A, Kapeller I, Winkler H (1995): Effects of haloperidol, clozapine and citalopram on messenger RNA levels of chromogranins A and B and secretogranin II in various regions of rat brain. *Neuroscience* 69:881–891
- Le Moine C, Normand E, Bloch B (1991): Phenotypical characterization of the rat striatal neurons expressing the D1 dopamine receptor gene. *Proc Natl Acad Sci USA* 88:4205–4209
- Leysen JE, Janssen PM, Schotte A, Luyten WH, Megens AA (1993): Interaction of antipsychotic drugs with neurotransmitter receptor sites in vitro and in vivo in relation to pharmacological and clinical effects: Role of 5HT₂ receptors. *Psychopharmacology (Berl)* 112:S40–S54
- Lieberman JA, Sheitman BB, Kinon BJ (1997): Neurochemical sensitization in the pathophysiology of schizophrenia: Deficits and dysfunction in neuronal regulation and plasticity. *Neuropsychopharmacology* 17:205–229
- MacArthur L, Eiden L (1996): Neuropeptide genes: targets of activity-dependent signal transduction. *Peptides* 17:721–728
- Mahata SK, Mahata M, Marksteiner J, Sperk G, Fischer-Colbrie R, Winkler H (1991): Distribution of mRNAs for chromogranin A and B and secretogranin II in rat brain. *Eur J Neurosci* 3:895–904
- Mahata SK, Mahata M, Parmer RJ, O'Connor DT (1999): Desensitization of catecholamine release. The novel catecholamine release-inhibitory peptide catestatin (chromogranin a344-364) acts at the receptor to prevent nicotinic cholinergic tolerance. *J Biol Chem* 274:2920–2928
- Marin C, Engber TM, Bonastre M, Chase TN, Tolosa E (1996): Effect of long-term haloperidol treatment on striatal neuropeptides: Relation to stereotyped behavior. *Brain Res* 731:57–62
- Marksteiner J, Kirschmair R, Mahata SK, Mahata M, Fischer-Colbrie R, Hogue-Angeletti R, Saria A, Winkler H (1993a): Distribution of secretoneurin, a peptide derived from secretogranin II, in rat brain: An immunocytochemical and radioimmunological study. *Neuroscience* 54:923–944
- Marksteiner J, Saria A, Miller CH, Krause JE (1993b): Increased synthesis of neurokinin B and enkephalin after chronic haloperidol treatment. *Regul Pept* 46:349–351
- Marksteiner J, Sperk G (1988): Concomitant increase of somatostatin, neuropeptide Y and glutamate decarboxylase in the frontal cortex of rats with decreased seizure threshold. *Neuroscience* 26:379–385
- Mercugliano M, Chesselet MF (1992): Clozapine decreases enkephalin mRNA in rat striatum. *Neurosci Lett* 136:10–14
- Mijnster MJ, Ingham CA, Meredith GE, Docter GJ, Arbuthnott GW (1996): Morphological changes in met(5)-enkephalin-immunoreactive synaptic boutons in the rat neostriatum after haloperidol decanoate treatment. *Eur J Neurosci* 8:716–726
- Mijnster MJ, Schotte A, Docter GJ, Voorn P (1998): Effects of risperidone and haloperidol on tachykinin and opioid precursor peptide mRNA levels in the caudate-putamen and nucleus accumbens of the rat. *Synapse* 28:302–312
- Needham PL, Atkinson J, Skill MJ, Heal DJ (1996): Zotepine: Preclinical tests predict antipsychotic efficacy and an atypical profile. *Psychopharmacol Bull* 32:123–128
- Nguyen TV, Kosofsky BE, Birnbaum R, Cohen BM, Hyman SE (1992): Differential expression of c-Fos and Zif268 in rat striatum after haloperidol, clozapine, and amphetamine. *Proc Natl Acad Sci USA* 89:4270–4274
- Normand E, Popovici T, Fellmann D, Bloch B (1987): Anatomical study of enkephalin gene expression in the rat forebrain following haloperidol treatment. *Neurosci Lett* 83:232–236
- O'Donnell P, Grace AA (1998): Dysfunctions in multiple inter-related systems as the neurobiological bases of schizophrenic symptom clusters. *Schizophr Bull* 24:267–283
- Paxinos G, Watson C (1986): The rat brain in stereotaxic coordinates, 2nd ed. Sydney, Academic Press
- Robertson GS, Fibiger HC (1992): Neuroleptics increase c-fos expression in the forebrain: Contrasting effects of haloperidol and clozapine. *Neuroscience* 46:315–328
- Robertson GS, Fibiger HC (1996): Effects of olanzapine on regional C-Fos expression in rat forebrain. *Neuropsychopharmacology* 14:105–110
- Robertson GS, Matsumura H, Fibiger HC (1994): Induction patterns of Fos-like immunoreactivity in the forebrain

- as predictors of atypical antipsychotic activity. *J Pharmacol Exp Ther* 271:1058–1066
- Romano GJ, Shivers BD, Harlan RE, Howells RD, Pfaff DW (1987): Haloperidol increases proenkephalin mRNA levels in the caudate-putamen of the rat: A quantitative study at the cellular level using in situ hybridization. *Brain Res* 388:33–41
- Schotte A, Janssen PF, Gommeren W, Luyten WH, Van GP, Lesage AS, De LK, Leysen JE (1996): Risperidone compared with new and reference antipsychotic drugs: In vitro and in vivo receptor binding. *Psychopharmacology (Berl)* 124:57–73
- Selemon LD, Goldman-Rakic PS (1999): The reduced neuro-pil hypothesis: A circuit based model of schizophrenia. *Biol Psychiatry* 45:17–25
- Shen PJ, Gundlach AL (1996): Chromogranin mRNA levels in the brain as a marker for acute and chronic changes in neuronal activity: Effect of treatments including seizures, osmotic stimulation and axotomy in the rat. *Eur J Neurosci* 8:988–1000
- Thiele C, Huttner WB (1998): The disulfide-bonded loop of chromogranins, which is essential for sorting to secretory granules, mediates homodimerization. *J Biol Chem* 273:1223–1231