
BRIEF REPORT

Δ^9 -Tetrahydrocannabinol and Synthetic Cannabinoids Prevent Emesis Produced by the Cannabinoid CB₁ Receptor Antagonist/Inverse Agonist SR 141716A

Nissar A. Darmani, Ph.D.

*There is substantial clinical evidence that Δ^9 -tetrahydrocannabinol (Δ^9 -THC) and its synthetic analogs (nabilone and levonantradol) can prevent emesis in cancer patients receiving chemotherapy. Limited available animal studies also support the antiemetic potential of these cannabinoids. The present study investigates the mechanism of antiemetic action of cannabinoids in an established animal model of emesis, the least shrew (*Cryptotis parva*). Since cannabinoid agonists prevent emesis, it was hypothesized that blockade of either the cannabinoid CB₁ receptor or the cannabinoid CB₂ receptor would induce vomiting. Thus, the emetic potential of SR 141716A (CB₁ receptor antagonist) or SR 144528 (CB₂ receptor antagonist) was investigated. Both intraperitoneal (0, 1, 2.5, 5, 10 and 20 mg/kg, n = 7–15 per group) and subcutaneous (0, 10, 20 and 40 mg/kg, n = 6–9 per group) administration of SR 141716A caused emesis (ED₅₀ = 5.52 \pm 1.23 and 20.2 \pm 1.02 mg/kg, respectively) in the least shrew in a dose-dependent manner. Indeed, both the frequency of emesis and the percentage of animals vomiting increased with increasing doses of SR 141716A. Significant effects were seen at the 10- and 20-mg/kg doses for the IP route, while only the 40-mg/kg dose produced significant emesis via the SC route. The CB₂ antagonist failed to*

produce emesis via either route of administration. SR 141716A at an IP dose of 20 mg/kg was used to induce emesis for drug interaction studies. Thus, varying doses of three different classes of cannabinoid agonists [CP 55, 940 (0, 0.1, 0.5 and 1 mg/kg), WIN 55, 212-2 (0, 1, 5 and 10 mg/kg), and Δ^9 -THC (0, 5, 10 and 20 mg/kg)], were administered IP to different groups of shrews 10 min prior to SR 141716A injection. The frequency of emesis was recorded for 30 min following the administration of SR 141716A. The order of potency for reducing both the frequency of emesis and the percentage of shrews vomiting was CP 55, 940 > WIN 55, 212-2 > Δ^9 -THC which is consistent with an action on the CB₁ receptor. These results suggest that the antiemetic activity of Δ^9 -THC and its synthetic analogs reside in their ability to stimulate the cannabinoid CB₁ receptor. Furthermore, the antiemetic potency of CP 55, 940 is 45 times greater than Δ^9 -THC. On the other hand, blockade of CB₁ receptors can induce vomiting, which implicates an important role for endogenous cannabinoids in emetic circuits.

[Neuropsychopharmacology 24:198–203, 2001]

© 2000 American College of Neuropsychopharmacology.
Published by Elsevier Science Inc.

From the Department of Pharmacology, Kirksville College of Osteopathic Medicine, Kirksville, Missouri, USA

Address correspondence to: Dr. Nissar A. Darmani, Department of Pharmacology, Kirksville College of Osteopathic Medicine, 800

West Jefferson Street, Kirksville, MO 63501, USA. Tel.: (660) 626-2326; Fax: (660) 626-2728.

E-mail address: ndarmani@kcom.edu

Received April 17, 2000; revised July 14, 2000; accepted August 21, 2000.

KEY WORDS: *Marijuana*; *Delta-9-tetrahydrocannabinol*; *CP 55, 940*; *WIN 55, 212-2*; *SR 141716A*; *SR 144528*; *Emesis*; *CB₁ receptor*; *CB₂ receptor*

In the early 1970s, accumulating anecdotal reports by young cancer patients suggested that smoking marijuana would alleviate the nausea and vomiting caused by chemotherapeutic agents. Since then, both government- and industry-sponsored clinical trials were initiated to test the antiemetic potential of Δ^9 -tetrahydrocannabinol (Δ^9 -THC) and some of its synthetic analogs such as nabilone and levonantradol (Gralla 1999; Voth and Schwartz 1997). This literature suggests that both Δ^9 -THC and its tested analogs are useful antiemetics in some patients for the prevention of nausea and vomiting associated with cancer chemotherapy. Though cannabinoids appear to be a more efficacious class of antiemetics than dopamine D₂ receptor antagonists for the prevention of chemotherapy-induced vomiting, the efficacy of tested cannabinoids to date seems not to be as high as the more potent antiemetics such as the selective 5-HT₃ receptor antagonists (Gralla 1999). However, one interesting advantage of cannabinoids is that many of the patients who are protected from the acute phase of emesis, also respond well during the delayed phase of chemotherapy-induced emesis which 5-HT₃ receptor antagonists poorly control (Abrahamov et al. 1995; Chan et al. 1987; Dalzell et al. 1986).

Unlike the relatively large body of clinical reports, only a few published animal studies on the antiemetic effects of cannabinoids are available. Several cannabinoids can block cisplatin- or apomorphine-induced emesis in a variety of animal species including the cat, the pigeon and the least shrew (*Cryptotis parva*) (McCarthy and Borison 1981; McCarthy et al. 1984; London et al. 1979; Stark 1982; Darmani 2000). Until recently, an animal model of emesis had not been employed for investigating the cannabinoid receptor involved in the antiemetic effect of Δ^9 -THC and other cannabinoids. In the present study, the least shrew (*Cryptotis parva*) has been used as an animal model of emesis. This species was recently introduced as a new serotonergic (Darmani 1998) and dopaminergic (Darmani et al. 1999) experimental model of vomiting. The least shrew is a small insectivore (adult weight 4–6 g) that lives in various ecological niches in Central and North America. The family Soricidae, to which shrews belong, constitute over 266 species (Churchfield 1990). Over the last decade, Japanese investigators have established the house musk shrew (*Suncus murinus*) as an experimental model for the various emetic stimuli (Matsuki et al. 1988; Torii et al. 1991). *Suncus murinus* is relatively a larger animal (adult being 50–100 g in weight) and is endogenous to Asia and Africa.

While the antiemetic effects of Δ^9 -THC appear to be receptor-mediated, it is unclear whether the cannabinoid CB₁ and/or CB₂ receptors are involved in emesis. In-

volvement of CB₁ receptor appears most likely since Δ^9 -THC produces most of its effects via this site (Pertwee 1997). If activation of cannabinoid receptors prevent emesis, then blockade of these receptors may produce vomiting. Thus, the present study investigated: (1) whether administration of the selective cannabinoid CB₁-receptor antagonist SR 141716A (Rinaldi-Carmona et al. 1994), or the CB₂-receptor antagonist SR 144528 (Rinaldi-Carmona et al. 1998), can induce emesis in the least shrew (Darmani 1998; Darmani et al. 1999); and (2) whether the induced emesis can be blocked by the cannabinoid agonists Δ^9 -THC and its newly introduced synthetic analogs CP 55, 940 and WIN 55, 212-2 (Pertwee 1997).

MATERIALS AND METHODS

Animals and Drugs

Shrews (*Cryptotis parva*) were bred and maintained in the animal facilities of the Kirksville College of Osteopathic Medicine, Kirksville, Missouri. Both male and female shrews (4–6 g, 45–70 days old) were used throughout the study. The animals were kept on a 14:10-h light-dark cycle at a humidity controlled room temperature of $21 \pm 1^\circ\text{C}$ with ad lib supply of food and water. The feeding and maintenance of shrews are fully described elsewhere (Darmani 1998; Darmani et al. 1999). The following drugs were purchased from Research Biochemicals Inc., Natick, MA: Δ^9 -tetrahydrocannabinol (Δ^9 -THC) and R(+)-WIN 55, 212-2 mesylate. CP 55, 940 was obtained from Pfizer (Groton, CT). SR 141716A and SR 144528 were generously donated by Professor B.R. Martin. All drugs were dissolved in a 1:1:18 solution of ethanol: emulphor: 0.9% saline to twice the stated drug concentrations. These drugs concentrations were further diluted by the addition of an equal volume of saline. This procedure was necessary, because the 1:1:18 vehicle mixture can cause emesis in up to 20% of animals by itself. The final vehicle mixture induced emesis only very rarely. All drugs were administered at a volume of 0.1 ml/10g of body weight. All animals received care according to the "Guide for the Care and Use of Laboratory Animals," DHSS Publication, revised, 1985.

Experimental Protocols

The present protocols were based upon our previous emesis studies in the least shrew (Darmani 1998; Darmani et al. 1999). On the test day, the shrews were transferred to the experimental room and were allowed to acclimate for at least 1 h prior to experimentation. To habituate the shrews to the test environment, each animal was randomly selected and transferred to a $20 \times 18 \times 21$ cm clean clear plastic cage and offered 4 meal worms (*Tenebrio* sp) 30 min prior to experimentation. Different groups of shrews were then injected either in-

traperitoneally or subcutaneously with vehicle ($n = 11$ – 12) or varying doses of the CB₁ antagonist SR 141716A (1, 2.5, 5, 10 and 20 mg/kg, $n = 7$ – 15), or the CB₂ antagonist SR 144528 (10, 20 and 40 mg/kg, $n = 8$ – 11 per group). Immediately following injection, each shrew was placed in the observation cage and the onset latency to first vomit as well as the frequency of vomiting (mean \pm SEM) were recorded for each individual shrew for the next 60 min. These data showed that intraperitoneal administration of 20 mg/kg of SR 141716A produced a robust frequency of emesis. This dose and route of administration of SR 141716A was chosen for subsequent studies in which the antiemetic effects of Δ^9 -THC as well as its synthetic analogs CP 55, 940 and WIN 55, 212-2 were investigated.

For these interaction studies, different doses of either CP 55, 940 (0, 0.1, 0.5 and 1 mg/kg, $n = 6$ – 8 per group), WIN 55, 212-2 (0, 1, 5 and 10 mg/kg, $n = 6$ – 9 per group), or Δ^9 -THC (0, 5, 10 and 20 mg/kg, $n = 7$ – 9 per group) were administered intraperitoneally to different groups of shrews 10 min prior to SR 141716A (20 mg/kg, IP) injection. Emesis was recorded for 30 min immediately following SR 141716A administration as described above.

Statistical Analysis

The data were analyzed by the Kruskal-Wallis nonparametric one-way analysis of variance (ANOVA) and posthoc analysis by Dunn's multiple comparisons test. A p -value of $<.05$ was necessary to achieve statistical significance. The ED₅₀ (the effective dose that produced emesis in 50% of animals) and ID₅₀ (the inhibitory dose that prevented emesis in 50% of shrews) were calculated by the use of a computerized program (Graph Pad InPlot, San Diego, CA).

RESULTS

The Kruskal-Wallis nonparametric ANOVA test indicated that intraperitoneal administration of SR 141716A dose-dependently increased the percentage of shrews vomiting (ED₅₀ = 5.52 ± 1.23) ($K_{W_{58,5}} = 28.1$, $p < .0001$) (Table 1). Dunn's multiple comparisons test showed that relative to the vehicle-injected control group, significant enhancements in the number of animals exhibiting emesis occurred in the groups injected with the

Table 1. Dose-dependent Emetogenic Effects of Different Routes of Administration of the Cannabinoid CB₁ (SR 141716A) and CB₂ (SR144528) Antagonists

| Drugs | Number of animals vomiting/tested | Vomiting frequency | Latency to first vomit (minutes) |
|-----------------|-----------------------------------|-----------------------|----------------------------------|
| SR 141716A | | | |
| Intraperitoneal | | | |
| 0 | 1/12 | 0.08 ± 0.08 | 19.20 |
| 1 | 2/10 | 0.5 ± 0.3 | 21.24 |
| 2.5 | 0/8 | 0 ± 0 | — |
| 5 | 8/15 | 0.93 ± 0.3 | 13.0 |
| 10 | 9/12* | $1.7 \pm 0.5^*$ | 8.78 |
| 20 | 7/7*** | $05.7 \pm 0.52^{***}$ | 5.28 |
| Subcutaneous | | | |
| 0 | 0/9 | 0 | — |
| 10 | 0/6 | 0 | — |
| 20 | 2/7 | 2 ± 1.3 | 10.32 |
| 40 | 4/7* | $2.9 \pm 1.5^*$ | 9.40 |
| SR 144528 | | | |
| Intraperitoneal | | | |
| 0 | 1/11 | 0.09 ± 0.09 | 10.21 |
| 10 | 0/8 | 0 ± 0 | — |
| 20 | 2/10 | 0.2 ± 0.13 | 12.28 |
| 40 | 2/10 | 0.3 ± 0.2 | 8.39 |
| Subcutaneous | | | |
| 0 | 0/9 | 0 ± 0 | — |
| 10 | 1/7 | 0.14 ± 0.14 | 3.49 |
| 20 | 0/7 | 0 ± 0 | — |
| 40 | 0/7 | 0 ± 0 | — |

Values for vomiting frequency and latency for first vomit are mean of those animals which exhibited vomiting. Animals which failed to vomit were not included in these mean values.

The above parameters were recorded for 60 min following either the intraperitoneal or subcutaneous administration of the cited selective cannabinoid antagonists.

* $p < .05$.

** $p < .01$.

*** $p < .001$ indicate significant differences relative to vehicle control by Dunn's multiple comparisons test.

10 ($p < .05$) and 20 mg/kg ($p < .001$) doses of SR 141716A. Moreover, the larger doses of SR 141716A induced emesis more rapidly. Indeed, depending upon the dose of SR 141716A administered, the mean latency to onset of first vomit in responsive animals varied from 5.28 to 19.20 minutes. In addition, the frequency of SR 141716A induced emesis also increased in a dose-dependent manner ($Kw_{58.5} = 34.9$, $p < .0001$) (Table 1). Again, significant enhancements in the number of vomiting episodes occurred at the 10 ($p < .05$) and 20 mg/kg ($p < .001$) doses. When administered subcutaneously, SR 141716A appeared to be a less efficacious emetogen (Table 1). Indeed, the Kruskal-Wallis ANOVA test indicated that although SR 141716A can increase both the frequency ($Kw_{25.3} = 9.5$, $p < .003$), and the percentage of animals vomiting ($ED_{50} = 20.2 \pm 1.02$ mg/kg) ($Kw_{25.3} = 9.5$, $p < .02$), a significant effect ($p < .05$) was only observed at the 40-mg/kg dose by the Dunn's multiple comparisons posthoc test (Table 1). The CB₂ antagonist, SR 144528, over the dose range of 10–40 mg/kg, IP and SC, failed to produce a significant degree of emesis in the least shrew (Table 1).

Table 2 shows the ability of three cannabinoids (CP 55, 940; WIN 55, 212-2 and Δ^9 -THC) in preventing emesis induced by the cannabinoid CB₁ receptor antagonist/inverse agonist SR 141716A. The cited doses of CP 55, 940 (0.1–1 mg/kg) attenuated both the percentage of animals vomiting ($ID_{50} = 0.35 \pm 2.2$ mg/kg) ($Kw_{25.3} = 19.8$, $p < .0002$) and the frequency of vomitings in a dose-dependent manner (Table 2). However, a significant decrease in both the percentage of shrews vomiting ($p < .01$) and the frequency of vomitings ($p < .01$) was seen at the 1 mg/kg dose of CP 55, 940. In a similar manner, intraperitoneal administration of WIN 55, 212-2 (1–10 mg/kg), dose-dependently antagonized the abil-

ity of SR 141716A to induce emesis in different shrews ($ID_{50} = 3.95 \pm 1.2$) ($Kw_{28.3} = 10.8$, $p < .01$) as well as reducing the frequency of the induced emesis ($Kw_{28.3} = 10.6$, $p < .01$) (Table 2). However, significant reductions ($p < .05$) were only observed for both emetic parameters at the 10 mg/kg dose. The Kruskal-Wallis nonparametric ANOVA test also showed that intraperitoneal injection of Δ^9 -THC can reduce the percentage of shrews vomiting in response to SR 141716A administration ($ID_{50} = 15.2 \pm 3.2$ mg/kg) ($Kw_{29.3} = 22.6$, $p < .0001$), as well as attenuating the frequency of the induced emesis ($Kw_{29.3} = 20.04$, $p < .0002$). However significant reductions in emetic parameters ($p < .001$) were only seen at the 20-mg/kg dose of Δ^9 -THC (Table 2).

DISCUSSION

The least shrew appears to be an excellent new animal model of emesis (Darmani 1998; Darmani et al. 1999). Both dopamine D₂- and serotonin 5-HT₃-receptor agonists induce vomiting in a potent manner in this species. Furthermore, 5-HT₃ receptor antagonists block cisplatin-induced emesis in these animals (Darmani 1998). There is overwhelming clinical evidence that Δ^9 -THC and some of its synthetic analogs (nabilone and levonantradol) can prevent emesis in some cancer patients receiving chemotherapy (Gralla 1999; Voth and Schwartz 1997). Likewise, as described in the introduction, some cannabinoids prevent cisplatin- or apomorphine-induced emesis in a number of different species including the least shrew.

Until the present study the mechanism of antiemetic action of cannabinoids was not known. Therapeutically effective antiemetics antagonize the emetic action of

Table 2. Dose-dependent Antiemetic Effects of Intraperitoneally Administered Cannabinoids Against the CB₁ Receptor Antagonist (SR 141716A)-induced Vomiting

| Drug | Dose (mg/kg) | Number of animals vomited/tested | Vomiting frequency |
|-----------------|--------------|----------------------------------|--------------------|
| Control | 0 | 8/8 | 2.9 \pm 0.2 |
| CP 55, 940 | 0.1 | 6/6 | 3 \pm 0.4 |
| | 0.5 | 3/7 | 1.1 \pm 0.5 |
| | 1 | 0/7*** | 0 \pm 0** |
| WIN 55, 212-2 | 1 | 5/6 | 2.5 \pm 1 |
| | 5 | 3/8 | 1 \pm 0.5 |
| | 10 | 3/9* | 0.9 \pm 0.5* |
| Δ^9 -THC | 5 | 6/7 | 3.1 \pm 0.7 |
| | 10 | 8/8 | 2 \pm 0.3 |
| | 20 | 1/9*** | 0.1 \pm 0.1*** |

Different shrews received intraperitoneally either vehicle or the cited doses of CP 55, 940; WIN 55, 212-2 or Δ^9 -THC 10 min prior to administration of SR 141716A (20 mg/kg, IP). The emetic parameters were recorded for the next 30 min immediately following the SR 141716A injection.

* $p < .05$.

** $p < .01$.

*** $p < .001$ indicate significant differences relative to vehicle control by Dunn's multiple comparisons test.

endogenous neurotransmitters such as acetylcholine, dopamine, histamine, serotonin or substance P (Naylor and Rudd 1996). Unlike these receptor antagonists, cannabinoids decrease emesis as a result of agonist action at cannabinoid CB₁ and/or CB₂ receptors (Pertwee 1997). Thus, it seemed reasonable that antagonism of one or both cannabinoid receptors should induce emesis. This proposal appears to be true since the intraperitoneal or subcutaneous administration of the selective CB₁ antagonist/inverse agonist, SR 141716A, caused emesis in a dose-dependent manner in the least shrew ($ED_{50} = 5.52 \pm 1.25$ mg/kg). Indeed, both the percentage of animals vomiting and the frequency of induced emesis were significantly increased in response to administration of increasing doses of SR 141617A. Intraperitoneal injection of SR 141716A caused profound emesis at 10 mg/kg or greater doses. However, SR 141716A is a less efficacious emetic ($ED_{50} = 20.2 \pm 1.02$ mg/kg) when administered via the subcutaneous route since only the largest dose employed (40 mg/kg) caused significant emesis. At 10 mg/kg or larger doses, SR 141716A has also been found to increase the spontaneous locomotor activity in mice following intravenous administration (Compton et al. 1996). Peripheral administration of such doses of SR 141716A also produces other motor behaviors such as the head-twitch response, lateral scratchings or wet-dog shakes in drug-naïve mice and rats (Aceto et al. 1998; Cook et al. 1998; Darmani and Pandya 2000; Rubino et al. 1998). The present and the cited studies suggest that SR 141716A produces these effects either by antagonizing the action of an endogenous cannabinoid(s), or by exerting an inverse agonist action. The selective cannabinoid CB₂ antagonist SR 144528 failed to produce significant emesis in the least shrew either via the intraperitoneal or the subcutaneous routes. Thus, the present results suggest that cannabinoid agonists may produce their antiemetic action via the activation of CB₁ receptors.

To show that SR 141716A-induced emesis is a cannabinoid receptor-mediated event, the antiemetic effect of three structurally different classes of cannabinoid receptor agonists (CP 55, 940; WIN 55, 212-2 and Δ^9 -THC) were investigated. The cited cannabinoids blocked the ability of SR 141716A to induce emesis in a dose-dependent manner with the following antiemetic ID_{50} rank order: CP 55, 940 < WIN 55, 212-2 < Δ^9 -THC. This antiemetic potency rank order in the least shrew, mirrors both the affinity of these cannabinoids for CB₁ and CB₂ receptors (Matsuda 1997; Pertwee 1997) and their ED_{50} potency values for the CB₁ receptor-mediated tetrad of behaviors in mice (Abood and Martin 1992). These results further support the suggestion that cannabinoids exert their antiemetic action via CB₁ receptors. However, the nonpsychoactive (HU-211) enantiomer of the potent cannabinoid HU-210, which lacks affinity for CB₁ receptors, has also been reported to prevent cis-

platin-induced emesis (Feigenbaum et al. 1989). Since HU-211 stereoselectively blocks the glutamate NMDA receptor, its antiemetic property is probably related to antagonism of this excitatory receptor. Indeed, NMDA antagonists can prevent emesis produced by a variety of emetic stimuli (Lehmann and Karrberg 1996; Lucot 1998).

Apart from the present study, very little is known regarding the antiemetic structure activity relationship of different cannabinoids in any species. Furthermore, as yet no clinical study has demonstrated a better margin of safety for one cannabinoid antiemetic versus another. Even though there is no information regarding the therapeutic index of CP 55, 940 in man, the present antiemetic potency order in the least shrew indicates that this cannabinoid possesses a superior antiemetic property than Δ^9 -THC. Indeed, relative to Δ^9 -THC, the index of psychoactivity of CP 55, 940 (i.e., its ED_{50} in producing the tetrad of behaviors) only varies from 4–25 times in mice (Abood and Martin 1992); whereas the antiemetic potency of this cannabinoid is 45 times greater than Δ^9 -THC in the least shrew. In the case of SR 141716A-induced emesis, the most potent tested cannabinoid in the present study, CP 55, 940, also caused the greatest degree of antiemetic activity. Indirect evidence from nonemetic species also support the involvement of CB₁ receptors in emesis. Thus, different classes of cannabinoid agonists (anandamide, methanandamide, nabilone, Δ^9 -THC and WIN 55, 212-2) reduce gastrointestinal motility and intestinal transit in rodents via SR 141716A-sensitive CB₁ receptors (Calignano et al. 1997; Colombo et al. 1998; Izzo et al. 1999; Krowicki et al. 1999; Shook and Burks 1989). Furthermore, these studies have shown that the emetic doses of SR 141716A employed in the present study, also promotes defecation and GI motility in rodents.

In summary, the present and published studies suggest that the antiemetic activity of cannabinoid agonists is mediated by the CB₁ receptor and endogenous cannabinoid neurotransmitter system may play an important regulatory role in emesis.

ACKNOWLEDGMENTS

This work was supported by a grant from the National Institute on Drug Abuse (DA 0767). The author would like to thank R. Chronister for typing the manuscript.

REFERENCES

- Abood ME, Martin BR (1992): Neurobiology of marijuana abuse. *Trends Pharmacol Sci* 13:201–206
- Abrahamov AS, Abrahamov A, Mechoulam R (1995): An efficient new cannabinoid antiemetic in pediatric oncology. *Life Sci* 56:2097–2102

- Aceto MD, Scates SM, Razdan RJ, Martin BR (1998): Anandamide, an endogenous cannabinoid, has very low physical dependence potential. *J Pharmacol Exp Therap* 287:598–605
- Calignano A, La Rana G, Makriyannis A, Lyn SY, Betramo M, Piomelli D (1997): Inhibition of intestinal motility by anandamide, an endogenous cannabinoid. *Eur J Pharmacol* 340:R7–R8
- Chan HSL, Correia JA, Macleod SM (1987): Nabilone versus prochlorperazine for control of cancer chemotherapy-induced emesis in children: A double blind, crossover trial. *Pediatrics* 79:946–952
- Churchfield S (1990): *The Natural History of Shrews*. Ithaca NY, Cornell University Press, Comstock Publishing Associates.
- Colombo G, Agabio R, Lobina C, Reali R, Gessa GL (1998): Cannabinoid modulation of intestinal propulsion in mice. *Eur J Pharmacol* 344:67–69
- Compton DR, Aceto MD, Lowe J, Martin BR (1996): In vivo characterization of a specific cannabinoid receptor antagonist (SR 141716A). Inhibition of Δ^9 -tetrahydrocannabinol-induced responses and apparent agonist activity. *J Pharmacol Exp Therap* 277:586–594
- Cook SA, Lowe JA, Martin BR (1998): CB₁ receptor antagonist precipitates withdrawal in mice exposed to Δ^9 -tetrahydrocannabinol. *J Pharmacol Exp Therap* 285:1150–1156
- Dalzell AM, Bartlett H, Lilleyman JS (1986): Nabilone: an alternative antiemetic for cancer chemotherapy. *Arch Disease Childhood* 161:502–505
- Darmani NA (2000): Δ^9 -Tetrahydrocannabinol prevents chemotherapy-induced vomiting via activation of cannabinoid CB₁ receptors. *Soc Neurosci Abs* 26(part 2):P2160
- Darmani NA (1998): Serotonin 5-HT₃ receptor antagonists prevent cisplatin-induced emesis in *Cryptotis parva*: A new experimental model of emesis. *J Neural Transm* 105:1143–1154
- Darmani NA, Pandya DK (2000): Involvement of other neurotransmitters in behaviors induced by the cannabinoid CB₁ receptor antagonist SR 141716A in naive mice. *J Neural Transm* 107:931–945
- Darmani NA, Zhao W, Ahmad B (1999): The role of D₂ and D₃ dopamine receptors in the mediation of emesis in *Cryptotis parva* (the least shrew). *J Neural Transm* 106:1045–1061
- Feigenbaum JJ, Richmond SA, Weissman Y, Mechoulam R (1989): Inhibition of cisplatin-induced emesis in the pigeon by a nonpsychotropic synthetic cannabinoid. *Eur J Pharmacol* 169:159–165
- Gralla RA (1999): Cannabinoids and the control of chemotherapy-induced nausea and vomiting. In Nahas GG, Sutin KM, Harvey DJ, Agurell S (eds), *Marijuana and Medicine*. Totowa, NJ, Humana Press, pp 599–610
- Izzo AA, Mascolo N, Borrelli F, Capasso F (1999): Defecation, intestinal fluid accumulation and motility in rodents: Implications of cannabinoid CB₁ receptors. *Naunyn-Schmied Arch Pharmacol* 359:65–70
- Krowicki ZK, Moerschbaecher JM, Winsauer PJ, Digavalli SV, Hornby PJ (1999): Δ^9 -Tetrahydrocannabinol inhibits gastric motility in the rat through cannabinoid CB₁ receptors. *Eur J Pharmacol* 371:187–196
- Lehmann A, Karrberg L (1996): Effects of N-methyl-D-aspartate receptor antagonists on cisplatin-induced emesis in the ferret. *Neuropharmacology* 35:475–481
- London SW, McCarthy LE, Borison HL (1979): Suppression of cancer chemotherapy-induced vomiting in the cat by nabilone, a synthetic cannabinoid. *Pro Soc Exp Biol Med* 160:437–440
- Lucot JB (1998): Effects of N-methyl-D-aspartate antagonists on different measures of motion sickness in cats. *Brain Res* 47:407–411
- Matsuda LA (1997): Molecular aspects of cannabinoid receptors. *Critical Rev Neurobiology* 11:143–166
- Matsuki N, Ueno S, Kaji T, Ishihara A, Wang C-H, Saito H (1988): Emesis induced by cancer chemotherapeutic agents in the *Suncus murinus*: A new experimental model. *Jpn J Pharmacol* 48:303–306
- McCarthy LE, Borison HL (1981): Antiemetic activity of N-methyllevonantradol and nabilone in cisplatin-treated cats. *J Clin Pharmacol* 21:305–375
- McCarthy LE, Flora KP, Vishnuvajjala R (1984): Antiemetic properties and plasma concentrations of delta-9-tetrahydrocannabinol against cisplatin vomiting in cats. In Agurell S, Dewey WL, Willette RD (eds), *The Cannabinoids: Chemical, Pharmacological and Therapeutic Aspects*. London, Academic Press, pp 895–902
- Naylor RJ, Rudd JA (1996): Mechanisms of chemotherapy/radiotherapy-induced emesis in animal models. *Oncology* 53(Suppl 1):8–17
- Pertwee RG (1997): Pharmacology of cannabinoid CB₁ and CB₂ receptors. *Pharmacol Therap* 74:129–180
- Rinaldi-Carmona M, Barth F, Healume M, Shire D, Calandra G, Congy C, Martinez S, Maruani J, Nélat G, Caput D, Ferrara P, Soubrie P, Breliere J-C, Le Fur G (1994): SR 141716A, a potent and selective antagonist of the brain cannabinoid receptor. *FEBS Lett* 350:240–244
- Rinaldi-Carmona M, Barth F, Millan J, Derocq JM, Casellas P, Congy C, Oustric D, Sanran M, Bouaboula M, Calandra P, Portier M, Shire D, Breliere IC, Le Fur JL (1998): SR 144528, the first potent and selective antagonist of CB₂ cannabinoid receptor. *J Pharmacol Exp Therap* 284:644–650
- Rubino T, Patrini G, Massi P, Fuzio D, Vigano D, Giagnoni G, Parolaro D (1998): Cannabinoid-precipitated withdrawal: A time-course study of the behavioral aspect and its correlation with cannabinoid receptors and G-protein expression. *J Pharmacol Exp Therap* 285:813–819
- Shook JE, Burks TF (1989): Psychoactive cannabinoids reduce gastrointestinal propulsion and motility in rodents. *J Pharmacol Exp Therap* 249:444–449
- Stark P (1982): The pharmacologic profile of nabilone: A new antiemetic agent. *Cancer Treat Rev* 9(Suppl B):11–16
- Torii Y, Saito H, Matsuki N (1991): Selective blockade of cytotoxic drug-induced emesis by 5-HT₃ receptor antagonists in *Suncus murinus*. *Jpn J Pharmacol* 55:107–113
- Voth EA, Schwartz RH (1997): Medicinal application of delta-9-tetrahydrocannabinol and marijuana. *Annals Inter Med* 126:791–798