
COMMENTARY

Where's the Excitement in Psychostimulant Sensitization?

Repeated psychostimulant exposure produces sensitized behavioral responses that persist into withdrawal and are thought to model increased drug craving in psychostimulant abusers (Post and Rose 1976; Robinson and Berridge 1993; Stewart 1983). A potential mechanism for inducing behavioral sensitization could involve a transient increase in AMPA glutamate receptor-mediated excitability in ventral tegmental area (VTA) dopamine neurons (White et al. 1995; Zhang et al. 1997). Previously, this hypersensitivity was thought to involve transient increases in AMPA (and NMDA) receptor subunits after multiple drug or stress-related treatment regimens known to induce behavioral sensitization (Churchill et al., 1999; Fitzgerald et al. 1996).

However, a study by Lu, Monteggia, and Wolf published in this issue of *Neuropsychopharmacology* directly challenges the idea that psychostimulant-induced hypersensitivity in VTA dopamine neurons involves an increase in the actual amount of glutamate receptors. In this study, multiple treatment regimens with amphetamine and cocaine, known to induce behavioral sensitization, and identical to protocols used in previous studies, all failed to increase AMPA subunits in the VTA and neighboring substantia nigra. Such discrepancies in the drug abuse literature usually are attributed to differences in dose, chronicity, withdrawal time, self- versus passive drug administration, or other methodological considerations.

The study by Lu et al. (1999, 2001) used quantitative immunohistochemical and reverse transcriptase-polymerase chain reaction methods to measure GluR1-4 levels throughout the rostral-caudal extent of the VTA and substantia nigra, whereas increases in AMPA subunits previously were detected in immunoblots of VTA tissue. After a series of experiments to verify epitope and sequence specificity of the GluR1 antibody and RNA probe, respectively, the authors conducted an exhaustive and thorough search for the precise anatomical site

in VTA where the elusive upregulation in GluR1 subunits was thought to occur. They found no evidence for upregulation of AMPA subunits at the mRNA or protein level in any region of VTA and substantia nigra, whether at 16 and 24 h or 3 and 14 days after multiple chronic treatment regimens with amphetamine and cocaine. Other studies have also failed to find evidence for upregulation in mRNAs for AMPA GluR1 or NMDA NR1 subunits, suggesting that reported increases in GluR1 and NR1 protein may not involve changes in GluR1 gene expression (Bardo et al., 2001; Ghasemzadeh et al., 1999). However, the present study also failed to find increases in GluR1 protein. One potential caveat to the immunohistochemical method used by Lu et al. is that GluR1 antibodies could recognize other endogenous proteins containing GluR1-like epitopes that dilute changes in GluR1 subunits themselves in native tissue slices; this problem was circumvented in previous studies in which GluR subunits were isolated by standard SDS PAGE techniques before immunolabeling. However, the authors note that their immunohistochemical method is capable of detecting changes as small as 15% in other brain regions. What may be apparent from this and other studies is that relatively similar chronic drug treatments produce markedly different effects on AMPA subunits in VTA across several laboratories (see Table 1).

In the present study, Lu et al. conclude that transient increases in GluR1 protein cannot account for transient increases in AMPA-mediated electrophysiological responses associated with the induction of behavioral sensitization, and they suggest phosphorylation or membrane trafficking of AMPA receptors as alternative hypotheses. There is considerable evidence for phosphorylation and membrane trafficking as important events in AMPA receptor plasticity in long-term potentiation (LTP) and long-term depression (LTD) (Carroll et al. 2001; Scannevin and Huganir 2000). Indeed, a recent

Table 1. Differential Alterations in Glutamate Receptor Subunits and Tyrosine Hydroxylase (TH) in VTA at Early and Late Psychostimulant Withdrawal

Chronic Cocaine	GluR1			GluR2/3			NR1			TH		
	mRNA	Protein		mRNA	Protein		mRNA	Protein		mRNA	Protein	
Early withdrawal ≤ 3 days	No Δ (Bardo et al. 2001; Lu et al. 2001)	↑ (Fitzgerald et al. 1996; Churchill et al. 1999)		No Δ (Bardo et al. 2001)	No Δ (Fitzgerald et al. 1996; Churchill et al. 1999, 2001; Lu et al. 2001)		—	↑ (Fitzgerald et al. 1996; Churchill et al. 1999; Loftis and Janowsky, 2000)		↑ (Vrana et al. 1993)	↑ (Beitner-Johnson et al. 1991; Sorg et al. 1993; Vrana et al. 1993)	
		No Δ (Lu et al. 2001)						No Δ (Lu et al. 1999)		No Δ (Sorg et al. 1993)	No Δ (Schmidt et al. 2001 ^a)	
Late withdrawal > 3 days	No Δ Chasemzadeh et al. 1999; Bardo et al. 2001)	No Δ (Churchill et al. 1999; Lu et al. 2001)		No Δ (Chasemzadeh et al., 1999) Bardo et al. 2001)	No Δ (Churchill et al. 1999; Lu et al. 2001)		No Δ (Ghasemzadeh et al. 1999)	No Δ (Churchill et al. 1999; Lu et al. 1999)		No Δ (Sorg et al. 1993)	↑ (Masserano et al. 1996)	
								↑ (Loftis and Janowsky, 2000)			No Δ (Sorg et al. 1993; Schmidt et al. 2001 ^a)	↓ (Trulson et al. 1987)

^aIndicates intravenous cocaine self-administration.

study found that a single injection of cocaine produces both behavioral sensitization and LTP in VTA dopamine neurons independent of detectable AMPA receptor up-regulation (Ungless et al. 2001), thereby implicating other possible trafficking and/or phosphorylation alterations. However, others have found that "synaptic" concentrations of AMPA receptors actually do increase after LTP in the hippocampus, and such subanatomical changes may not be detected in tissue slices or crude tissue homogenates (Heynen et al. 2001).

It is clear from these studies that drug-induced neuroadaptations in the VTA differ widely across research laboratories, even when drug regimens and withdrawal times are taken into account. Several studies on psychostimulant regulation of glutamate receptor subunits in the VTA are listed in Table 1, along with regulation of VTA tyrosine hydroxylase levels, which also differ among studies and psychostimulant treatment regimens. It is important to consider that several methodological variables differ between these studies, such as rat strains and vendors and whether animals received drug treatments in their home cages or in specific drug-associated test chambers. Drug-induced neuroplasticity is profoundly altered by certain contextual or other experience-related factors (e.g., Bell et al. 2000; Schmidt et al. 2001), and the potential influence of these factors should not be underestimated.

Results from behavioral studies are sometimes encumbered by the fact that certain effects clearly and easily obtained in one laboratory are difficult to repeat in another. Even when animal strains are held constant and procedures are carefully controlled, different, and even opposite, results can be found (Crabbe et al. 1999). The findings of Giorgetti and colleagues (Giorgetti et al. 2001), when taken in view of previous studies, highlight the fact that neurobiological studies can suffer a similar dilemma. However, in scientific research, these discrepancies serve well to remind us that a single study cannot, and should not, ever be considered definitive. Scientific progress benefits from both convergent and divergent lines of evidence, and discrepant findings such as these ultimately help to illuminate intricate facets of neural and behavioral plasticity that may have important implications for drug addiction research and treatment.

David W. Self, Ph.D.
Department of Psychiatry
UT Southwestern Medical Center
Dallas, TX

REFERENCES

- Bardo MT, Robinet PM, Mattingly BA, Marguilles JE (2001): Effect of 6-hydroxydopamine or repeated amphetamine treatment on mesencephalic mRNA levels for AMPA glutamate receptor subunits in the rat. *Neurosci Lett* 302:133–136
- Beitner-Johnson D, Guitart X, Nestler EJ (1991): Dopaminergic brain reward regions of Lew's and Fischerrats display different levels of tyrosine hydroxylase and other morphine and cocaine-regulated phosphoproteins. *Brain Res* 561: 147–150
- Bell K, Duffy P, Kalivas PW (2000): Context-specific enhancement of glutamate transmission by cocaine. *Neuropsychopharmacology* 23:335–344
- Carroll RC, Beattie EC, von Zastrow M, Malenka RC (2001): Role of AMPA receptor endocytosis in synaptic plasticity. *Nature Rev Neurosci* 2:315–324
- Churchill L, Swanson CJ, Urbina M, Kalivas PW (1999): Repeated cocaine alters glutamate receptor subunit levels in the nucleus accumbens and ventral tegmental area of rats that develop behavioral sensitization. *J Neurochem* 72:2397–2403
- Crabbe JC, Wahlsten D, Dudek BC (1999): Genetics of mouse behavior: Interactions with laboratory environment. *Science* 284:1670–1672
- Fitzgerald LW, Ortiz J, Hamedani AG, Nestler EJ (1996): Regulation of glutamate receptor subunit expression by drugs of abuse and stress: Common adaptations among cross-sensitizing agents. *J Neurosci* 16:274–282
- Ghasemzadeh MB, Nelson LC, Lu X-Y, Kalivas PW (1999): Neuroadaptations in ionotropic and metabotropic glutamate receptor mRNA produced by cocaine treatment. *J Neurochem* 72:157–165
- Giorgetti M, Hotsenpiller G, Ward P, Tepper T, Wolf ME (2001): Amphetamine-induced plasticity of AMPA receptors in the ventral tegmental area: Effects on extracellular levels of dopamine and glutamine in freely moving rats. *J Neurosci* 21:6362–6369
- Heynen AJ, Quinlan EM, Bae DC, Bear MF (2001): Bidirectional, activity-dependent regulation of glutamate receptors in the adult hippocampus in vivo. *Neuron* 28:527–536
- Loftis JM, Janowsky A (2000): Regulation of NMDA receptor subunits and nitric oxide synthase expression during cocaine withdrawal. *J Neurochem* 75:2040–2050
- Lu W, Monteggia LM, Wolf ME (1999): Withdrawal from repeated amphetamine administration reduces NMDAR1 expression in the rat substantia nigra, nucleus accumbens and medial prefrontal cortex. *Eur J Neurosci* 11:3167–3177
- Lu W, Monteggia LM, Wolf ME (2001) Repeated administration of amphetamine and cocaine does not alter AMPA receptor subunit expression in the rat midbrain. *Neuropsychopharmacology* 26:1–13.
- Masserano JM, Baker I, Natsukari N, Wyatt RJ (1996): Chronic cocaine administration increases tyrosine hydroxylase activity in the ventral tegmental area through glutaminergic- and dopaminergic D2-receptor mechanisms. *Neurosci Lett* 217:73–76
- Post R, Rose H (1976): Increasing effects of repetitive cocaine administration in rats. *Nature* 260:731–734
- Robinson TE, Berridge KC (1993): The neural basis of drug craving: An incentive-sensitization theory of addiction. *Brain Res Rev* 18:247–291
- Scannevin RH, Haganir RL (2000): Postsynaptic organiza-

- tion and regulation of excitatory synapses. *Nature Rev Neurosci* 1:133–141
- Schmidt EF, Sutton MA, Schad CA, Karanian DA, Brodtkin ES, Self DW (2001): Extinction training regulates tyrosine hydroxylase during withdrawal from cocaine self-administration. *J Neurosci* 21:RC137
- Sorg BA, Chen S-Y, Kalivas PW (1993): Time course of tyrosine hydroxylase expression after behavioral sensitization to cocaine. *J Pharmacol Exp Ther* 266:424–430
- Stewart J (1983): Conditioned and unconditioned drug effects in relapse to opiate and stimulant drug self-administration. *Prog Neuro-Psychopharmacol Biol Psychiatry* 7:591–597
- Trulson ME, Cannon MS, Faegg TS, Raese JD (1987): Tyrosine hydroxylase immunochemistry and quantitative light microscopic studies of the mesolimbic dopamine system in rat brain: effects of chronic methamphetamine administration. *Brain Res Bull* 18:269–277
- Ungless MA, Whistler JL, Malenka RC, Bonci A (2001): Single cocaine exposure in vivo induces long-term potentiation in dopamine neurons. *Nature* 411:583–587
- Vrana SL, Vrana KE, Koves TR, Smith JE, Dworkin SI (1993): Chronic cocaine administration increases CNS tyrosine hydroxylase enzyme activity and mRNA levels and tryptophan hydroxylase enzyme activity levels. *J Neurochem* 61:2262–2268
- White FJ, Hu XT, Zhang XF, Wolf ME (1995): Repeated administration of cocaine or amphetamine alters neuronal responses to glutamate in the mesoaccumbens dopamine system. *J Pharmacol Exp Ther* 273:445–454
- Zhang X-F, Hu X-T, White FJ, Wolf ME (1997): Increased responsiveness of ventral tegmental area dopamine neurons to glutamate after repeated administration of cocaine or amphetamine is transient and selectively involves AMPA receptors. *J Pharmacol Exp Ther* 281:699–706