

Activation of α_2 Adrenergic Receptors Suppresses Fear Conditioning: Expression of c-Fos and Phosphorylated CREB in Mouse Amygdala

M Frances Davies^{*1}, Janet Tsui², Judy A Flannery², Xiangqi Li¹, Timothy M DeLorey³ and Brian B Hoffman²,

¹Department of Anesthesia, Stanford University of Anesthesiology, Veterans Affairs, Palo Alto Health Care System, Palo Alto, CA, USA;

²Department of Medicine, Stanford University of Medicine, Veterans Affairs, Palo Alto Health Care System, Palo Alto, CA, USA; ³Molecular Research Institute, Mountain View, CA, USA

α_2 adrenergic agonists such as dexmedetomidine generally suppress noradrenergic transmission and have sedative, analgesic, and antihypertensive properties. Considering the importance of the neurotransmitter norepinephrine in forming memories for fearful events, we have investigated the acute and chronic effects of dexmedetomidine on discrete cue and contextual fear conditioning in mice. When administered before training, dexmedetomidine (10–20 $\mu\text{g/kg}$, i.p.) selectively suppressed discrete cue fear conditioning without affecting contextual memory. This behavioral change was associated with a decrease in memory retrieval-induced expression of c-Fos and P-CREB in the lateral, basolateral, and central nuclei of the amygdala. Dexmedetomidine's action on discrete cue memory did not occur in α_{2A} adrenoceptor knockout (KO) mice. When dexmedetomidine was administered after training, it suppressed contextual memory, an effect that did not occur in α_{2A} adrenoceptor KO mice. We conclude that dexmedetomidine, acting at α_{2A} adrenoceptors, must be present during the encoding process to decrease discrete cue fear memory; however, its ability to suppress contextual memory is likely the result of blocking the consolidation process. The ability of α_2 agonists to suppress fear memory may be a valuable property clinically in order to suppress the formation of memories during stressful situations.

Neuropsychopharmacology (2004) **29**, 229–239, advance online publication, 29 October 2003; doi:10.1038/sj.npp.1300324

Keywords: contextual memory; discrete cue memory; transcription factors; c-Fos; P-CREB; α_{2A} adrenoceptor subtype knockout mice; chronic administration

INTRODUCTION

Stressful situations can trigger the typical 'fight and flight' response that activates the peripheral sympathetic and central noradrenergic systems to release norepinephrine. The increased central noradrenergic release can promote memory formation evoked by the fearful stimuli (Vaandrager and de Jonge, 1996), and may explain why we remember exciting or traumatic events more vividly than everyday ones (Cahill and McGaugh, 1998). However, fear learning can be detrimental in some situations, such as for patients undergoing medical procedures, or where the persistent fear response evokes chronic stress (Svensson, 1987). Consequently, a means to selectively reduce fear learning is desirable in such circumstances.

*Correspondence: MF Davies, Department of Anesthesia, VAPAHCS, 3801 Miranda Avenue, Palo Alto, CA 94304, USA, Tel: +1 650 493 5000 ext. 64854, Fax: +1 650 852 3423, E-mail: fdavies@stanford.edu
Received 15 July 2003; revised 08 September 2003; accepted 12 September 2003

Online publication: 12 September 2003 at <http://www.acnp.org/citations/Npp09120303314/default.pdf>

Fear conditioning is a form of associative learning in which defense responses can be elicited by a neutral conditioned stimulus (CS) such as an audible tone, when it has been previously paired with an aversive unconditioned stimulus (US) such as an electric footshock. The amygdala is the major brain region responsible for pairing the CS with the US and then relaying this association onto other regions (Fanselow and Kim, 1994). Discrete cue learning to an auditory tone largely occurs in the amygdala (Rogan *et al*, 1997) whereas contextual learning also involves the hippocampus (Anagnostaras *et al*, 2001). The basolateral amygdala has reported to be crucial in the formation and storage of fear memories (Maren, 1999). In the amygdala, long-term potentiation (LTP), hypothesized to be the mechanism behind some forms of learning, has been demonstrated to require a strong noradrenergic input in order to be established (Huang *et al*, 2000; Schafe *et al*, 2001), whereas norepinephrine plays mainly a modulatory role in the hippocampus LTP (Dunwiddie *et al*, 1982; Sarvey *et al*, 1989). In both areas, phosphorylation of cyclic AMP response element-binding protein (CREB) and expression of c-Fos are critical in changing gene expression that is

important to memory formation (Dragunow *et al*, 1989; Hall *et al*, 2001; Moller *et al*, 1994). The transcription factors P-CREB and c-Fos increase during memory encoding (Tischmeyer and Grimm, 1999), consolidation (Kang *et al*, 2001), and retrieval (Hall *et al*, 2001). Whenever information is being actively processed in the brain, the memory is unstable and is vulnerable to drugs that interfere with paths leading to protein synthesis (Nader *et al*, 2000). The nature by which a drug affects these processes provides insight into the mechanisms by which the drug is likely to influence memory formation.

Stimulation of α_2 adrenoceptors (ARs) decreases norepinephrine release in the amygdala (Fendt *et al*, 1994), and therefore may decrease fear learning. However, this hypothesis has not been rigorously tested. We have addressed this problem and found that the α_2 agonist dexmedetomidine decreases fear memory but not contextual memory. To further understand the mechanism of action by which α_2 ARs may influence fear memory (Hall *et al*, 2001; Moller *et al*, 1994), we have assessed dexmedetomidine's affect on c-Fos and P-CREB production in the amygdala. In doing so, we have further elucidated the α_2 adrenoceptor subtype responsible for the observed behavioral action.

METHODS

Mice

The experimental protocol was approved by the Animal Care and Use Committee at the Veterans Administration Palo Alto Health Care System. Wild-type C57BL/6J mice (Jackson Laboratories, Davis, CA), α_{2A} AR knockout (KO) adult male mice originated by Dr Brian Kobilka (Altman *et al*, 1999) and D79N mice originated by Dr Lee Limbird (MacMillan *et al*, 1996), which contains a point mutation that functionally inactivates the α_{2A} AR, were bred in house. Both were backcrossed onto a single C57BL/6J background. They were housed in groups of eight, kept on a 12:12 h light/dark cycle and were allowed free access to food and water.

Behavioral Apparatus

Four identical chambers ($20 \times 25 \times 22 \text{ cm}^3$) (MED Associates, Georgia, VT) were used for Pavlovian fear conditioning and testing. Each chamber contained metal floor bars through which an electrical current could be passed. Before each trial, floor bars were tested for correct amperage and correct decibel levels were assessed for both the tone generator and white noise generator.

Pavlovian Fear Conditioning

Before testing each day, the mice were moved to a holding room adjacent to the testing room and allowed to acclimate for at least 30 min. The mice were placed individually in one of the four identical experimental chambers (MED Associates, St Albans, VT) that had been scented with 1% ammonium hydroxide solution before testing. Chambers were backlit with diffuse fluorescent light and a white noise generator provided 70 dB background noise. Mice were

allowed to explore the experimental chamber for 4 min, after which they were exposed to a loud tone (85 dB, 2.9 kHz) for 33 s with the last 3 s coupled with a 0.75-mA 'scrambled footshock'. This procedure was repeated for a total of three episodes with a 1-min period separating each episode. At 1 min after the final footshock, the mice were returned to their home cages. After 24 h, contextual memory was assessed by placing the mice back into the freshly rescented (1% ammonium hydroxide) conditioning chambers in which they were trained, for a 4-min test period in the absence of footshock. Conditioned fear to the context was assessed by measuring the freezing response, according to the methods of Fanselow and Bolles (Fanselow and Bolles, 1979). Freezing was defined as the absence of all visible movements of the body and vibrissae aside from those necessitated by respiration. For the discrete memory assessment, the context was altered by changing the visual, tactile, and olfactory cues. The new context consisted of chambers scented with 2.5% lemon juice concentrate and reconfigured with three white opaque plastic inserts that convert the inner dimensions of the chamber to be triangular in shape rather than the original rectangular shape. Fluorescent backlighting and white noise generation were switched off. Baseline fear to the new context was assessed by recording the freezing response for 4 min. Following this baseline exploration period, the tone originally presented in the training session was sounded for 4 min during which time freezing behavior was again recorded. An observer, blind to the genotype of the mice, scored each mouse every 8 s, for a total of 4 min, for the presence or absence of freezing. These data were transformed to a percentage of total observations. Data were analyzed by one-way analysis of variance (ANOVA) using GraphPad PRISM 3 (GraphPad Software). Separate treatment effects between groups were analyzed *post hoc* using Dunnett's comparisons.

Immunohistochemistry

At 1 h after discrete cue memory assessment, mice were euthanized with carbon dioxide and perfused with 20 ml of ice-cold phosphate-buffered saline (PBS) followed by 4% paraformaldehyde in PBS injected into the left ventricle of the heart. Brains were then harvested, fixed in 4% paraformaldehyde-PBS for 6 h, and cryoprotected in 30% sucrose-phosphate buffer (PB) overnight. The brains were sliced coronally into 3–4-mm thick slices in the regions of the hippocampus and amygdala using an adult mouse coronal 1 mm brain matrix (Ted Pella, Redding, CA), embedded, and stored at -80°C until sectioning. After being cut into 40- μm sections with a cryostat microtome (Leica, CM 1850 Wetzlar, 35578 Germany) and placed in 0.1 M PB, these floating sections were washed with PB twice and then blocked with a 3% solution of normal goat serum (NGS) in buffer 1 (0.1 PBS, NGS, and Triton X-100) for 5 h followed by overnight incubation on a shaker with either anti-c-Fos (1:10 000, Santa Cruz Biotechnology, CA) or anti-P-CREB (1:500, Cell Signaling, Beverly, MA) polyclonal rabbit IgG. Tissues were then washed with buffer 1 and incubated with secondary biotinylated anti-rabbit IgG (Vector Labs, Burlingame, CA) for 2 h. After washing with buffer 1, VECTASTAIN ABC reagent (Vector Labs, Burlin-

game, CA) was added and allowed to adhere to the secondary antibody for 1 h. The sections were subsequently washed with buffer 2 (0.1 PBS and Triton X-100), stained with DAB (Vector Labs, Burlingame, CA) substrate for 1–4 min, and washed in deionized water. Sections were mounted onto precoated Frost+ slides (Fisher Scientific, Pittsburgh, PA) and air-dried overnight. Slides were dehydrated in 100% alcohol for 10 min, submerged in a series of xylene washes, and covered with glass coverslips using Permount mounting medium (Fisher Scientific, Pittsburgh, PA). Slides were viewed and photographed with a light microscope connected to a SPOT RT Slider Camera (Diagnostic Instruments, Sterling Heights, MI). Amygdala and hippocampal areas were identified with the help of *Mouse Brain atlas* (2001) (Paxinos and Franklin, 2001), and positive c-Fos and P-CREB spots were counted using NIH Image. Digital photographs of every fourth section were counted and individual spots were counted and marked on the photographs. The averages were taken of all sections that corresponded to one atlas coronal section. This data point was then averaged across mice treated in the same manner.

Drugs

Dexmedetomidine (10–20 μ g/kg) or saline vehicle was injected intraperitoneally in a volume of 10 ml/kg.

Statistics

Statistical significance was tested with either one-way ANOVA with Dunnett's *post hoc* test or Student's *t*-test where appropriate, using Prism version 3.0a for Macintosh (GraphPad Software, San Diego, CA, USA).

RESULTS

Changes in norepinephrine release could potentially augment fear learning in a variety of ways. Information gathered through the five senses is initially stored in a sensory buffer. If these impressions persist, they may be encoded into short-term memory storage. These memories in turn may be consolidated into long-term memories. When information from long-term storage is required to perform certain tasks, memories are retrieved. To determine the processes that are sensitive to drug administration, it is necessary to examine its effect at each phase of the memory creation process.

It is critical to memory formation that the US and CS are both perceived. Since dexmedetomidine has analgesic actions in a variety of models (Guo *et al*, 1996; Hayashi *et al*, 1995, 1996), we were concerned that it might suppress the noxious nature of the US, namely electric footshock. Consequently, we undertook preliminary experiments to assess whether the footshock was sufficiently salient to elicit pain when dexmedetomidine was administered. To do this, we determined the shock threshold required to elicit flinching responses, 30 min after the test dose of dexmedetomidine (20 μ g/kg i.p.) was administered. Unexpectedly, dexmedetomidine actually decreased thresholds in C57BL/6J wild-type mice (Table 1) although it had no effect in the D79N or α_{2A} AR KO mice. To ensure that the mice exhibited

Table 1 Evaluation of Pain Perception for C57BL/6, D79N, and α_{2A} AR KO Mice Injected with Dexmedetomidine

	Flinch threshold (mA)
C57BL/6	
Saline	0.17 \pm 0.02
Dexmedetomidine	0.16 \pm 0.01
D79N	
Saline	0.18 \pm 0.01
Dexmedetomidine	0.20 \pm 0.02
α_{2A} AR KO	
Control	0.21 \pm 0.01
Dexmedetomidine	0.21 \pm 0.01

Mice were injected with either dexmedetomidine (20 μ g/kg i.p.) or saline and then placed in the experimental chambers 30 min after injection. The intensity of the footshock was increased in increments of 0.05 mA and the intensity required to evoke flinching was recorded. *N* = 4.

nociceptive behavior to the US in subsequent experiments, a supramaximal shock of 0.75 mA was used for all strains. We have demonstrated in a previous study (Davies *et al*, 2003) that applying the tone alone or shock without pairing leads to a lack of freezing to the tone when sounded during discrete cue testing.

Acute Effect of Dexmedetomidine on Fear Conditioning

In assessing the effect of a drug on fear memory, it is first necessary to ascertain that the drug does not alter the perception of either the context or tone used in the training period. Figure 1a shows the exploratory behavior of the mice when they are first exposed to the training cage and for the three intervals after the tone-shock pairing. The footshock itself significantly reduced exploratory behavior in saline-treated mice during all three intershock intervals (Figure 1a), indicating that it was salient enough to be remembered in the short term. Mice administered dexmedetomidine (20 μ g/kg) exhibited a significant decrease in the initial chamber exploration prior to the footshock when compared to saline-treated mice (Figure 1a). However, after the first shock, the dexmedetomidine-treated group moved significantly less than the saline-treated nonshocked and shocked mice. It was clear that the exploratory behavior of dexmedetomidine-treated mice was likewise decreased by the first footshock, indicating that the mice experienced both pain and demonstrated short-term memory.

When saline- and dexmedetomidine-treated mice are placed in a novel contextual environment, different than the context in which they were trained, they both exhibited less than 10% freezing (Figure 1b) suggesting that neither group has generalized their fear of the original training context to the new test context. Mice placed in the novel environment but not shocked during the training period exhibited minimal freezing in the novel context whether with (2.6 \pm 1.5%) or without (1.6–1.6%) the tone. Wild-type mice that were fear conditioned to a tone froze approximately 50% of the total possible time when re-exposed to




Figure 1 Effect of preadministering dexmedetomidine (20 µg/kg i.p.) on fear-conditioned C57BL/6 wild-type mice. (a) Chamber exploration during the initial 4-min acclimation period in the training chamber was significantly decreased by administration of dexmedetomidine (20 µg/kg i.p.) 30 min before presentation of the first shock. The initial exploration data are expressed as the average centerline crosses per minute. During subsequent 1-min intervals following each tone and shock presentation, saline-treated mice that were shocked (white bars) exhibited a significant reduction in exploration compared to saline-treated mice that were not shocked (light gray bar) (### $p < 0.001$, # $p < 0.05$, $N = 6-8$). Dexmedetomidine-treated mice that were shocked (black bars) explored significantly less than saline-treated mice that were shocked (white bars) (*** $p < 0.001$, ** $p < 0.01$, $N = 6-8$). (b) Dexmedetomidine had no effect on freezing when 24 h after training the mice were put into a novel environment different than the one trained in and allowed to freely explore for 4 min. (c) Dexmedetomidine reduced freezing in the novel environment when the tone was presented for 4 min. Bars represent mean \pm SEM. Statistically significant difference to saline-treated controls: ** $p < 0.01$, $N = 6$.

the tone when placed in a novel context (discrete cue) 24 h after training (Figure 1c). Mice injected with dexmedetomidine (20 µg/kg i.p.) 30 min before footshock tone pairing, froze significantly less to the tone presentation compared to




Figure 2 Effect of preadministration of a nonsedating dose of dexmedetomidine before fear-conditioned training in C57BL/6 wild-type mice. Dexmedetomidine (10 µg/kg i.p.) was administered 30 min before fear-conditioned training. (a) Chamber exploration during training was not affected by administration of this dose of dexmedetomidine. (b) When tested 24 h after training, the mice treated with dexmedetomidine froze significantly less in response to the tone. Bars represent mean \pm SEM. Statistically significant difference to saline-treated controls: * $p < 0.05$, $N = 6$.

controls when tested 24 h after training. To determine whether the sedative action of dexmedetomidine (20 µg/kg) was responsible for the suppression of the discrete cue fear conditioning, a lower, nonsedating dose of 10 µg/kg dexmedetomidine was also tested. This dose did not significantly affect the initial exploration of the chamber (Figure 2a), but significantly reduced freezing behavior compared to control in the presence of the tone when presented 24 h after training (Figure 2b). This result suggests that the sedative action of dexmedetomidine administration is not responsible for its fear reducing action.

To determine whether dexmedetomidine suppression of discrete cue freezing is attributable to an effect on the encoding or on the consolidation phase of fear memory, dexmedetomidine (20 µg/kg) was also given immediately after training to a separate set of wild-type mice. As expected, the exploratory behavior during training in the two groups of mice was the same (Figure 3a). Under these conditions, the drug had no effect on discrete cue freezing behavior compared to control (Figure 3b). Taken together with results presented in Figure 2, these findings support the hypothesis that dexmedetomidine influences the encoding rather than the consolidation processes.

Induction of P-CREB Immunoreactivity and c-Fos Immunoreactivity Within the Amygdala by Fear Conditioning

Transcription factors regulating gene expression in the lateral, basolateral nucleus, and the central nucleus of the amygdala, are likely to be important to the development of fear memory (Ferry *et al*, 1999b). Auditory and other




Figure 3 Effect of dexmedetomidine (20 μ g/kg i.p.) administered immediately after training on fear-conditioned C57BL/6 wild-type mice. Dexmedetomidine (20 μ g/kg i.p.) was administered immediately after training. (a) No significant difference between dexmedetomidine treated and untreated mice was observed during the initial chamber exploration period prior to sounding of the tone and presentation of the shock. (b) There was no significant difference between the two groups in freezing response to the tone during discrete cue assessment tested one day after drug administration and training. Bars represent mean \pm SEM. Statistically significant difference to saline-treated controls: * $p < 0.05$, ** $p < 0.01$, $N = 6$.

sensory inputs generally terminate in the lateral amygdala (LeDoux *et al*, 1990; Romanski *et al*, 1993), and destruction of this area interferes with discrete cue conditioning (LeDoux *et al*, 1990). The lateral amygdala is the region proposed to be responsible for associating the CS, the auditory tone, with the nociceptive information generated by the US, the footshock (LeDoux *et al*, 1990). The central nucleus is essential for autonomic and emotional responses to stressful stimuli (LeDoux *et al*, 1988). The basolateral amygdala appears to play a role in conditioned aversion (Amorapanth *et al*, 2000) or inhibitory avoidance (Wilensky *et al*, 2000). Long-term memory has been reported to be facilitated by CREB protein overexpression in the basolateral amygdala (Josselyn *et al*, 2001). Moreover, fear memory retrieval has been shown to induce phosphorylation of CREB and expression of c-Fos in the basolateral amygdala (Hall *et al*, 2001) as well as the induction of a protein-synthesis-dependent reconsolidation of memories within the amygdala (Nader *et al*, 2000).

We wondered if the fear suppressive action of acute administration of dexmedetomidine given before training is associated with changes in phosphorylation of CREB and c-Fos expression in the three amygdala nuclei implicated in fear conditioning following retrieval. In untrained wild-type mice, there was no indication of P-CREB (Figure 4a) or c-Fos (Figure 5a) positive neurons in the lateral, basolateral, or central nucleus of the amygdala. In fear-conditioned mice killed 1 h after discrete cue testing, to stimulate fear




Figure 4 Effect of preadministration of dexmedetomidine on the number of cells expressing P-CREB in the amygdala evoked by discrete cue retrieval tested 24 h later. Immunohistochemical staining of P-CREB in a coronal section of the amygdala of C57BL6 mice (a) exposed to the training box and tone but not given a footshock, (b) given saline then trained, or (c) given dexmedetomidine (20 μ g/kg, i.p.) 30 min before fear conditioning and then tested on the following day. Mice were killed 1 h after discrete cue testing. The numbers in the gray boxes refer to the areas that are enlarged in the side figures labeled. The (1) lateral, (2) basolateral, and (3) central nucleus of the amygdala of dexmedetomidine-treated mice had fewer P-CREB positive cells than saline-treated mice. Bregma -1.58 mm; interaural 2.22 mm; Bar = 100 μ m.

retrieval, the number of P-CREB (Figure 4b) and c-Fos (Figure 5b) positive neurons significantly increased in all three amygdaloid nuclei. When dexmedetomidine was administered before training, the number of P-CREB (Figure 6a–c) and c-Fos (Figure 6d–f) positive neurons were significantly reduced in all three amygdaloid regions.

As the hippocampus is thought to play a role in contextual memory, we endeavored to examine P-CREB and c-Fos expression in the hippocampus upon retrieval. We observed no expression of either transcription factor in the hippocampus following context re-exposure in the presence or absence of dexmedetomidine (data not shown).




Figure 5 Effect of preadministration of dexmedetomidine on the number of cells expressing c-Fos in the amygdala evoked by discrete cue retrieval tested 24 h later. Immunohistochemical staining of c-Fos in a coronal section of the amygdala of C57BL6 mice (a) exposed to the training box and tone but not given a footshock, (b) given saline then trained, or (c) given dexmedetomidine (20 µg/kg) 30 min before fear conditioning and then tested on the following day. Mice were killed 1 h after discrete cue testing. The numbers in the gray boxes refer to the areas that are enlarged in the side figures labeled. The (1) lateral, (2) basolateral, and (3) central nucleus of the amygdala of dexmedetomidine-treated mice had fewer c-Fos positive cells than saline-treated mice. Bregma -1.58 mm; interaural 2.22 mm; Bar = 100 µm.

Effect of Dexmedetomidine on Discrete Cue Memory in Mice Deficient in α_{2A} ARs

There are three α_2 AR subtypes in the brain A, B and C (Blaxall *et al*, 1994; Flordellis *et al*, 1991; Regan *et al*, 1988), which could potentially mediate the amnesic effects of the nonselective α_2 agonist dexmedetomidine. As many of the central effects of α_2 agonists are mediated by the α_{2A} AR (Lakhlani *et al*, 1997), we sought to determine whether this AR subtype was also the major mediator of the fear amnesic effect by which dexmedetomidine acted. To test this, dexmedetomidine (20 µg/kg) was administered to α_{2A} AR KO mice prior to training. This dose was found not to

affect the initial chamber exploration compared to saline (Figure 7a). At 24 h after training, mice were placed in a novel context, different from the one they were trained in. Both dexmedetomidine-treated α_{2A} AR KO mice and wild-type mice exhibited similar low levels of freezing in the novel context prior to tone (Figure 7b). In the presence of the tone dexmedetomidine-treated α_{2A} AR KO mice exhibited robust freezing in the discrete cue test compared to the control (Figure 7c). To further test the role of the α_{2A} AR, the effect of dexmedetomidine was also assessed in transgenic D79N mice that are functionally deficient in the α_{2A} AR (MacMillan *et al*, 1998). A similar behavioral pattern was observed in the transgenic D79N mice with dexmedetomidine having no effect on initial chamber exploration prior to training (Figure 8a), there was no change in freezing behavior in response to a context different from training (Figure 8b), and failed to reduce freezing behavior to the tone presentation in comparison to wild-type mice receiving dexmedetomidine (Figure 8c). Combined, these results demonstrate that the inhibitory effects of dexmedetomidine on discrete cue freezing are mediated through α_{2A} ARs.

Effect of Dexmedetomidine on Contextual Memory: The Role of α_{2A} ARs

Contextual memory differs from discrete cue memory in that the hippocampus plays a crucial role in contextual memory (Anagnostaras *et al*, 2001). Dexmedetomidine (10 µg/kg i.p.) administered 30 min before training had no effect on contextual memory when assessed 24 h later (Figure 9a). Unexpectedly, we found that administration of dexmedetomidine (20 µg/kg i.p.) to wild-type mice immediately following training decreased their freezing behavior in the contextual memory test (Figure 9b). To determine if the α_{2A} AR was responsible for this retrograde amnesic effect of dexmedetomidine, this experiment was repeated using α_{2A} AR KO mice (Figure 9c) and D79N mice (Figure 9d). Both α_{2A} AR KO and D79N mice given dexmedetomidine and placed in the same context froze to the same extent as controls not given dexmedetomidine, indicating that the α_{2A} AR mediates this retrograde amnesic action.

DISCUSSION

This study demonstrates that activation by dexmedetomidine of α_{2A} AR suppresses discrete cue fear conditioning only when administered prior to training, but not when given immediately after training. This suggests that α_2 AR agonists modulate the encoding rather than the consolidation process of memory formation. The lack of effect of dexmedetomidine on contextual memory when administered prior to training supports the concept that the noradrenergic system is not essential for the encoding of contextual memory. However, its inhibitory effect when administered after training points to a role for the noradrenergic system in the consolidation phase of spatial memory as has been observed by others when exogenous noradrenergic agents were administered after training (Cahill, 2000; Hatfield and McGaugh, 1999). Our results




Figure 6 Effect of dexmedetomidine (20 $\mu\text{g/kg}$ i.p.) administered 30 min before fear-conditioning training session on retrieval-induced expression of P-CREB (a–c) and c-Fos (d–f) in the lateral (a, d), basolateral (b, e), and central (c, f) nuclei. On day 1 wild-type mice were either injected with saline or dexmedetomidine (20 $\mu\text{g/kg}$ i.p.) 30 min before training, on day 2 the mice were placed in a novel environment and presented with the conditioning stimulus tone in the absence of footshock for 4 min. Mice were then removed from the chamber and killed 1 h after discrete cue testing. Bars represent mean \pm SEM. Statistically significant difference to saline-treated controls * $p < 0.05$, ** $p < 0.01$, $N = 3$ –4.

strongly suggest that the α_{2A} AR is largely responsible for the fear suppressing action of α_2 adrenergic agonists as shown by the studies performed in the α_{2A} AR KO and D79N mice. Furthermore, when dexmedetomidine was administered prior to training, we observed a reduction in phosphorylation of CREB and expression of c-Fos in the lateral, basolateral, and central nuclei of the amygdala evoked by memory retrieval in the discrete cue test, suggesting that it was able to reduce the establishment of long-term memory formation in these structures.

Central release of norepinephrine plays an important role in the reinforcement of fear memory likely through activation of β (Ferry and McGaugh, 1999) and α_1 ARs (Ferry *et al*, 1999b). It has been established that locus coeruleus noradrenergic neurons are activated by the US of the footshock (Ishida *et al*, 2002; Sara, 1985) as well as other stressful stimuli (Abercrombie and Jacobs, 1987). Activation of either the β_2 (Ferry and McGaugh, 1999) or α_1 (Ferry *et al*, 1999b) ARs increases attention and arousal, and enhances memory. CREB-dependent transcription is required for cellular events underlying long-term memory in *Aplysia*, *Drosophila*, mice, and rats (Impey *et al*, 1998; Lamprecht *et al*, 1997). Both the behavioral signs of fear freezing and the increase in P-CREB in the amygdala are sensitive to administration of β adrenergic antagonists (Przybylski *et al*, 1999). A cascade of intracellular events that culminate in increased intracellular $[\text{Ca}^{2+}]$ or $[\text{cAMP}]$ can lead to phosphorylation and activation of CREB (Lin *et al*, 1998; Takuma *et al*, 1997). Norepinephrine is ideally suited to initiate phosphorylation of CREB as it can activate G_s -coupled β ARs to increase cAMP levels (Reisine *et al*, 1983) and activate α_1 ARs to increase intracellular $[\text{Ca}^{2+}]$ (Nelemans and den Hertog, 1987). P-CREB bound to promoter CRE sites may then enhance gene transcription including the *fos* gene (Simpson and McGinty, 1994).

Furthermore, it has been demonstrated by immunocytochemical studies in lateral amygdala that both repeated tetanizing stimulation and forskolin administration, which enhances adenylyl cyclase activity, stimulate the phosphorylation of CREB (Huang *et al*, 2000). The ability of dexmedetomidine to suppress discrete fear memory and inhibit retrieval-induced P-CREB and c-Fos expression in the amygdala is consistent with its ability to suppress norepinephrine release or by functionally antagonizing the β AR mediated increase in cAMP (Mori-Okamoto *et al*, 1991).

The site of action of dexmedetomidine in the neuronal circuits involved in fear conditioning is unclear. Our results point to the α_{2A} AR being the prime receptor subtype mediating the suppressant action of dexmedetomidine on discrete cue and contextual fear learning. α_{2A} ARs are expressed in the amygdala where they are located on both the postsynaptic amygdala neurons or presynaptically on noradrenergic terminals (Talley *et al*, 1996). Immunocytochemical studies of the rat brainstem have shown that α_{2A} receptors are found on the axon terminals and dendrites of noradrenergic locus coeruleus neurons that project to the lateral and basolateral amygdala nuclei (Lee *et al*, 1998). Noradrenergic neurons within the nucleus tractus solitarius (NTS) project to the central nucleus (Asan, 1998), and the activation of the NTS (Zardetto-Smith and Gray, 1990) contributes to an increased norepinephrine release in the amygdala and enhances memory (Williams *et al*, 2000). Norepinephrine release from the ascending noradrenergic tracts is also under presynaptic control of both α_{2C} and α_{2A} ARs (Bucheler *et al*, 2002), where they act as an autoreceptors on noradrenergic terminals. The absence of α_{2A} AR-mediated presynaptic inhibition in α_{2A} AR-deficient mice would likely promote a stronger release of norepinephrine evoked by the US to promote discrete cue




Figure 7 Effect of preadministration of dexmedetomidine administered 30 min before training on fear-conditioned α_{2A} AR KO mice. Dexmedetomidine (20 μ g/kg) administered 30 min before training on α_{2A} AR KO mice and discrete cue memory was tested the next day (a) Cage exploration during training was not significantly affected by dexmedetomidine. (b) Dexmedetomidine did not significantly affect freezing behavior of α_{2A} AR KO mice compared to wild-type controls placed in the novel environment. (c) There was no significant difference in freezing response between the two mouse groups to the tone during the discrete cue assessment. Bars represent mean \pm SEM; $N = 4$.

fear encoding (Bucheler *et al*, 2002). Indeed, we have found that α_{2A} AR-deficient mice exhibit a higher level of freezing than controls in the discrete test (Davies *et al*, 2003).

The ability of dexmedetomidine to suppress c-Fos and P-CREB expressions in the amygdala of mice fear conditioned to a discrete cue tone suggests that the norepinephrine release is also reduced in all three amygdaloid nuclei. α_2 Adrenergic agonists have been found to block fear-startle when locally injected into the lateral nucleus of the amygdala (Schulz *et al*, 2002), whereas local injection of the nonselective α_2 antagonist atipamezole into the junction of the central and lateral amygdaloid nucleus induces c-Fos expression in the amygdala (Stone *et al*, 1997). Afferents from the central nucleus project to and affect brainstem structures involved in the control of the autonomic nervous system (LeDoux *et al*, 1988). These reports taken together with our results support the idea that dexmedetomidine affects both the emotional and autonomic expression of fear (Asan, 1998).




Figure 8 Effect of preadministration of dexmedetomidine administered 30 min before training on fear-conditioned D79N mice. Dexmedetomidine (20 μ g/kg i.p.) was administered 30 min before fear conditioned-training began. (a) Dexmedetomidine did not significantly decrease initial chamber exploration during the training session. (b) There was no difference in the freezing behavior between saline- and dexmedetomidine-treated mice placed 24 h later in a new context. (c) There was no significant difference in freezing response to the tone presentation during discrete cue assessment. Bars represent mean \pm SEM; $N = 4$.

We have found that dexmedetomidine selectively decreases discrete cue freezing without affecting encoding of contextual memory. This implies that administration of an α_2 adrenergic agonist selectively affects auditory cue memory but not contextual memory. This is consistent with studies in the mice deficient in tyrosine hydroxylase, the rate-limiting enzyme in the synthesis of norepinephrine, in which these mice displayed impairment in a water-finding task, active avoidance, cued fear conditioning, and conditioned taste aversion but exhibited normal spatial learning and hippocampal LTP (Kobayashi *et al*, 2000). Others have found that blockade of β ARs by propranolol affects spatial memory when given after training (Cahill, 2000). Therefore, suppression of the noradrenergic system may cause retrograde spatial amnesia. There has been an ongoing controversy on whether the noradrenergic system affects acquisition or retention of memory, where auditory cued fear conditioning points to a noradrenergic influence on acquisition (Wilensky *et al*, 1999), the more spatially




Figure 9 Effect of dexmedetomidine on contextual memory: (a) dexmedetomidine (10 μ g/kg i.p.) administered 30 min before training on fear-conditioned C57BL/6 wild-type mice had no effect on contextual memory when tested 24 h later; (b) dexmedetomidine (20 μ g/kg i.p.), administered to mice immediately after training which were then tested 24 h later in the same context, significantly reduced freezing compared to saline-treated C57BL/6 wild-type mice; (c) in α_{2A} AR KO mice, dexmedetomidine (20 μ g/kg i.p.) administered immediately after fear-conditioned training had no effect on freezing induced by exposure to the training environment when tested 24 h; (d) in D79N mice, dexmedetomidine (20 μ g/kg i.p.) administered immediately after training had no effect on freezing when tested 24 h. Bars represent mean \pm SEM. Statistically significant difference to saline-treated controls: * $p < 0.05$, $N = 5-6$.

oriented task of inhibitory avoidance indicates that consolidation is affected (LaLumiere *et al*, 2003; Weisskopf *et al*, 1999). Our results support the concept that the noradrenergic system influences the acquisition of auditory cued memory while alternatively influencing the consolidation of contextual memory.

Attempts have been made to enhance memory by modulating the adrenergic system to affect memory with most of the attention focused on the use α_2 antagonists, such as yohimbine (O'Carroll *et al*, 1999), idazoxan (Coull *et al*, 1996), and dexefaroxan (Chopin *et al*, 2002). Unfortunately, yohimbine has been reported to increase anxiety in PTSD patients (Southwick *et al*, 1999b) and to enhance fear memory when infused into rat amygdala (Ferry *et al*, 1999a). On the other hand, amnesia to stressful situations such as surgical procedures is a valuable effect of anesthetic agents and many anesthetic agents such as halothane (Rosman *et al*, 1992), propofol (Pang *et al*, 1993), midazolam (Barros *et al*, 1998), isoflurane (Dutton *et al*, 2002), and thiopental do cause anterograde but not retrograde amnesia (Pandit, 1992). However, α_2 agonists appear to behave differently in that they preferentially suppress discrete fear memory over contextual memory. Therefore, α_2 agonists such as dexmedetomidine may reduce anxiety and the responses to the memories of a stressful situation. This may be one of the additional benefits of dexmedetomidine administration in the ICU environment.

There is mounting evidence that the noradrenergic system is dysregulated in post-traumatic stress disorder (PTSD), where traumatic events are vividly recalled with concomitant emotional and autonomic responses (Southwick *et al*, 1999a). The ability of α_2 adrenergic agonists to limit the production of transcription factors generated in

response to memory retrieval indicates that they may be effective in breaking the vicious cycle of PTSD. Clearly, the ability of α_2 agonists to curb noradrenergic transmission may ultimately be useful in many clinical settings where fear memory is detrimental to the health of the patient.

ACKNOWLEDGEMENTS

We acknowledge the National Institutes of Health, USA and the Department of Veterans Affairs.

REFERENCES

- Abercrombie ED, Jacobs BL (1987). Single-unit response of noradrenergic neurons in the locus coeruleus of freely moving cats. I. Acutely presented stressful and nonstressful stimuli. *J Neurosci* 7: 2837-2843.
- Altman JD, Trendelenburg AU, MacMillan L, Bernstein D, Limbird L, Starke K *et al* (1999). Abnormal regulation of the sympathetic nervous system in α_{2A} -adrenergic receptor knockout mice. *Mol Pharmacol* 56: 154-161.
- Amorapanth P, LeDoux JE, Nader K (2000). Different lateral amygdala outputs mediate reactions and actions elicited by a fear-arousing stimulus. *Nat Neurosci* 3: 74-79.
- Anagnostaras SG, Gale GD, Fanselow MS (2001). Hippocampus and contextual fear conditioning: recent controversies and advances. *Hippocampus* 11: 8-17.
- Asan E (1998). The catecholaminergic innervation of the rat amygdala. *Adv Anat Embryol Cell Biol* 142: 1-118.
- Barros DM, Izquierdo LA, Quevedo J, Rodrigues C, Madruga M, Medina JH *et al* (1998). Interaction between midazolam-induced anterograde amnesia and memory enhancement by treatments given hours later in hippocampus, entorhinal cortex or posterior parietal cortex. *Behav Pharmacol* 9: 163-167.
- Blaxall HS, Cerutis DR, Hass NA, Iversen LJ, Bylund DB (1994). Cloning and expression of the α_{2C} -adrenergic receptor from the OK cell line. *Mol Pharmacol* 45: 176-181.
- Bucheler MM, Hadamek K, Hein L (2002). Two α_2 adrenergic receptor subtypes, α_{2A} and α_{2C} , inhibit transmitter release in the brain of gene-targeted mice. *Neuroscience* 109: 819-826.
- Cahill L (2000). Modulation of long-term memory storage in humans by emotional arousal: adrenergic activation and the amygdala. In Aggleton JP (eds). *The Amygdala: A Functional Analysis* 2nd edn. Oxford University Press: New York. pp 426-445.
- Cahill L, McGaugh JL (1998). Mechanisms of emotional arousal and lasting declarative memory. *Trends Neurosci* 21: 294-299.
- Chopin P, Colpaert FC, Marien M (2002). Effects of acute and subchronic administration of dexefaroxan, an α_2 adrenoceptor antagonist, on memory performance in young adult and aged rodents. *J Pharmacol Exp Ther* 301: 187-196.
- Coull JT, Sahakian BJ, Hodges JR (1996). The α_2 antagonist idazoxan remedies certain attentional and executive dysfunction in patients with dementia of frontal type. *Psychopharmacology (Berlin)* 123: 239-249.
- Davies MF, Tsui J, Flannery J, DeLorey TM, Li X, Hoffman BB (2003). Anatomical enlargement augmentation of the noradrenergic system in α_{2A} adrenergic receptor deficient mice is associated with enhanced fear memory. *Brain Res* 986: 157-165.
- Dragunow M, Abraham WC, Goulding M, Mason SE, Robertson HA, Faull RL (1989). Long-term potentiation and the induction of c-fos mRNA and proteins in the dentate gyrus of unanesthetized rats. *Neurosci Lett* 101: 274-280.
- Dunwiddie TV, Roberson NL, Worth T (1982). Modulation of long-term potentiation: effects of adrenergic and neuroleptic drugs. *Pharmacol Biochem Behav* 17: 1257-1264.

- Dutton RC, Maurer AJ, Sonner JM, Fanselow MS, Laster MJ, Eger II EI (2002). Isoflurane causes anterograde but not retrograde amnesia for Pavlovian fear conditioning. *Anesthesiology* **96**: 1223–1229.
- Fanselow MS, Bolles RC (1979). Naloxone and shock-elicited freezing in the rat. *J Comp Physiol Psychol* **93**: 736–744.
- Fanselow MS, Kim JJ (1994). Acquisition of contextual Pavlovian fear conditioning is blocked by application of an NMDA receptor antagonist D,L-2-amino-5-phosphonovaleric acid to the basolateral amygdala. *Behav Neurosci* **108**: 210–212.
- Fendt M, Koch M, Schnitzler HU (1994). Amygdaloid noradrenaline is involved in the sensitization of the acoustic startle response in rats. *Pharmacol Biochem Behav* **48**: 307–314.
- Ferry B, McGaugh JL (1999). Clenbuterol administration into the basolateral amygdala post-training enhances retention in an inhibitory avoidance task. *Neurobiol Learn Mem* **72**: 8–12.
- Ferry B, Roozendaal B, McGaugh JL (1999a). Involvement of α_1 -adrenoceptors in the basolateral amygdala in modulation of memory storage. *Eur J Pharmacol* **372**: 9–16.
- Ferry B, Roozendaal B, McGaugh JL (1999b). Role of norepinephrine in mediating stress hormone regulation of long-term memory storage: a critical involvement of the amygdala. *Biol Psychiatry* **46**: 1140–1152.
- Flordellis CS, Handy DE, Bresnahan MR, Zannis VI, Gavras H (1991). Cloning and expression of a rat brain α_{2B} adrenergic receptor. *Proc Natl Acad Sci USA* **88**: 1019–1023.
- Guo TZ, Jiang JY, Buttermann AE, Maze M (1996). Dexmedetomidine injection into the locus coeruleus produces antinociception. *Anesthesiology* **84**: 873–881.
- Hall J, Thomas KL, Everitt BJ (2001). Fear memory retrieval induces CREB phosphorylation and Fos expression within the amygdala. *Eur J Neurosci* **13**: 1453–1458.
- Hatfield T, McGaugh JL (1999). Norepinephrine infused into the basolateral amygdala posttraining enhances retention in a spatial water maze task. *Neurobiol Learn Mem* **71**: 232–239.
- Hayashi Y, Guo TZ, Maze M (1995). Desensitization to the behavioral effects of α_2 -adrenergic agonists in rats. *Anesthesiology* **82**: 954–962.
- Hayashi Y, Guo TZ, Maze M (1996). Hypnotic and analgesic effects of the α_2 -adrenergic agonist dexmedetomidine in morphine-tolerant rats. *Anesth Analg* **83**: 606–610.
- Huang YY, Martin KC, Kandel ER (2000). Both protein kinase A and mitogen-activated protein kinase are required in the amygdala for the macromolecular synthesis-dependent late phase of long-term potentiation. *J Neurosci* **20**: 6317–6325.
- Impey S, Smith DM, Obrietan K, Donahue R, Wade C, Storm DR (1998). Stimulation of cAMP response element (CRE)-mediated transcription during contextual learning. *Nat Neurosci* **1**: 595–601.
- Ishida Y, Hashiguchi H, Takeda R, Ishizuka Y, Mitsuyama Y, Kannan H et al (2002). Conditioned-fear stress increases Fos expression in monoaminergic and GABAergic neurons of the locus coeruleus and dorsal raphe nuclei. *Synapse* **45**: 46–51.
- Josselyn SA, Shi C, Carlezon Jr WA, Neve RL, Nestler EJ, Davis M (2001). Long-term memory is facilitated by cAMP response element-binding protein overexpression in the amygdala. *J Neurosci* **21**: 2404–2412.
- Kang H, Sun LD, Atkins CM, Soderling TR, Wilson MA, Tonegawa S (2001). An important role of neural activity-dependent CaMKIV signaling in the consolidation of long-term memory. *Cell* **106**: 771–783.
- Kobayashi K, Noda Y, Matsushita N, Nishii K, Sawada H, Nagatsu T et al (2000). Modest neuropsychological deficits caused by reduced noradrenaline metabolism in mice heterozygous for a mutated tyrosine hydroxylase gene. *J Neurosci* **20**: 2418–2426.
- Lakhlani PP, MacMillan LB, Guo TZ, McCool BA, Lovinger DM, Maze M et al (1997). Substitution of a mutant α_{2A} -adrenergic receptor via 'hit and run' gene targeting reveals the role of this subtype in sedative, analgesic, and anesthetic-sparing responses *in vivo*. *Proc Natl Acad Sci USA* **94**: 9950–9955.
- LaLumiere RT, Buen TV, McGaugh JL (2003). Post-training intrabasolateral amygdala infusions of norepinephrine enhance consolidation of memory for contextual fear conditioning. *J Neurosci* **23**: 6754–6758.
- Lamprecht R, Hazvi S, Dudai Y (1997). cAMP response element-binding protein in the amygdala is required for long- but not short-term conditioned taste aversion memory. *J Neurosci* **17**: 8443–8450.
- LeDoux JE, Cicchetti P, Xagoraris A, Romanski LM (1990). The lateral amygdaloid nucleus: sensory interface of the amygdala in fear conditioning. *J Neurosci* **10**: 1062–1069.
- LeDoux JE, Iwata J, Cicchetti P, Reis DJ (1988). Different projections of the central amygdaloid nucleus mediate autonomic and behavioral correlates of conditioned fear. *J Neurosci* **8**: 2517–2529.
- Lee A, Wissekerke AE, Rosin DL, Lynch KR (1998). Localization of α_{2C} -adrenergic receptor immunoreactivity in catecholaminergic neurons in the rat central nervous system. *Neuroscience* **84**: 1085–1096.
- Lin RZ, Chen J, Hu ZW, Hoffman BB (1998). Phosphorylation of the cAMP response element-binding protein and activation of transcription by α_1 adrenergic receptors. *J Biol Chem* **273**: 30033–30038.
- MacMillan LB, Hein L, Smith MS, Piascik MT, Limbird LE (1996). Central hypotensive effects of the α_{2A} -adrenergic receptor subtype. *Science* **273**: 801–803.
- MacMillan LB, Lakhlani P, Lovinger D, Limbird LE (1998). Alpha 2-adrenergic receptor subtypes: subtle mutation of the α_{2A} -adrenergic receptor *in vivo* by gene targeting strategies reveals the role of this subtype in multiple physiological settings. *Recent Prog Horm Res* **53**: 25–42.
- Maren S (1999). Neurotoxic basolateral amygdala lesions impair learning and memory but not the performance of conditional fear in rats. *J Neurosci* **19**: 8696–8703.
- Moller C, Bing O, Heilig M (1994). c-fos expression in the amygdala: *in vivo* antisense modulation and role in anxiety. *Cell Mol Neurobiol* **14**: 415–423.
- Mori-Okamoto J, Namii Y, Tatsuno J (1991). Subtypes of adrenergic receptors and intracellular mechanisms involved in modulatory effects of noradrenaline on glutamate. *Brain Res* **539**: 67–75.
- Nader K, Schafe GE, Le Doux JE (2000). Fear memories require protein synthesis in the amygdala for reconsolidation after retrieval. *Nature* **406**: 722–726.
- Nelemans A, den Hertog A (1987). Calcium translocation during activation of α_1 -adrenoceptor and voltage-operated channels in smooth muscle cells. *Eur J Pharmacol* **140**: 39–46.
- O'Carroll RE, Drysdale E, Cahill L, Shajahan P, Ebmeier KP (1999). Stimulation of the noradrenergic system enhances and blockade reduces memory for emotional material in man. *Psychol Med* **29**: 1083–1088.
- Pandit SK (1992). Anterograde, not retrograde, amnesia after thiopental. *Anesthesiology* **77**: 214–215.
- Pang R, Quartermain D, Rosman E, Turndorf H (1993). Effect of propofol on memory in mice. *Pharmacol Biochem Behav* **44**: 145–151.
- Paxinos G, Franklin KBJ (2001). *The Mouse Brain in Stereotaxic Coordinates* 2nd edn Academic Press: San Diego.
- Przybylski J, Roulet P, Sara SJ (1999). Attenuation of emotional and nonemotional memories after their reactivation: role of β adrenergic receptors. *J Neurosci* **19**: 6623–6628.
- Regan JW, Kobilka TS, Yang-Feng TL, Caron MG, Lefkowitz RJ, Kobilka BK (1988). Cloning and expression of a human kidney cDNA for an α_2 -adrenergic receptor subtype. *Proc Natl Acad Sci USA* **85**: 6301–6305.

- Reisine TD, Heisler S, Hook VY, Axelrod J (1983). Activation of β_2 -adrenergic receptors on mouse anterior pituitary tumor cells increases cyclic adenosine 3':5'-monophosphate synthesis and adrenocorticotropin release. *J Neurosci* 3: 725-732.
- Rogan MT, Staubli UV, LeDoux JE (1997). Fear conditioning induces associative long-term potentiation in the amygdala. *Nature* 390: 604-607.
- Romanski LM, Clugnet MC, Bordi F, LeDoux JE (1993). Somatosensory and auditory convergence in the lateral nucleus of the amygdala. *Behav Neurosci* 107: 444-450.
- Rosman E, Quartermain D, Pang R, Turndorf H (1992). Halothane anesthesia causes state-dependent retrieval failure in mice. *Physiol Behav* 52: 449-453.
- Sara SJ (1985). Noradrenergic modulation of selective attention: its role in memory retrieval. *Ann NY Acad Sci* 444: 178-193.
- Sarvey JM, Burgard EC, Decker G (1989). Long-term potentiation: studies in the hippocampal slice. *J Neurosci Methods* 28: 109-124.
- Schafe GE, Nader K, Blair HT, LeDoux JE (2001). Memory consolidation of Pavlovian fear conditioning: a cellular and molecular perspective. *Trends Neurosci* 24: 540-546.
- Schulz B, Fendt M, Schnitzler HU (2002). Clonidine injections into the lateral nucleus of the amygdala block acquisition and expression of fear-potentiated startle. *Eur J Neurosci* 15: 151-157.
- Simpson JN, McGinty JF (1994). Forskolin increases phosphorylated-CREB and fos immunoreactivity in rat striatum. *Neuroreport* 5: 1213-1216.
- Southwick SM, Bremner JD, Rasmusson A, Morgan III CA, Arnsten A, Charney DS (1999a). Role of norepinephrine in the pathophysiology and treatment of posttraumatic stress disorder. *Biol Psychiatry* 46: 1192-1204.
- Southwick SM, Morgan III CA, Charney DS, High JR (1999b). Yohimbine use in a natural setting: effects on posttraumatic stress disorder. *Biol Psychiatry* 46: 442-444.
- Stone EA, Zhang Y, Hiller JM, Simon EJ, Hillman DE (1997). Activation of fos in mouse amygdala by local infusion of norepinephrine or atipamezole. *Brain Res* 778: 1-5.
- Svensson TH (1987). Stress, central neurotransmitters, and the mechanism of action of α_2 -adrenoceptor agonists. *J Cardiovasc Pharmacol* 10(Suppl 12): S88-S92.
- Takuma T, Tajima Y, Ichida T (1997). Regulation of CREB phosphorylation by cAMP and Ca^{2+} in parotid acinar cells. *Biochem Mol Biol Int* 43: 563-570.
- Talley EM, Rosin DL, Lee A, Guyenet PG, Lynch KR (1996). Distribution of α_{2A} -adrenergic receptor-like immunoreactivity in the rat central nervous system. *J Comp Neurol* 372: 111-134.
- Tischmeyer W, Grimm R (1999). Activation of immediate early genes and memory formation. *Cell Mol Life Sci* 55: 564-574.
- Vaandrager AB, de Jonge HR (1996). Signalling by cGMP-dependent protein kinases. *Mol Cell Biochem* 157: 23-30.
- Weisskopf MG, Bauer EP, LeDoux JE (1999). L-type voltage-gated calcium channels mediate NMDA-independent associative long-term potentiation at thalamic input synapses to the amygdala. *J Neurosci* 19: 10512-10519.
- Wilensky AE, Schafe GE, LeDoux JE (1999). Functional inactivation of the amygdala before but not after auditory fear conditioning prevents memory formation. *J Neurosci* 19: RC48.
- Wilensky AE, Schafe GE, LeDoux JE (2000). The amygdala modulates memory consolidation of fear-motivated inhibitory avoidance learning but not classical fear conditioning. *J Neurosci* 20: 7059-7066.
- Williams CL, Men D, Clayton EC (2000). The effects of noradrenergic activation of the nucleus tractus solitarius on memory and in potentiating norepinephrine release in the amygdala. *Behav Neurosci* 114: 1131-1144.
- Zardetto-Smith AM, Gray TS (1990). Organization of peptidergic and catecholaminergic efferents from the nucleus of the solitary tract to the rat amygdala. *Brain Res Bull* 25: 875-887.