

# Regulation of Working Memory by Dopamine D<sub>4</sub> Receptor in Rats

Kehong Zhang<sup>\*,1,2</sup>, Christopher J Grady<sup>1,2</sup>, Evangelia M Tsapakis<sup>1,2</sup>, Susan L Andersen<sup>1,2</sup>, Frank I Tarazi<sup>1,2</sup> and Ross J Baldessarini<sup>1,2</sup>

<sup>1</sup>Mailman Research Center, McLean Division of Massachusetts General Hospital, Belmont, MA, USA; <sup>2</sup>Department of Psychiatry and Neuroscience Program, Harvard Medical School, Boston, MA, USA

Working memory is regulated by neurotransmitters in prefrontal cortex (PFC), including dopamine and norepinephrine. Previous studies of dopamine function in working memory have focused on the D<sub>1</sub> and D<sub>2</sub> receptors, with most evidence suggesting a dominant role for the D<sub>1</sub> receptor. Since the dopamine D<sub>4</sub> receptor is highly expressed in PFC, we hypothesize that it may also contribute to working memory. To test this hypothesis, we examined behavioral effects of L-745,870, a highly selective, centrally active, D<sub>4</sub> antagonist, using a delayed alternation task in rats. Task performance was dose-dependently affected by the D<sub>4</sub> antagonist, depending on individual baseline functional status of working memory. In rats with good baseline performance, the D<sub>4</sub> antagonist had no effects at low doses, whereas high doses disrupted working memory. In rats with poor baseline working memory, the D<sub>4</sub> antagonist significantly improved working memory at low doses, and higher doses were not distinguishable from vehicle controls. Effects of the D<sub>4</sub> antagonist among poor performers were most robust when task demand for working memory was high, with lesser effects at lower demand level, suggesting that such effects were selective for working memory. The present findings indicate a significant role of the D<sub>4</sub> receptor in working memory, and suggest innovative, D<sub>4</sub>-based, treatment of cognitive deficits associated with neuropsychiatric disorders.

*Neuropsychopharmacology* (2004) 29, 1648–1655, advance online publication, 12 May 2004; doi:10.1038/sj.npp.1300491

**Keywords:** cognition; delayed alternation; dopamine; D<sub>4</sub> receptor; L-745,870; working memory

## INTRODUCTION

Working memory is a mechanism that maintains information over a short period of time (Baddeley, 1986). Such information is held 'on-line' temporarily, and used to guide future response selection. Working memory is essential for virtually all complex behaviors. A trivial example is using a telephone number several seconds after it was read from a directory. Without the guidance of working memory, components of higher-order behavior become temporally disconnected from each other. As a component of complex cognitive processes, working memory has become a central construct in cognitive neuroscience.

As with many other forms of memory, working memory is dependent on many interconnected brain regions, but a great deal of research supports the conclusion that prefrontal cortex (PFC) is critically important. Working memory is impaired in human subjects with lesions in the frontal cortex, and particularly the dorsolateral PFC (Freed-

man and Oscar-Berman, 1986; Verin *et al*, 1993). Ablation of PFC in non-human primates leads to poor performance in tasks that require working memory (Battig *et al*, 1960; Goldman *et al*, 1971). Deficits of working memory are also characteristic of rats with lesions to the medial PFC—the rodent equivalent of the dorsolateral PFC in primates (Kolb *et al*, 1974; Larsen and Divac, 1978).

Working memory is regulated by various neurotransmitters found in PFC, particularly dopamine. Lesioning the mesocortical dopamine pathway from the ventral tegmental area to PFC results in impaired working memory in both primates (Brozoski *et al*, 1979) and rats (Bubser and Schmidt, 1990; Simon, 1981). Deficits induced as such can be rescued with the dopamine precursor L-DOPA, or the dopamine agonist apomorphine, to directly implicate dopaminergic mechanisms (Brozoski *et al*, 1979; Stam *et al*, 1989). Recent studies in human subjects have also provided evidence indicating that working memory is modulated by dopaminergic transmission (Mattay *et al*, 2000).

The receptor basis for dopamine action in working memory has been extensively studied with selective ligands. These studies have demonstrated an important role of the D<sub>1</sub>-like (D<sub>1</sub>, D<sub>5</sub>) receptors. Blockade of the D<sub>1</sub>-like receptors in PFC disrupts working memory (Arnsten *et al*, 1994; Sawaguchi and Goldman-Rakic, 1991). In animals with

\*Correspondence: Dr K Zhang, Mailman Research Center, McLean Hospital, 115 Mill Street, Belmont, MA 02478, USA, Tel: +1 617 855 3222, Fax: +1 617 855 3479, E-mail: kz@mclean.harvard.edu  
Received 2 January 2004; revised 30 March 2004; accepted 19 April 2004

Online publication: 20 April 2004 at <http://www.acnp.org/citations/Npp04200404002/default.pdf>

dopamine depletion, occurring either naturally with aging, or induced by reserpine treatment or chronic stress, the D<sub>1</sub> partial agonist SKF-38393 improves working memory (Arnsten *et al*, 1994; Mizoguchi *et al*, 2000). However, an overflow of dopamine activity, either by excessive dopamine release or overstimulation of postsynaptic D<sub>1</sub> receptor, can impair working memory (Arnsten and Goldman-Rakic, 1998; Cai and Arnsten, 1997; Murphy *et al*, 1996; Zahrt *et al*, 1997). These findings suggest that optimal functioning of PFC requires an intermediate level of dopamine input.

Studies of the D<sub>2</sub>-like (D<sub>2</sub>, D<sub>3</sub>, D<sub>4</sub>) receptors have yielded inconsistent, and sometimes conflicting results. Chronic exposure to the D<sub>2</sub>-like receptor antagonist haloperidol results in disrupted working memory (Castner *et al*, 2000). Acute challenge with various D<sub>2</sub>-like receptor antagonists has been reported to impair working memory and delay-specific PFC neuronal activity in some studies (Arnsten and Goldman-Rakic, 1998; Didriksen, 1995; Murphy *et al*, 1996), but not others (Aultman and Moghaddam, 2001; Bushnell and Levin, 1993; Sawaguchi and Goldman-Rakic, 1994; Verma and Moghaddam, 1996; Williams and Goldman-Rakic, 1995). Working memory deficits induced by the noncompetitive NMDA antagonist ketamine (Verma and Moghaddam, 1996), the benzodiazepine inverse agonist FG-7142, or physiological stress (Arnsten and Goldman-Rakic, 1998) are attenuated by D<sub>2</sub>-like receptor antagonists. These observations suggest that, although the D<sub>1</sub> (or D<sub>5</sub>) receptor is a major constituent of working memory, D<sub>2</sub>-like receptors may also participate in its regulation.

Dopamine D<sub>4</sub> receptor is a member of the D<sub>2</sub>-like receptor family with limited anatomical distribution (Van Tol *et al*, 1991). In mammalian species, D<sub>4</sub> receptor is detected mainly in the corticolimbic areas, with particularly high levels in PFC (Oaks *et al*, 2000; Tarazi and Baldessarini, 1999). Because of this unique distribution, we hypothesize that D<sub>4</sub> receptor may play a significant role in working memory. In the present study, we examined behavioral effects of L-745,870, a highly selective D<sub>4</sub> receptor antagonist, using a continuous delayed alternation task, a commonly used paradigm to assess working memory in rat (Dember and Fowler, 1958).

## MATERIALS AND METHODS

### Subjects

Male Sprague–Dawley rats (4–8 months old; Charles River Labs., Wilmington, MA) were housed in groups of 2–3 under a 12-h artificial daylight/dark schedule (on, 0700–1900). A total of 18 rats were included in the experiment. Daily food intake was restricted to approximately 17 g of standard rat chow, and body weight was maintained at 85–90% of free-feeding weight. Water was freely available. Rats were handled extensively before training.

Working memory was assessed using a standard T-maze. The main alley (18 × 60 cm<sup>2</sup>) was separated from a starting box (18 × 24 cm<sup>2</sup>) and two goal arms (13 × 40 cm<sup>2</sup>) by opaque guillotine doors. At 2 cm from the end of each goal arm, a barrier blocked a food reward (1/6 Froot Loop cereal) from view. A large amount of reward was placed outside both goal arms to mask olfactory cues. The maze was located in the same position in a room with several easily

identifiable visual cues, and cleaned with 50% ethanol between each animal.

### Training

In the first 3–5 days, cage-mates were placed on the maze in pairs, and allowed to explore and consume rewards spread in the goal arms. In the following 3–5 days, rats were placed in the maze individually, with rewards behind barriers.

Training sessions were conducted 5 days/week (Monday through Friday). Each session consisted of 11 trials (a forced trial followed by 10 choice trials). In the forced trial, a randomly selected goal arm was blocked by a guillotine door, and a reward was placed in the other arm. A rat was placed in the starting box, and the guillotine door separating the starting box from the main alley was raised immediately. Once the rat entered the open arm, the guillotine door was closed behind it. In choice trials, both arms were accessible, but reward was available only in the arm not entered in the previous trial. Entry into the arm visited in the previous trial was registered as an error of working memory. Rats were allowed 10 s to consume the reward.

Habituation to injection (needle poke, twice/week for 4 weeks) started when performance of the rats in sessions without delay reached ≤1 error/session on 2 consecutive days. During this period, rats were placed in a holding cage for approximately 10 min after the needle poke but prior to the sessions so that they became fully habituated to the procedure. Initially, needle poke significantly decreased the number of correct trials. However, towards the end of the 4-week habituation, needle poke no longer had any effect on performance.

### Testing

Effects of L-745,870 (Merck; Rahway, NJ) were examined with three different delays between trials (0, 30, or 120 s; each tested in a different session), during which rats were placed in a holding cage next to the maze. L-745,870 was administered intraperitoneally (i.p.) at doses of 0 (20%  $\beta$ -hydroxypropyl-cyclodextrin as vehicle control), 15, 50, 150, 500, or 5000  $\mu$ g/kg at 40 min prior to the start of behavioral testing. At this dose range, L-745,870 is centrally active and selective for the D<sub>4</sub> receptor, and has negligible action at other receptors (Patel *et al*, 1997).

Testing was carried out on Tuesdays and Fridays, with at least a 72-h washout period between testing sessions to minimize carry-over artifacts of previous drug administration. Performance was maintained by training sessions that did not involve delays on the remaining weekdays. The dosing sequence was randomized using a balanced Latin square design. All rats received all doses of L-745,870 (and a vehicle) once for each delay condition.

### Data Analysis


Since rats were used repeatedly, data were analyzed using ANOVA of repeated measures with *post hoc* Dunnett's *t*-test. Probability ≤0.05 was the criterion for statistically significant effect. Data are presented as mean ± SEM. Since working memory is dependent on an optimal level of

dopamine activity (see Introduction), we hypothesize that the D<sub>4</sub> antagonist may have different actions depending upon basal dopamine activity, and hence baseline performance. Accordingly, an average-split analysis was carried out. In this analysis, data obtained from rats with good baseline performance (above average) were separated from rats with poor baseline performance (below average). Nonparametric Spearman rank correlation analysis was performed to determine whether effect of D<sub>4</sub> antagonist was associated with individual baseline performance.

## RESULTS

Performance of rats decreased as task demand for working memory increased (Figure 1). The mean number of correct trials was  $9.96 \pm 0.12$ ,  $8.17 \pm 0.22$ , and  $6.69 \pm 0.22$  at 0, 30, and 120 s delay, respectively. In theory, the number of correct trials at chance performance is 5 ( $50\% \times$  total of 10 choice trials). To truly reflect working memory, we designed a working memory index (WMI), calculated as  $[(\text{correct trials} - 5)/5] \times 100$ . Accordingly, WMI was 99.2, 63.4, and 33.8 at 0, 30, and 120 s delay, respectively.

At 0-s delay, performance was not affected at any dose of L-745,870 (Figure 2a). However, at 30 (Figure 2b) and 120 s delay (Figure 2c), performance was significantly improved at doses of 15 and 150  $\mu\text{g/kg}$ , respectively.




**Figure 1** Performance of the rats in delayed alternation task as a function of delay. (\*\*),  $p \leq 0.01$ .  $N = 18$ .

Next, we separated data from rats with relatively good (above average; Figure 3) vs poor (below average; Figure 4) baseline working memory, as determined with data obtained with vehicle injection at 30 and 120 s delays. In rats with good baseline performance, L-745,870 resulted in a significant *decrease* of WMI in sessions with a 30 s delay at 500 and 5000  $\mu\text{g/kg}$ , with no effect at lower doses of 15–150  $\mu\text{g/kg}$  (Figure 3a). In sessions with a 120 s delay, WMI was significantly decreased at 5000  $\mu\text{g/kg}$  (Figure 3b). In rats with relatively poor baseline performance, treatment with L-745,870 at doses of 15 and 50  $\mu\text{g/kg}$  yielded significantly *increased* WMI in sessions with a 30 s delay, whereas performance returned toward control values at higher doses (Figure 4a). In sessions with a 120 s delay, WMI was significantly increased at 15, 50, and 150  $\mu\text{g/kg}$  (Figure 4b).


Improvement in WMI in the rats with poor baseline performance by low doses of L-745,870 also was dependent on the delay. At a 30 s delay, WMI was increased by a maximum of 77% (from  $45.2 \pm 4.8$  to  $80.0 \pm 9.0$ ) at 15  $\mu\text{g/kg}$ , and somewhat less at higher doses (Figure 4a). At a 120 s delay, WMI was increased by up to 3.86-fold (from  $10.0 \pm 2.8$  to  $48.6 \pm 13.0$ ) at 150  $\mu\text{g/kg}$ , with lesser increases at lower or higher doses (Figure 4b). A Spearman correlation analysis indicated that improvement of working memory performance with the D<sub>4</sub> antagonist was significantly inversely correlated with individual baseline performance (Figure 5).

## DISCUSSION


Dopamine D<sub>4</sub> receptor is expressed preferentially in the limbic system and cortical areas, particularly in PFC (Oaks *et al*, 2000; Tarazi and Baldessarini, 1999). Several previous studies have examined cognitive effects of D<sub>4</sub> antagonists. Using an object retrieval/detour task in monkeys (a test of frontostriatal function), Jentsch *et al* (1999) found that behavioral deficits induced by the NMDA antagonist phencyclidine could be reversed with the selective D<sub>4</sub> antagonist NGD94-1. The D<sub>4</sub>-selective antagonist PNU-101,387G has been reported to prevent working memory deficits induced by the benzodiazepine inverse agonist FG-7142 in monkeys (Arnsten *et al*, 2000). These studies implicate D<sub>4</sub> receptor in cognitive functions generally, but



**Figure 2** Effects of L-745,870 on delayed alternation task performance at various delays. (\*),  $p \leq 0.05$ .  $N = 18$ .



**Figure 3** Effects of L-745,870 on working memory in rats with above-average baseline performance. (\*),  $p \leq 0.05$  in comparison to vehicle controls.  $N = 8$  at 30 s, and 9 at 120 s delay.




**Figure 4** Effects of L-745,870 on working memory in rats with below-average baseline performance. (\*),  $p \leq 0.05$  in comparison to vehicle controls.  $N = 10$  at 30 s, and 9 at 120 s delay.

explicit involvement in working memory *per se* was uncertain since the D<sub>4</sub> antagonist may have produced improvements in cognitive function by interacting with the agents used to disrupt working memory.

In the present study, we used a delayed alternation task to test for a role of the D<sub>4</sub> receptor in working memory in rats. Consistent with previous reports (Dember and Fowler, 1958), performance of rats in this task was dependent on the length of the intertrial delay, during which the rats must remember which arm of the T-maze was visited in the preceding trial. The number of correct trials and the WMI decreased significantly with increasing delays between trials, indicating construct validity of the procedure as a paradigm for working memory.

The present results indicate that dopamine regulates working memory by multiple mechanisms that evidently include a D<sub>4</sub> component. Changes of WMI by L-745,870 were not randomly distributed among various doses of L-745,870. Instead, such changes followed particular patterns (inverted U-shape dose-response in poor performers, disruption by high doses in good performers). These observations suggest that the observed drug effects were related to altered levels of D<sub>4</sub> receptor activation.

In addition, we found that effects of the D<sub>4</sub> antagonist were dependent on both individual baseline working memory level and dosage. In rats with above-average baseline performance, low doses of L-745,870 had little effect, but disrupted working memory at higher doses. In rats with below-average baseline performance, L-745,870



**Figure 5** Relationship between baseline performance and effect of 50 µg/kg L-745,870 on working memory.  $N=18$ . Spearman  $r_s=0.59$ ,  $p\leq 0.05$ .

produced an inverted U-shape dose-response. At low doses, L-745,870 (15–150 µg/kg) resulted in a significant improvement of working memory. At higher doses (500 and 5000 µg/kg), working memory returned toward vehicle control levels. These findings suggest that working memory is differentially regulated by the D<sub>4</sub> receptor, depending on baseline working memory function.

Based on the inverted U-shape dose-response found with the D<sub>4</sub> antagonist in rats with relative poor baseline, it is reasonable to conclude that optimal working memory requires an intermediate level of D<sub>4</sub> receptor stimulation. We further speculate that signal transduction mediated by D<sub>4</sub> receptor may be overactive in subjects with poor working memory. Thus, reducing D<sub>4</sub> receptor stimulation with low doses of an antagonist in these rats enhanced working memory.

Enhancement of performance in the delayed alternation task by L-745,870 was dependent on task demand for working memory. At a 0 s delay (minimal demand for working memory), performance of the rats was not affected by any dose of L-745,870. At a 120 s delay (relatively high demand), the WMI in rats with poor baseline performance was increased by up to nearly four-fold by the D<sub>4</sub> antagonist. At an intermediate level of demand for working memory (30 s delay), L-745,870 produced a significant, but somewhat smaller increase in WMI (up to 80%) in rats with poor baseline performance. These findings suggest that the effects of L-745,870 on delayed alternation task performance reflect changes in working memory.

An important distinction between the present experiments and several previous studies that reported behavior effects of L-745,870 (Mansbach *et al*, 1998; Patel *et al*, 1997; Zhang *et al*, 2001, 2002) was the finding of significant effects at much lower doses, with presumably greater selectivity at the D<sub>4</sub> receptor. Although we do not have direct evidence

for the *in vivo* specificity for the D<sub>4</sub> receptor, other studies using surrogate markers such as plasma prolactin level (Patel *et al*, 1997) indicate that L-745,870 is devoid of any noticeable affect on the D<sub>2</sub> receptor at doses used in this study.

Consistent with the view that an optimal level of dopaminergic functioning is required for working memory functioning are human studies indicating that working memory, and function of PFC in general, may depend on individual levels of dopamine signaling. One such observation is the inverse correlation of working memory to the activity of catechol-*O*-methyltransferase (COMT, an enzyme that converts dopamine to the inactive metabolite 3-methoxytyramine), based on inheritance of specific alleles of the COMT gene (Mattay *et al*, 2003). In order to establish a relationship between working memory and the D<sub>4</sub> receptor, further experiments using reliable methods to quantify the density and function of the D<sub>4</sub> receptor are required. These might include specific testing of memory and other cognitive functions with a range of doses of D<sub>4</sub> antagonists that have already been proven to be safe and well tolerated by human subjects, including L-745,870 (Kramer *et al*, 1997).

Specific mechanisms by which D<sub>4</sub> antagonists produce cognitive effects are unknown. We propose that PFC is a leading candidate for such actions since PFC is vitally important for working memory (Goldman-Rakic, 2001) and since D<sub>4</sub> receptor mRNA and proteins are highly expressed in PFC (Oaks *et al*, 2000; Tarazi and Baldessarini, 1999). During working memory tasks, certain groups of pyramidal neurons in PFC increase firing rate after presentation of a spatial cue (for a review, see Goldman-Rakic, 2001). These neurons remain activated in the delay period (during which the spatial cue is removed), and cease firing when a response is executed. Failure of these neurons to maintain activity during the delay period is associated invariably with performance errors. These observations demonstrate that working memory is in part encoded and maintained by pyramidal neurons in PFC. There is evidence that the same neurons are modulated by D<sub>4</sub> receptor. Notably, several immunocytochemical studies indicate that D<sub>4</sub> receptor resides in pyramidal neurons in PFC (Ariano *et al*, 1997; Mrzljak *et al*, 1996; Wedzony *et al*, 2000). The presence of D<sub>4</sub> receptor in pyramidal neurons places this receptor in an ideal location to modulate working memory. Indeed, electrophysiological studies using D<sub>4</sub> selective ligands or knockout mice provided strong evidence that physiological property of PFC pyramidal neurons is regulated by D<sub>4</sub> receptor in normal condition (Rubinstein *et al*, 2001; Wang *et al*, 2002).

By simultaneously recording multiple PFC neurons in behaving monkeys, Goldman-Rakic and her colleagues provided evidence that pyramidal neurons representing working memory inhibit activity of surrounding units via GABAergic interneurons (Constantinidis *et al*, 2002; Rao *et al*, 1999). Through this collateral pathway, GABAergic interneurons sharpen the presentation of working memory. The presence of D<sub>4</sub> receptor in the GABAergic interneurons in PFC (Mrzljak *et al*, 1996; Wedzony *et al*, 2000) raises the possibility that D<sub>4</sub> antagonists may also regulate working memory indirectly through GABAergic interneurons.

A subpopulation of the D<sub>4</sub> receptor is present in presynaptic terminals in nucleus accumbens as evidenced by experiments that combined electronic microscopy with immunocytochemistry (Svingos *et al*, 2000). Although such findings have not been replicated in PFC, dopamine release in PFC has been shown to be increased in D<sub>4</sub> receptor knockout mice (Rubinstein *et al*, 1997), and in response to D<sub>4</sub> antagonists (Broderick and Piercey, 1998; Millan *et al*, 1998). These observations suggest that D<sub>4</sub> antagonist may affect working memory by modulating dopamine release in PFC.

Lesions to hippocampus (Aggleton *et al*, 1986), and closely associated structures, including the fornix (Pisa, 1981) and the septal area (Brito and Thomas, 1981), also produce a deficit in working memory as measured using delayed alternation. Significant levels of the D<sub>4</sub> receptor in hippocampus (Oaks *et al*, 2000; Tarazi and Baldessarini, 1999) raise the possibility that behavioral effects of L-745,870 observed in the present study may include actions in the hippocampal complex. Adding one more dimension of complexity, the D<sub>4</sub> receptor has high affinity for both dopamine and norepinephrine (Lanau *et al*, 1997). Since hippocampus as well as PFC is innervated by both catecholamines, it is conceivable that the behavioral effects of the D<sub>4</sub> antagonist observed in this study may reflect combined effects of both catecholamines in both brain regions.

Finally, the present findings suggest that the D<sub>4</sub> receptor might play a role in neuropsychiatric disorders involving working memory deficits, such as schizophrenia and attention-deficit/hyperactivity disorder (ADHD). D<sub>4</sub> receptor was initially implicated in schizophrenia by the relatively high D<sub>4</sub> receptor affinity for clozapine, an atypical antipsychotic agent (Oaks *et al*, 2000; Tarazi and Baldessarini, 1999). Subsequent clinical trials with D<sub>4</sub> receptor antagonists have failed to show antipsychotic efficacy of these agents (Corrigan *et al*, 2004; Kramer *et al*, 1997). However, possible effects of such treatment on cognitive deficits might have been overlooked since these studies were conducted in acutely ill psychotic patients, using clinical symptom rating scales. Further studies on chronically ill schizophrenia patients, aimed at quantitative assessments of specific cognitive functions and social functioning, are needed to adequately evaluate D<sub>4</sub> receptor as a potentially important therapeutic target. Genetic studies have strongly implicated D<sub>4</sub> receptor polymorphism in ADHD (La Hoste *et al*, 1996; Faraone *et al*, 2001). Using young rats with neonatal 6-hydroxydopamine lesions as a model for ADHD, we reported that locomotor hyperactivity in this model was inhibited by several selective D<sub>4</sub> antagonists as by the stimulant methylphenidate, but not antagonists that bind preferentially to the D<sub>2</sub> or 5-HT<sub>2A</sub> receptor (Davids *et al*, 2002; Zhang *et al*, 2001, 2002). Given a hypothesized role of working memory deficits in ADHD (Pennington and Ozonoff, 1996; Denney and Rapport, 2001), beneficial effects of L-745,870 on cognition identified in the current study encourage further consideration of D<sub>4</sub> antagonists as novel treatments for clinical ADHD and other disorders of attention and cognition.

## ACKNOWLEDGEMENTS

This work was supported, in part, by a grant from the Bruce J Anderson Foundation and by the McLean Private Donors Neuropsychopharmacology Research Fund (to RJB). L-745,870 was generously donated by Merck (Rahway, NJ).

## REFERENCES

- Aggleton JP, Hunt PR, Rawlins JNP (1986). The effects of hippocampal lesions upon spatial and non-spatial tests of working memory. *Behav Brain Res* 19: 133–146.
- Ariano MA, Wang J, Noblett KL, Larson ER, Sibley DR (1997). Cellular distribution of the rat D<sub>4</sub> dopamine receptor protein in the CNS using anti-receptor antisera. *Brain Res* 752: 26–34.
- Arnsten AFT, Cai JX, Murphy BL, Goldman-Rakic PS (1994). Dopamine D<sub>1</sub> receptor mechanisms in the cognitive performance of young adult and aged monkeys. *Psychopharmacology* 116: 143–151.
- Arnsten AFT, Goldman-Rakic PS (1998). Noise stress impairs prefrontal cortical cognitive function in monkeys: evidence for a hyperdopaminergic mechanism. *Arch Gen Psychiatry* 55: 362–368.
- Arnsten AFT, Murphy B, Merchant K (2000). The selective dopamine D<sub>4</sub> receptor antagonist, PNU-101387G, prevents stress-induced cognitive deficits in monkeys. *Neuropsychopharmacology* 4: 405–410.
- Aultman JM, Moghaddam B (2001). Distinct contributions of glutamate and dopamine receptors to temporal aspects of rodent working memory using a clinically relevant task. *Psychopharmacology* 15: 353–364.
- Baddeley AD (1986). *Working Memory*. Clarendon: Oxford.
- Battig K, Rosvold HE, Mishkin M (1960). Comparison of the effects of frontal and caudate lesions on delayed response and alternation in monkeys. *J Comp Physiol Psychol* 53: 400–404.
- Brito GNO, Thomas GJ (1981). T-maze alternation, response patterning, and septo-hippocampal circuitry in rats. *Behav Brain Res* 3: 319–340.
- Broderick PA, Piercey MF (1998). Clozapine, haloperidol, and the D<sub>4</sub> antagonist PNU-101387G: *in vivo* effects on mesocortical, mesolimbic, and nigrostriatal dopamine and serotonin release. *J Neural Transm* 105: 749–767.
- Brozoski TJ, Brown RM, Rosvold HE, Goldman PS (1979). Cognitive deficit caused by regional depletion of dopamine in prefrontal cortex of rhesus monkey. *Science* 205: 929–932.
- Bubser M, Schmidt WJ (1990). 6-Hydroxydopamine lesion of the rat prefrontal cortex increases locomotor activity, impairs acquisition of delayed alternation tasks, but does not affect uninterrupted tasks in the radial maze. *Behav Brain Res* 37: 157–168.
- Bushnell PJ, Levin ED (1993). Effects of dopaminergic drugs on working and reference memory in rats. *Pharmacol Biochem Behav* 45: 765–776.
- Cai JX, Arnsten AFT (1997). Dose-dependent effects of the dopamine D<sub>1</sub> receptor agonists A77636 or SKF81297 on spatial working memory in aged monkeys. *J Pharmacol Exp Ther* 283: 183–189.
- Castner SA, Williams GV, Goldman-Rakic PS (2000). Reversal of antipsychotic-induced working memory deficits by short-term dopamine D<sub>1</sub> receptor stimulation. *Science* 287: 2020–2022.
- Constantinidis C, Williams GV, Goldman-Rakic PS (2002). A role for inhibition in shaping the temporal flow of information in prefrontal cortex. *Nat Neurosci* 5: 175–180.
- Corrigan MH, Gallen CC, Bonura ML, Merchant KM (2004). Effectiveness of the selective D<sub>4</sub> antagonist sonepipirazole in schizophrenia: a placebo-controlled trial. *Biol Psychiatry* 55: 445–451.

- Davids E, Zhang K, Baldessarini RJ (2002). Stereoselective effects of methylphenidate on hyperactivity in juvenile rats induced by neonatal 6-hydroxydopamine lesioning. *Psychopharmacology* 160: 92–98.
- Dember WN, Fowler H (1958). Spontaneous alternation behavior. *Psychol Bull* 55: 412–428.
- Denney CB, Rapport MD (2001). Cognitive pharmacology of stimulants in children with ADHD. In: Solanto MV, Arnsten AFT, Castellanos FX (eds). *Stimulant Drugs and ADHD*. Oxford University Press: New York. pp 283–302.
- Didriksen M (1995). Effects of antipsychotics on cognitive behaviour in rats using the delayed non-match to position paradigm. *Eur J Pharmacol* 281: 241–250.
- Faraone SV, Doyle AE, Mick E, Biederman J (2001). Meta-analysis of the association between the 7-repeat allele of the dopamine D<sub>4</sub> receptor gene and attention deficit hyperactivity disorder. *Am J Psychiatry* 158: 1052–1057.
- Freedman M, Oscar-Berman M (1986). Bilateral frontal lobe disease and selective delayed response deficits in humans. *Behav Neurosci* 100: 337–342.
- Goldman PS, Rosvold HE, Vest B, Galkin TW (1971). Analysis of the delayed-alternation deficit produced by dorsolateral prefrontal lesions in the rhesus monkey. *J Comp Physiol Psychol* 77: 212–220.
- Goldman-Rakic PS (2001). Working memory dysfunction in schizophrenia. In: Salloway SP, Malloy PF, Duffy JD (eds). *The Frontal Lobes and Neuropsychiatric Illness*. American Psychiatric Publishing: Washington, DC. pp 71–82.
- Jentsch JD, Taylor JR, Redmond Jr ED, Elsworth JD, Youngren KD, Roth RH (1999). Dopamine D<sub>4</sub> receptor antagonist reversal of subchronic phencyclidine-induced object retrieval/detour deficits in monkeys. *Psychopharmacology* 142: 78–84.
- Kolb B, Nonneman AJ, Singh RK (1974). Double dissociation of spatial impairments and perseveration following selective prefrontal lesions in rats. *J Comp Physiol Psychol* 87: 772–780.
- Kramer MS, Last B, Geston A, Reines SA (1997). The effects of a selective D<sub>4</sub> dopamine receptor antagonist (L-745,870) in acutely psychotic inpatients with schizophrenia. *Arch Gen Psychiatry* 54: 567–572.
- La Hoste GJ, Swanson JM, Wigal SB, Glabe C, Wigal T, King N, Kennedy JL (1996). Dopamine D<sub>4</sub> receptor gene polymorphism is associated with attention deficit hyperactivity disorder. *Mol Psychiatry* 1: 128–131.
- Lanau F, Zenner MT, Civelli O, Hartman DS (1997). Epinephrine and norepinephrine act as potent agonists at the recombinant human dopamine D<sub>4</sub> receptor. *J Neurochem* 68: 804–812.
- Larsen JK, Divac I (1978). Selective ablations within the prefrontal cortex of the rat and performance of delayed alternation. *Physiol Psychol* 6: 15–17.
- Mansbach RS, Brooks EW, Sanner MA, Zorn SH (1998). Selective dopamine D<sub>4</sub> receptor antagonists reverse apomorphine-induced blockade of prepulse inhibition. *Psychopharmacology* 135: 194–200.
- Mattay VS, Callicott JH, Bertolino A, Heaton I, Frank JA, Coppola R et al (2000). Effects of dextroamphetamine on cognitive performance and cortical activation. *Neuroimage* 12: 268–275.
- Mattay VS, Goldberg TE, Fera F, Hariri AR, Tessitore A, Egan MF et al (2003). Catechol O-methyltransferase val158-met genotype and individual variation in the brain response to amphetamine. *Proc Natl Acad Sci USA* 100: 6186–6191.
- Millan MJ, Newman-Tancredi A, Brocco M, Gobert A, Lejeune F, Audinot A et al (1998). S 18126 (2-[4-(2,3-dihydrobenzo[1,4]-dioxin-6-yl)piperazin-1-yl methyl]indan-2-yl), a potent, selective and competitive antagonist at dopamine D<sub>4</sub> receptors: an *in vitro* and *in vivo* comparison with L 745,870 (3-(4-[4-chlorophenyl]-piperazin-1-yl)methyl-1H-pyrrolo[2,3b]pyridine) and raclopride. *J Pharmacol Exp Ther* 287: 167–186.
- Mizoguchi K, Yuzurihara M, Ishige A, Sasaki H, Chui D, Tabira T (2000). Chronic stress induces impairment of spatial working memory because of prefrontal dopaminergic dysfunction. *J Neurosci* 20: 1568–1574.
- Mrzljak L, Bergson C, Pappy M, Huff R, Levenson R, Goldman-Rakic PS (1996). Localization of dopamine D<sub>4</sub> receptors in GABAergic neurons of the primate brain. *Nature* 381: 245–248.
- Murphy BL, Arnsten ATF, Goldman-Rakic PS, Roth RH (1996). Increased dopamine turnover in the prefrontal cortex impairs spatial working memory performance in rats and monkeys. *Proc Natl Acad Sci USA* 93: 1325–1329.
- Oaks JN, Oldenhof J, Van Tol HHM (2000). The dopamine D<sub>4</sub> receptor: one decade of research. *Eur J Pharmacol* 405: 303–327.
- Patel S, Freedman S, Chapman KL, Emms F, Flectcher AE, Knowles M et al (1997). Biological profile of L-745,870, a selective antagonist with high affinity for the dopamine D<sub>4</sub> receptor. *J Pharmacol Exp Ther* 283: 636–647.
- Pennington BF, Ozonoff S (1996). Executive functions and developmental psychopathology. *J Child Psychol Psychiat* 37: 51–87.
- Pisa M (1981). Amnesia for discrete events in rats with fornixotomies: effects of interference on spatial alternation performance. *J Comp Physiol Psychol* 95: 924–942.
- Rao SG, Williams GV, Goldman-Rakic PS (1999). Isodirectional tuning of adjacent interneurons and pyramidal cells during working memory: evidence for microcolumnar organization in PFC. *J Neurophysiol* 81: 1903–1916.
- Rubinstein M, Cepeda C, Hurst RS, Flores-Hernandez J, Ariano MA, Falzone T et al (2001). Dopamine D<sub>4</sub> receptor-deficient mice display cortical hyperexcitability. *J Neurosci* 21: 3756–3763.
- Rubinstein M, Phillips TJ, Bunzow JR, Falzone TL, Dziewczapolski G, Zhang G et al (1997). Mice lacking dopamine D<sub>4</sub> receptors are supersensitive to ethanol, cocaine, and methamphetamine. *Cell* 90: 991–1001.
- Sawaguchi T, Goldman-Rakic PS (1991). D<sub>1</sub> dopamine receptors in prefrontal cortex: involvement in working memory. *Science* 251: 947–950.
- Sawaguchi T, Goldman-Rakic PS (1994). The role of D<sub>1</sub>-dopamine receptor in working memory: local injections of dopamine antagonists into the prefrontal cortex of rhesus monkeys performing an oculomotor delayed-response task. *J Neurophysiol* 71: 515–528.
- Simon H (1981). Dopaminergic A10 neurons and frontal system. *J Physiol* 77: 81–95.
- Stam CJ, de Bruin JP, van Haelst AM, van der Gugten J, Kalsbeek A (1989). Influence of the mesocortical dopaminergic system on activity, food hoarding, social-agonistic behavior, and spatial delayed alternation in male rats. *Behav Neurosci* 103: 24–35.
- Svingos AL, Periasamy S, Pickel VM (2000). Presynaptic dopamine D<sub>4</sub> receptor localization in the rat nucleus accumbens. *Synapse* 36: 222–232.
- Tarazi FI, Baldessarini RJ (1999). Dopamine D<sub>4</sub> receptors: significance for molecular psychiatry at the millennium. *Mol Psychiatry* 4: 529–538.
- Van Tol HHM, Bunzow JR, Guan H-C, Sunahara RK, Seeman P, Niznik HB et al (1991). Cloning of a human dopamine D<sub>4</sub> receptor gene with high affinity for the antipsychotic clozapine. *Nature* 350: 614–619.
- Verin M, Partoit A, Pillon B, Malapani C, Agid Y, Dubois B (1993). Delayed response tasks and prefrontal lesions in man—evidence for self-generated patterns of behavior with poor environmental modulation. *Neuropsychologia* 31: 1279–1296.
- Verma A, Moghaddam B (1996). NMDA receptor antagonists impair prefrontal cortex function as assessed via spatial delayed

- alternation performance in rats: modulation by dopamine. *J Neurosci* **16**: 373–379.
- Wang X, Zhong P, Yan Z (2002). Dopamine D<sub>4</sub> receptor modulate GABAergic signaling in pyramidal neurons of prefrontal cortex. *J Neurosci* **22**: 9185–9193.
- Wedzony K, Chocyk A, Mackowiak M, Fijał K, Czyrak A (2000). Cortical localization of dopamine D<sub>4</sub> receptors in the rat brain-immunocytochemical study. *J Physiol Pharmacol* **51**: 205–221.
- Williams GV, Goldman-Rakic PS (1995). Modulation of memory fields by dopamine D<sub>1</sub> receptors in prefrontal cortex. *Nature* **37**: 572–575.
- Zahrt J, Taylor JR, Mathew RG, Arnsten AF (1997). Supranormal stimulation of D<sub>1</sub> dopamine receptors in the rodent prefrontal cortex impairs spatial working memory performance. *J Neurosci* **17**: 8528–8535.
- Zhang K, Davids E, Tarazi FI, Baldessarini RJ (2002). Effects of dopamine D<sub>4</sub> receptor-selective antagonists on motor hyperactivity in rats with neonatal 6-hydroxydopamine lesions. *Psychopharmacology* **161**: 100–106.
- Zhang K, Tarazi FI, Baldessarini RJ (2001). Role of dopamine D<sub>4</sub> receptors in motor hyperactivity induced by neonatal 6-hydroxydopamine lesions in rats. *Neuropsychopharmacology* **25**: 624–632.