

## Letter to the Editor

# Clinical Implications of Cocaine-Induced Cortical Depression

Charles A Dackis<sup>\*1</sup> and Charles P O'Brien<sup>1,2</sup>

<sup>1</sup>University of Pennsylvania School of Medicine, Treatment Research Center, Philadelphia, PA, USA; <sup>2</sup>Department of Veterans Affairs Medical Center, Philadelphia, PA, USA

Neuropsychopharmacology (2005) 30, 1033. doi:10.1038/sj.npp.1300701

Sir

(Trantham-Davidson and Lavin, 2004) found that the acute administration of cocaine decreased spontaneous firing and VTA-evoked responses in the prefrontal cortex (PFC), and compromised the membrane bistability of PFC pyramidal neurons. The authors previously reported similar findings after repeated cocaine administration, and should be commended for their pursuit of cocaine-induced neuroadaptations through *in vivo* single-cell recording studies. In other animal studies, chronic exposure to cocaine also produces dysfunctional morphological changes (Robinson *et al*, 2001) and reduced numbers (Lidow and Song, 2001) of PFC pyramidal neurons, which is consistent with reports (reviewed elsewhere) that cocaine-dependent patients have reductions in PFC metabolism and gray matter density, and perform poorly on standardized tests that assess frontal function (Dackis and O'Brien, 2003). While the authors reasonably concluded that electrophysiological alterations of pyramidal neurons might affect attention, working memory, and information processing, we believe that PFC dysfunction may also be a core component of cocaine addiction, and contribute to many baffling characteristics of addicted patients that were once thought to be purely psychological.

Denial, a hallmark of cocaine addiction, classically involves minimization, rationalization, and poor insight into cocaine-related hazards. These examples of nonadaptive executive function could stem from cocaine-induced PFC imbalances, as might problems with decision-making, impulse control, and motivation that are characteristic of cocaine-addicted patients. Also, it has been firmly established by a large number of controlled studies that regions of the PFC (especially the anterior cingulate cortex) are robustly activated during cue-induced cocaine craving. Remarkably, these hypermetabolic responses occur in regions that are hypometabolic at baseline, creating an enhanced 'delta' response that might underlie cue salience,

an important yet poorly understood phenomenon that often leads to recidivism in the clinical arena. Pharmacological interventions that diminish the magnitude of this metabolic response, either by normalizing hypofrontality (eg glutamate-enhancing agents like modafinil) or dampening hypermetabolic responses to cocaine-conditioned cues (eg GABA-enhancing agents like topiramate or baclofen), are currently under intense investigation (Dackis, 2004). Human and animal studies also suggest that delta responses are important in cocaine reward. Patients with baseline negative affective symptoms (Newton *et al*, 2003), or baseline cocaine withdrawal (Sofuoglu *et al*, 2003), report enhanced cocaine-induced euphoria under controlled conditions. Additionally, cocaine reinstatement is obliterated in animals after baseline glutamate depletion has been pharmacologically normalized, diminishing the delta release (peak minus baseline) of glutamate after cocaine administration (Baker *et al*, 2003). Given these considerations, the evolving controversy over glutamate-enhancing vs GABA-enhancing agents for cocaine dependence could ultimately resolve in relative agreement.

## REFERENCES

- Baker DA, McFarland K, Lake RW, Shen H, Toda S, Kalivas PW (2003). N-acetyl cysteine-induced blockade of cocaine-induced reinstatement. *Ann NY Acad Sci* 1003: 349–351.
- Dackis C, O'Brien C (2003). Glutamatergic agents for cocaine dependence. *Ann NY Acad Sci* 1003: 328–345.
- Dackis CA (2004). Recent advances in the pharmacotherapy of cocaine dependence. *Curr Psychiatry Rep* 6: 323–331.
- Lidow MS, Song ZM (2001). Primates exposed to cocaine *in utero* display reduced density and number of cerebral cortical neurons. *J Comp Neurol* 435: 263–275.
- Newton TF, Kalechstein AD, Tervo KE, Ling W (2003). Irritability following abstinence from cocaine predicts euphoric effects of cocaine administration. *Addict Behav* 28: 817–821.
- Robinson TE, Gorny G, Mitton E, Kolb B (2001). Cocaine self-administration alters the morphology of dendrites and dendritic spines in the nucleus accumbens and neocortex. *Synapse* 39: 257–266.
- Sofuoglu M, Dudish-Poulsen S, Brown S, Hatsukami K (2003). Association of cocaine withdrawal symptoms with more severe dependence and enhanced subjective response to cocaine. *Drug Alcohol Depend* 69: 273–282.
- Trantham-Davidson H, Lavin A (2004). Acute cocaine administration depresses cortical activity. *Neuropsychopharmacology* 29: 2046–2051.

\*Correspondence: Dr CA Dackis, University of Pennsylvania School of Medicine, Treatment Research Center, 3900 Chestnut Street, Philadelphia, PA 19104, USA; Tel: +1 215 662 8752; Fax: +1 215 243 4665; E-mail: dackis@mail.med.upenn.edu

Received 18 November 2004; revised 11 January 2005; accepted 16 January 2005

Online publication: 19 January 2005 at <http://www.acnp.org/citations/Npp011905040542/default.pdf>