

Evidence for the Preferential Involvement of 5-HT_{2A} Serotonin Receptors in Stress- and Drug-Induced Dopamine Release in the Rat Medial Prefrontal Cortex

Elizabeth A Pehek^{*1,2,3}, Christine Nocjar^{1,3}, Bryan L Roth^{1,2,4}, Tara A Byrd³ and Omar S Mabrouk^{1,3}

¹Department of Psychiatry, Case Western Reserve School of Medicine, Cleveland, OH, USA; ²Department of Neurosciences, Case Western Reserve School of Medicine, Cleveland, OH, USA; ³Louis Stokes Cleveland VA Medical Center, Cleveland, OH, USA; ⁴Department of Biochemistry, Case Western Reserve School of Medicine, Cleveland, OH, USA

The mechanism(s) by which serotonin modulates dopamine release in the medial prefrontal cortex is not known, although studies suggest an involvement of 5-HT₂ family receptors. We employed *in vivo* microdialysis and putatively selective 5-HT_{2A} antagonists (M100907, MDL 11,939, SR46349B) to determine if 5-HT_{2A} receptors are responsible for both drug- and stress-induced DA release in the medial prefrontal cortex. MDL 11,939 and SR46349B receptor-binding studies indicated, for the first time, that only MDL 11,939 had greater selectivity for the 5-HT_{2A} vs the 5-HT_{2C} receptor subtypes similar to M100907, and that both showed low or no affinity for non-5-HT₂ receptors. Reverse dialysis with 5-HT_{2A} antagonists had little or no effect on basal dopamine efflux. However, intracortical administration of MDL 11,939 or M100907 attenuated dopamine release induced by systemic administration of the 5-HT₂ agonist DOI. Dual-probe microdialysis demonstrated that systemic DOI also increased glutamate concentrations in the ventral tegmental area (VTA). This was blocked by intracortical M100907. Cortical perfusion with M100907, or the atypical antipsychotic drug risperidone, but not the 5-HT_{2B/C} ligand SB 206553, also decreased dopamine release induced physiologically by stress. These results indicate that stimulation of cortical 5-HT_{2A} receptors increases the release of dopamine from the mesocortical system. They suggest that this effect may be mediated by increases in glutamate release from corticotegmental projections to the VTA. Additionally, they indicate that cortical 5-HT_{2A} receptors modulate evoked dopamine release, such as that observed physiologically following mild stress. These findings may have implications for the pharmacological treatment of disorders resulting from or exacerbated by stress. *Neuropsychopharmacology* (2006) 31, 265–277. doi:10.1038/sj.npp.1300819; published online 6 July 2005

Keywords: DOI; glutamate; M100907; medial prefrontal cortex; schizophrenia; ventral tegmental area

INTRODUCTION

A common property of atypical antipsychotic drugs such as clozapine or risperidone is high affinity for the serotonin-2A (5-HT_{2A}) receptor subtype. However, it is clear that atypical antipsychotic drugs have appreciable affinities for most biogenic amine receptors (Roth *et al*, 2004). The apparent preferentially high affinity for 5-HT_{2A} receptors led to the hypothesis that antipsychotic efficacy may be related to the ability of a drug to antagonize 5-HT_{2A} receptors (Meltzer, 1989). Studies show that atypical antipsychotic drugs regulate dopamine (DA) release in the

medial prefrontal cortex (mPFC) (eg Moghaddam and Bunney, 1990; Pehek and Yamamoto, 1994). Since abnormalities in the mesocortical DA system have been associated with the negative symptoms of schizophrenia (Weinberger, 1987), it is possible that a causal link between 5-HT_{2A} receptor antagonism, alterations in mesocortical DA release, and clinical efficacy exists.

Neurons in the mesocortical DA system originate in the ventral tegmental area (VTA) and project to the mPFC. Both the mPFC and VTA are innervated by serotonergic neurons originating in the dorsal raphe (Azmitia and Segal, 1978; Beart and McDonald, 1982). Immunohistochemical studies indicate the presence of 5-HT_{2A} receptors in both the PFC (Willins *et al*, 1997; Jakab and Goldman-Rakic, 1998; Miner *et al*, 2003) and VTA (Doherty and Pickel, 2000; Nocjar *et al*, 2002) where they could, at least in theory, modulate DA neuronal activity. In the mPFC, the vast majority of 5-HT_{2A} receptors are localized to the apical dendrites of pyramidal cells and, to a lesser extent, GABAergic interneurons (Willins *et al*, 1997; Jakab and Goldman-Rakic, 1998; Miner *et al*, 2003). Thus, regulation of

*Correspondence: Dr EA Pehek, Departments of Psychiatry and Neurosciences, Case Western Reserve University School of Medicine, VA Medical Center 151 (VV), 10701 East Blvd., Room K219, Cleveland, OH 44106, USA, Tel: +216 791 3800 ext 4237, Fax: +216 229 8509, E-mail: eap6@cwru.edu

Received 9 February 2005; revised 29 April 2005; accepted 2 May 2005

Online publication: 3 June 2005 at <http://www.acnp.org/citations/Npp060305050087/default.pdf>

mesocortical DA by cortical 5-HT_{2A} receptors could be a local effect or possibly involve a polysynaptic neural circuit.

Previous studies have suggested that 5-HT₂ family receptors may regulate mesocortical DA neuronal activity. Systemic administration of DOI, a nonsubtype-selective 5-HT_{2A/B/C} agonist, increased cortical DA efflux which could be blocked by the 5-HT_{2A}-preferring antagonist M100907 ($K_i = 1.92$ for r5-HT_{2A}; see on-line database: <http://kiddb.case.edu/pdsp.php>; Gobert and Millan, 1999). This effect is also observed when M100907 is infused directly into the cortex, suggesting a role for mPFC 5-HT_{2A} receptors (Pehek et al, 2001). Unfortunately, M100907 has appreciable affinities for 5-HT_{2C} ($K_i = 88$ nM; <http://kiddb.case.edu/pdsp.php>) and many other biogenic amine receptors including α_1 - and α_2 -adrenergic and D1-dopamine receptors (<http://kiddb.case.edu/pdsp.php>). The primary goal of the present study was to determine the local 5-HT₂ receptor subtype responsible for regulating phasic DA release in the mPFC. To accomplish this, we examined a panel of potent 5-HT_{2A} receptor antagonists (M100907, MDL 11,939, and SR46349B) along with a highly selective 5-HT_{2B/C} inverse agonist SB 206553. Since the actual selectivity of these putatively 'selective' compounds is not known, receptor-binding studies were conducted to determine the preferential 5-HT_{2A} vs 5-HT_{2C} receptor affinities of MDL 11,939 and SR46349B, compared to M100907. The effect of intracortical administration of these antagonists on basal DA efflux was first determined to see whether the reported negligible role of cortical 5-HT_{2A} receptors for basal DA function (Pehek et al, 2001) would be further confirmed. 5-HT₂ receptor subtype involvement in DA release evoked by administration of the 5-HT₂ agonist DOI was then determined using the 5-HT₂ receptor antagonists that showed preferential binding for the 5-HT_{2A} vs 2C receptor in the receptor-binding study above. Glutamate was also measured in the VTA after DOI and M100907 administration to begin to assess whether corticostriatal mechanisms might potentially mediate the effects of DOI on cortical DA release. And finally, we assessed whether mPFC 5-HT₂ family receptors may modulate mesocortical DA release induced by natural physiological phenomena. Since mesocortical dopamine neurons are extremely responsive to stress (Thierry et al, 1976; Abercrombie et al, 1989; Sorg and Kalivas, 1993; Feenstra and Botterblom, 1996; Inglis and Moghaddam, 1999), and stress increases 5-HT in the mPFC (eg Kawahara et al, 1993), we examined the effect of cortical administration of the 5-HT_{2A} receptor antagonist M100907 on DA release induced physiologically by stress (20 min of gentle handling). To assess the potential involvement of other 5-HT₂ receptor subtypes, these results were contrasted with those following cortical administration of the selective 5-HT_{2B/C} inverse agonist, SB 206,553. To extend the results to agents employed clinically, the atypical antipsychotic drug risperidone, which is a potent 5-HT_{2A} antagonist ($K_i = 0.26$ at r5-HT_{2A}) and weak 5-HT_{2C} antagonist ($K_i = 87$ nM), was also tested (<http://kiddb.case.edu/pdsp.php>). It was hypothesized that intracortical infusions of 5-HT_{2A} antagonists would attenuate DOI and stress-induced DA release. It was further posited that DOI administration would increase VTA dialysate glutamate levels through an interaction with cortical 5-HT_{2A} receptors. Our results demonstrate that 5-HT_{2A} receptors

modulate both drug- and stress-induced DA release in the mPFC and that some of the effects of atypical antipsychotic drugs such as risperidone on DA release are most likely mediated via 5-HT_{2A} receptor blockade.

MATERIALS AND METHODS

Receptor-Binding Studies

Receptors. Cell lines stably transfected with recombinant cDNA encoding receptors or cell lines that express endogenous receptors were used for the comprehensive screening using the resources of the National Institute of Mental Health's Psychoactive Drug Screening Program (PDSP) as previously detailed (Roth et al, 2002). The recombinant receptors included (1) human adrenergic receptors, α_1A , α_1B , α_2A , α_2B , α_2C , and rat adrenergic receptors, β_1 and β_2 ; (2) rat cannabinoid CB₁ receptor; (3) dopaminergic receptors, hD₁, rD₂, rD₃, rD₄, and hD₅; (4) human histamine receptors, H₁, H₂, and H₄; (5) rat imidazoline receptor; (6) human muscarinic acetylcholine receptors, M₁, M₂, M₃, M₄, and M₅; (7) human nicotinic acetylcholine receptors, α_2/β_2 , α_2/β_4 , α_3/β_2 , α_3/β_4 , α_4/β_2 , and α_4/β_4 ; (8) human opiate receptors, μ , δ , and κ ; (9) human peptide receptors, V₁, V₂, V₃, and OT; (10) serotonergic receptors, h5-HT_{1A}, r5-HT_{1B}, h5-HT_{1E}, r5-HT_{2A}, h5-HT_{2B}, r5-HT_{2C}, h5-HT₃, h5-HT_{5A}, h5-HT₆, and h5-HT₇; (11) human transporters of serotonin, norepinephrine, and dopamine as previously described (Roth et al, 1998; Rothman et al, 2000; Roth et al, 2001; Roth et al, 2002; Shapiro et al, 2002); and (12) rat metabotropic glutamate receptors, mGluR_{1a}, mGluR₂, mGluR₄, mGluR_{5a}, mGluR₆, and mGluR₈ as previously described (Gomez et al, 1996; Wroblewska et al, 1997; Kozikowski et al, 1998; Shi et al, 2003). The endogenous receptors included (1) GABA receptors, GABA_A, GABA_B, and GABA_{BZP} from rat forebrain; (2) histamine receptor H₁ from rat forebrain; (3) rat nicotinic acetylcholine receptor, α_4/β_2 ; (4) ionotropic NMDA glutamate receptor from rat forebrain; and (5) voltage-sensitive Ca²⁺ channel from rat heart. For the cloned receptors, HEK-293 cells were either stably or transiently transfected with the receptor of interest with the following exceptions: muscarinic receptors were expressed in CHO cells (see Davies et al (2005) for details) and r5HT_{2A} and r5HT_{2C} receptors were expressed in stable cell NIH 3T3 cell lines.

Radioligand competitive binding assays. Receptor-binding affinities were determined by radioligand competitive binding experiments. The assays were performed as previously detailed using cloned receptor preparations (Roth et al, 1998, 2001; Rothman et al, 2000; Shapiro et al, 2002), or with forebrain membrane preparations containing the endogenous receptors of interest (Roth et al, 1991).

Competitive binding assays were carried out under standard conditions as detailed previously (Roth et al, 2001, 2002; Rothman et al, 2000). The conditions for the different binding assays and K_i values for reference ligands are as described previously (Shi et al, 2003). On-line protocols for binding assays are available at <http://>

pdsp.cwru.edu/nimh/binding.htm. Binding data were performed in quadruplicate.

Microdialysis Studies

Animals. Male Sprague–Dawley rats (Zivic Miller, Alison Park, PA or Harlan, Indianapolis, IN), weighing between 200 and 350 gm at the time of surgery, were used for all experiments. Rats were housed two per cage in a temperature-controlled room with a 12/12 h light/dark cycle. Food and water were available *ad libitum*. All animal procedures were in strict accordance with the NIH Guide for the Care and Use of Laboratory Animals and were approved by the local animal care committee. All efforts were made to minimize the number of animals used and their suffering.

Surgery. Rats were anesthetized with a mixture of xylazine and ketamine (6 and 70 mg/kg, respectively; administered i.m.) and mounted on a stereotaxic frame. After dura was removed, 21 g stainless steel guide cannulae were mounted on the brain surface above the mPFC (+3.2 AP, ML 0.8) (Paxinos and Watson, 1998). For dual-probe studies, a second cannula was mounted above the VTA (−5.60 AP, ML 0.6). The cannulae, with wire obturators, were secured in place with three set screws covered with cranioplastic cement. Animals were returned to the colony room, singly housed and allowed to recover for 2–5 days prior to the experiments. Probe locations were verified histologically at the completion of the experiments. If improperly placed, the animal was excluded from the experiment.

Microdialysis. Microdialysis probes were of a concentric flow design (Yamamoto and Pehek, 1990). mPFC probes were constructed with a 5.0 mm active dialyzing surface membrane (Spectra/Por Hollow, MW cutoff = 13 000, diameter = 240 μ m) to effectively dialyze from the dorsal anterior cingulate to the most ventral region of the infralimbic mPFC. VTA probes were constructed with a 1.0 mm active dialyzing surface at the most ventral extension of the probe to effectively dialyze the mediolateral parabrachial and paranigral VTA. The afternoon prior to microdialysis experiments animals were placed in clear plexiglas microdialysis chambers (Harvard Apparatus) with food and water available *ad libitum*. Probes were lowered slowly through the guide cannulae of awake rats and secured in place with cyanoacrylate glue. Animals were then returned to their dialysis chambers and tethered to counterbalance arms (Instech, Plymouth Meeting, PA) that permitted free movement about the chamber. A microinfusion pump (PHD 2000TM, Harvard Apparatus) and liquid swivel (Instech) were used to perfuse a modified Dulbecco's artificial cerebrospinal fluid (aCSF) buffer solution (137 mM NaCl, 3 mM KCl, 1.2 mM MgSO₄, 0.4 mM KH₂PO₄, with 1.2 mM CaCl₂ and 10 mM glucose; pH 7.4) through the probes. After baseline collections, drugs dissolved in the aCSF were administered by reverse dialysis. Tubing connections were switched manually while maintaining a constant flow rate and collection volume. Samples were immediately analyzed for DA content and frozen for later assay of glutamate.

Drugs. (±)-DOI hydrochloride, (±)-2,5-Dimethoxy-4-iodoamphetamine hydrochloride, was obtained from Sigma-Aldrich (St Louis, MO). M100907, R-(+)-4-[1-hydroxy-1-(2,3-dimethoxyphenyl)methyl]-N-2-(4-fluoro-phenylethyl)-piperidine, was kindly donated by Marion Merrell Dow; risperidone was kindly donated by Janssen Pharmaceutica; SB 206553, N-3-Pyridinyl-3,5-dihydro-5-methylbenzo(1,2-b:4,5-b')dipyrrole-1(2H)carboxamide hydrochloride 5-methyl-1-(3-pyridylcarbonyl)-1,2,3,5-tetrahydropyrrolo[2,3-f]indole, was kindly donated by Smithkline and Beecham; and SR 46349B, trans, 4-([3Z]3-(2-dimethylaminoethyl)oxyimino-3(2-fluorophenyl + + +)propen-1-yl]phenol hemifumarate, was kindly donated by Sanofi Recherche. MDL 11,939, α -Phenyl-1-(2-phenylethyl)-4-piperidinemethanol, was purchased from Tocris Cookson (Bristol, UK). For intracortical administration, all drugs were initially dissolved in water containing 1.5–5 μ l of glacial acetic acid to make a 10 mM stock solution. They were then further diluted to 1.0–10.0 μ M concentrations with aCSF. Concentrations refer to the base forms of the antagonists. The pH of all aCSF solutions was adjusted to 7.4. DOI was injected systemically (s.c.) and was dissolved in water (2.5 mg/kg/ml, dose refers to the salt).

Chromatography. HPLC with electrochemical detection was used to measure dialysate DA, 5-HT, and glutamate content.

DA and 5-HT: DA and 5-HT concentrations in dialysate samples were measured by reverse phase HPLC coupled with electrochemical detection. In total, 20 μ l samples were injected immediately after collection onto a 2 \times 100 mm Phenomenex (Torrance, CA) column (UltracarbTM, 3 μ m particle size, ODS 20). The mobile phase consisted of 32 mM anhydrous citric acid, 54 mM sodium acetate trihydrate, 0.074 mM EDTA, 0.215 mM octylsulfonic acid, and 3% methanol (vol/vol), pH 4.2. To maintain separation of DA and 5-HT from its metabolites and 5-hydroxyindoleacetic acid, the pH of the mobile phase and the concentration of the octylsulfonic acid were adjusted as needed. A BAS LC-4C or Antec INTRO electrochemical detector with a glassy carbon electrode, maintained at a potential of +0.60 V relative to an Ag/AgCl reference electrode, was employed.

Glutamate: Frozen dialysate samples were defrosted and 14 μ l was placed into a refrigerated autosampler (ThermoQuest, Thermo Separation Products, San Jose, CA). Samples were precolumn derivatized with ophthaldehyde (OPA) using a slightly modified version of the procedure of Donzanti and Yamamoto (1988). The derivatizing solution was prepared by mixing 27 mg OPA with 1 ml methanol. Then, 9 ml of 0.05 M sodium tetraborate and 5.0 μ l of β mercaptoethanol were added. This mixture was diluted further with sodium tetraborate (1:3). Next, 7 μ l of this solution was automatically mixed with the sample and then 10 μ l was injected onto a 2 \times 100 mm Phenomenex column (UltracarbTM, 3 μ m particle size, ODS 20), maintained at 35°C. The mobile phase consisted of 14.2 g Na₂HPO₄, 0.5 g Na₂EDTA, 25% methanol, and pH 4.2. A Bioanalytical Systems LC-4C electrochemical detector with a glassy carbon electrode was used to detect the derivatized amino acids and was maintained at a potential of 0.6 V (range 20 nA/V).

Histology. After microdialysis experiments were completed, probe placements were verified histologically. Only animals, whose probe placements were verified to be in the mPFC, or the mPFC and the VTA for the dual probe study, were used in this study.

Data analysis. Microdialysis data were expressed as the percentage of the average baseline level (last three baseline samples prior to treatment). Two-way ANOVAs, with drug group as the between factor and time as the within factor, were used to analyze data between drug treatment groups. One-way repeated measures ANOVAs, with time as the within factor, were used to analyze data from individual drug treatment groups. Following significant F-values, Dunnett's *post hoc* tests for comparing treatment means with a control were employed.

Procedures

Receptor-binding properties of two 5-HT_{2A} antagonists: MDL 11,939 and SR 46349B. This experiment screened the receptor-binding properties of MDL 11,939 and SR 46349B. If a compound displaced radioligand binding by 50% or more, then its K_i was determined as detailed above.

Effects of the 5-HT_{2A} antagonists MDL 11,939 and SR 46349B on cortical DA release. This experiment investigated the effects of intracortical administration of MDL 11,939 and SR 46349B on basal DA efflux in the mPFC. In addition, since Experiment 1 determined that MDL 11,939, but not SR46349B, showed strong preferential selectivity for the 5-HT_{2A} vs the 5-HT_{2C} receptor, we examined the effects of this ligand on 5-HT₂ agonist (DOI)-stimulated DA release. At 2–4 days after rats had been implanted with cannulae above the mPFC, microdialysis probes were lowered slowly through the cannulae. After 18–24 h, aCSF perfusion began at a rate of 1.0 μ l/min. Samples were collected every 30 min for immediate determination of DA in the mPFC. After baselines had stabilized (approximately 3 h), either SR 46349B or MDL 11,939 was perfused through the mPFC of two separate groups of rats to test the effect of 5HT_{2A} receptor antagonism on basal DA efflux. SR 46349B was perfused through the mPFC for 1 h at a concentration of 1.0 μ M followed by 1 h at a concentration of 10 μ M as similarly tested with M100907 (Pehek et al, 2001). MDL 11,939 was perfused at a 10 μ M concentration for 3 h. To test the effect of 5HT_{2A} receptor antagonism on drug-induced DA efflux, one group of rats was injected with DOI (2.5 mg/kg s.c.) 30 min after perfusion with 1.0 μ M MDL 11,939 began (Pehek et al, 2001). This group was compared to a group receiving DOI injection and drug-free aCSF and a control group that received vehicle injections but no drug.

5-HT_{2A} receptor regulation of stress-induced DA release in the PFC. This study determined whether intracortical infusions of the 5-HT_{2A} antagonist M100907 would attenuate DA efflux elicited by a mild stress. Since M100907 has been extensively utilized in the literature to assess 5-HT_{2A} receptor function compared to MDL 11,939, and we previously found that cortical 10 μ M M100907 blocked DOI-induced DA release in the mPFC (Pehek et al,

2001), it was implemented in this study to begin to assess 5-HT_{2A} receptor mechanisms in stress-induced cortical DA release. This concentration of M100907 does not alter basal DA levels in the mPFC (Pehek et al, 2001). Probes were lowered into guide cannulae placed over the mPFC in awake rats 18–24 h before the experiment began. At the start of the experiment, probes were perfused at a rate of 1.5 ml/min and samples were collected every 20 min until baselines were stable. A 10 μ M concentration of M100907 or drug-free aCSF was then perfused through the probes in two separate groups of rats. After 20 min, rats were stressed for 20 min. The stressor consisted of gentle stroking ('petting'), a procedure which reliably increases DA efflux in rats (Inglis and Moghaddam, 1999). In M100907-treated animals, drug infusion continued during the stressor and until the end of the experiment. These results were compared to those following 10 μ M infusions of the selective 5-HT_{2B/C} inverse agonist SB 206553. Local administration of 10 μ M SB206553 does not alter basal DA levels in the mPFC (Alex et al, 2005).

Effects of risperidone on stress-induced DA release in the MPFC. This experiment examined whether administration of risperidone by reverse dialysis would alter handling-induced cortical DA efflux. Procedures were the same as the previous experiment except risperidone was infused beginning 40 min before the stressor was applied, to show that basal DA efflux was stable prior to stress application. Risperidone increases DA release in the mPFC when administered systemically, but cortical application of selective 5-HT_{2A} antagonists do not (see Discussion). These results were compared to a control group that received aCSF without drug.

Cortical 5-HT_{2A} receptor regulation of mesocortical DA and corticostriatal glutamate. This experiment investigated the regulation of DA efflux in the mPFC and glutamate in the VTA by 5-HT_{2A} receptors in the mPFC. Due to the extensive use of M100907 in the literature, and to tie the findings in this study to previous results (Pehek et al, 2001), we used 10 μ M M100907 to assess cortical 5-HT_{2A} receptor involvement. Rats were implanted with two cannulae: one above the mPFC and one above the VTA. After 2–4 days, on the morning of the experiment, microdialysis probes were lowered slowly through these cannulae. aCSF was then perfused immediately at a rate of 1.0 μ l/min through both probes. This procedure of initiating the experiment immediately following probe insertion was utilized because we and others have found that brain injury-derived, that is, non-neuronal, levels of glutamate are higher 18–24 h following probe implantation (Obrenovitch et al, 1993; Xue et al, 1996). Samples were collected every 30 min for immediate determination of DA in the mPFC and subsequent determination of glutamate in the VTA. The VTA dialysate was frozen for later analysis of glutamate. Samples were collected until baseline dopamine levels were stable (approximately 3 h after probe implantation). A 10 μ M solution of M100907 or aCSF without drug (control) was then infused into the mPFC. After 30 min, half of each group also received either an injection of the 5-HT₂ agonist DOI (2.5 mg/kg s.c.) or vehicle (water). This created four treatment groups: (1) intracortical M100907 and s.c. DOI,

(2) intracortical aCSF and s.c. DOI, (3) intracortical M100907 and s.c. vehicle, and (4) intracortical aCSF and s.c. vehicle. Only animals with correct placements in both the VTA and mPFC were included in data analysis.

RESULTS

Receptor-Binding Properties of Two 5-HT₂ Antagonists: MDL 11,939 and SR 46349B

Both MDL 11,939 and SR 46349B had high affinities for 5-HT_{2A} receptors (K_i (nM) = 2.89 and 0.95, respectively, see Table 1). However, these ligands differed with respect to their affinities for 5-HT_{2C} receptors. While SR 46349B had high affinity for 5-HT_{2C} receptors (K_i = 8.24), MDL 11,939 had relatively low affinity (K_i = 853.6). Thus, of the two ligands, MDL 11,939 was the most selective. For comparison, the putatively selective 5-HT_{2A} antagonist M100907 has higher affinity for the 5-HT_{2A} (K_i = 1.92) vs the 5-HT_{2C} (K_i = 88) receptor (data from PDSP data base: <http://kidb.case.edu/pdsp.php>). Both MDL 11,939 and SR 46349B had low or no affinity for non-5-HT₂ receptors (see Table 1). M100907 has been previously reported to have appreciable affinity for various adrenergic receptors, H1-histamine, and sigma receptors and low affinities for D1 and D4 dopamine receptors (<http://kidb.case.edu/pdsp.php>).

Effects of the 5-HT_{2A} Antagonists MDL 11,939 and SR 46349B on Cortical DA Release

Figure 1a illustrates the location of dialysis probes within the mPFC for this and subsequent experiments. Figure 2a shows that intracortical perfusion with either 1.0 or 10 μ M SR 46349B (n = 7) or 10 μ M MDL 11,939 (n = 5) did not significantly alter basal DA efflux (average basal levels of DA for these two groups were 0.56 ± 0.07 pg/20 μ l). As shown in Figure 2b, intracortical MDL 11,939 pretreatment blocked DOI-induced DA release in the mPFC. A two-way ANOVA (drug group \times time) indicated that extracellular DA levels were significantly different in animals treated systemically with vehicle, DOI alone, or DOI plus intracortical MDL 11,939 (overall drug group effect: $F(2,19) = 4.47$, $p = 0.03$). Indeed, rats treated with DOI alone showed significantly higher DA levels than MDL 11,939 + DOI-cotreated rats (drug group effect: $F(1,13) = 4.67$, $p = 0.05$). *Post hoc* one-way ANOVAs (time) revealed that animals treated with DOI alone showed significantly increased dialysate DA levels following treatment (time effect: $F(6,54) = 4.18$, $p = 0.002$; $n = 10$; see DOI group Figure 2b), while vehicle controls ($n = 7$) and animals cotreated with DOI plus intracortical MDL 11,939 (1 μ M, $n = 5$) did not (p 's > 0.05). Average basal DA levels for these last three groups were 0.41 ± 0.04 pg/20 μ l, $n = 22$).

5-HT_{2A} Receptor Regulation of Stress-Induced DA Release in the mPFC

As shown in Figure 3, mild handling (stroking) for 20 min increased DA to 182% of baseline (time effect: $F(5,50) = 4.73$, $p = 0.001$, $n = 11$, average basal levels = 0.38 ± 0.06 pg/20 μ l). 5-HT was also increased to 197% (time effect: $F(5,55) = 8.30$, $p < 0.001$, $n = 12$, see Figure 3, basal

Table 1 K_i Values of MDL 11,939, SR 46349B (mean \pm SEM) and M100907^a in Competitive Binding Assays

Membrane-bound proteins	Drugs		
	MDL 11,939	SR 46349B	M100907
Receptors			
Serotonin			
5-HT _{1A}	3416 \pm 1069	2500 \pm 410	> 10 000
r5-HT _{1B}	70.46 \pm 23.0	914 \pm 314	> 10 000
5-HT _{1E}	No binding	No binding	> 10 000
r5-HT _{2A}	2.893 \pm 0.56	0.9507 \pm 0.11	1.92
5-HT _{2B}	1419 \pm 150	NT	261
r5-HT _{2C}	853.6 \pm 185.6	8.243 \pm 1.37	88
5-HT ₃	No binding	No binding	> 10 000
5-HT _{5a}	No binding	1750 \pm 435	> 10 000
5-HT ₆	2752 \pm 1014	3868 \pm 933	5000
Dopamine			
D1	327.2 \pm 43.8	> 10 000	2964
rD2	No binding	No binding	> 10 000
rD3	No binding	> 10 000	> 10 000
D4	3934 \pm 1334	NT	> 10 000
D5	2869 \pm 646	NT	> 10 000
Acetylcholine (muscarinic)			
M1	No binding	732.7 \pm 44.30	> 10 000
M2	No binding	13 800 \pm 1081	> 10 000
M3	No binding	2387 \pm 255.2	> 10 000
M4	No binding	3672 \pm 513.2	> 10 000
M5	No binding	3015 \pm 203.0	> 10 000
Adrenergic			
Alpha _{2A}	1586 \pm 263.7	725.9 \pm 86.06	1900
Alpha _{2C}	1186 \pm 118.6	2118 \pm 84.37	1442
Alpha _{1A}	1876 \pm 51.72	No binding	128
Alpha _{1B}	1579 \pm 117.9	No binding	424.7
rBeta ₂	No binding	No binding	> 10 000
Transporters			
Serotonin	2598 \pm 417	No binding	> 10 000
Norepinephrine	> 10 000	739.5 \pm 136.4	> 10 000
Dopamine	> 10 000	5619 \pm 1817.3	> 10 000
Histamine			
hH1	1604 \pm 282.6	No binding	66.7
hH4	No binding	No binding	> 10 000

No binding: did not displace radioligand binding by 50% or more; NT: not tested; r: rat.

^aFrom PDSP website: <http://kidb.case.edu/pdsp.php>.

levels = 0.22 ± 0.04 pg/20 μ l). As shown in Figure 4a, mild handling stress increased mPFC DA levels again in a subsequent experiment, and intracortical M100907 pre-




Figure 1 Microdialysis probe placements. (a) Schematic of representative probe placements in the medial prefrontal cortex. (b) Schematic of representative probe placements in the ventral tegmental area (VTA). The 1.0 mm active dialyzing surface of VTA probes was located at the most ventral extension of the probe.

treatment blocked this effect. A two-way ANOVA (drug group \times time) of this data revealed a significant drug group \times time interaction ($F(6,102) = 3.89$, $p = 0.002$). Indeed, mild handling significantly increased mPFC DA levels in vehicle-treated animals (time effect: $F(6,36) = 3.97$, $p = 0.004$, $n = 7$; Figure 4a), but not in animals treated with intracortical 10 μ M M100907 ($F(6,66) = 1.306$, $p > 0.05$, $n = 12$). As seen in Figure 4b, mild handling still increased mPFC DA levels in animals treated intracortically with 10 μ M SB 206553 (time effect: $F(6,30) = 5.37$, $p < 0.001$, $n = 6$). The average basal levels of DA for these last three groups were 0.43 ± 0.04 pg/20 μ l, $n = 25$). Thus, cortical 5-HT_{2A} receptor blockade diminished stress-induced increases in mPFC DA, while 5-HT_{2B/2C} receptor blockade did not.

Effects of Risperidone on Stress-Induced DA Release in the mPFC

As shown in Figure 4c, 20 min handling stress again increased cortical DA levels in another group of rats, and local risperidone treatment blocked this effect. A two-way ANOVA (drug group \times time) of this data revealed a significant drug group \times time interaction ($F(6,66) = 3.68$, $p = 0.003$) between these two groups. Indeed, vehicle-treated rats showed significantly increased cortical DA efflux following handling (time effect: $F(6,36) = 6.27$, $p < 0.001$, $n = 7$; Figure 4c), while animals pretreated locally with 10 μ M risperidone infusion did not ($F(6,30) = 0.21$, $p > 0.05$, $n = 6$). The average basal levels of DA for this experiment were: 0.68 ± 0.08 pg/20 μ l, $n = 13$.

Cortical 5-HT_{2A} Receptor Regulation of Mesocortical DA and Corticostriatal Glutamate

Prefrontal cortical dopamine. A total of 26 rats were used for this experiment but, due to chromatographic problems with the DA assay, only 18 were used for the DA measurements (all 26 were used for the glutamate measurements, see below). Both the VTA and mPFC probes

were functional in all 26 rats (see probe locations in Figure 1b and a, respectively). As seen in Figure 5a, systemic injections of DOI alone increased DA levels in the mPFC of rats, but this was not seen in animals cotreated with DOI and intracortical M100907. A two-way ANOVA (drug group \times time) revealed that mPFC DA levels were significantly different between animals in the four treatment groups shown in Figure 5a (overall drug group effect: $F(3,14) = 3.79$, $p = 0.035$). *Post hoc* analysis indicated that rats treated with DOI alone showed significantly increased dialysate DA levels in the mPFC following treatment (time effect: $F(6,18) = 2.92$, $p = 0.04$, $n = 4$). These subsided and returned to baseline levels by 2 h post injection. However, rats given DOI plus intracortical M100907 cotreatment showed the reverse ($F(6,24) = 2.77$, $p < 0.03$, $n = 5$). Indeed, infusions of M100907 reversed the early DA increases by DOI and gradually diminished DA levels once the effect of DOI had subsided. There were no changes in DA levels in vehicle controls ($p > 0.05$, $n = 4$). Although DA levels diminished in animals treated with intracortical M100907 alone (time effect: $F(6,24) = 3.47$, $p < 0.01$, $n = 5$), significant decreases relative to controls were only shown at two later time points across the 3 h treatment (p 's < 0.05 , see Figure 5a). Average basal levels for this experiment were 0.50 ± 0.03 pg/20 μ l, $n = 18$.

VTA glutamate. The above four treatment groups also had an additional microdialysis probe located in the VTA, which allowed simultaneous assessment of VTA glutamate response to systemic DOI treatment. As can be seen in Figure 5b, systemic injections of DOI increased glutamate in the VTA and intracortical M100907 cotreatment reversed this effect. A two-way ANOVA (drug group \times time) revealed that glutamate levels were significantly different between animals in the four treatment groups (overall drug group effect: $F(3,22) = 3.08$, $p = 0.048$). A marginal drug group \times time interaction was also shown ($F(18,132) = 3.68$, $p = 0.053$). Indeed, systemic DOI significantly increased dialysate glutamate levels (time effect: $F(6,36) = 3.20$, $p = 0.01$, $n = 7$), while DOI plus intracortical M100907 cotreatment did not




Figure 2 Effects of intracortical administration of the 5-HT_{2A} antagonists SR 46349B and MDL 11,939 on DA release in the medial prefrontal cortex. (a) Intracortical infusions of SR 46349B or MDL 11,939 did not alter basal DA release. SR 46349B was infused at a 1.0 μ M concentration for 1 h followed by a 10.0 μ M concentration for 1 h. MDL 11,939 (10 μ M) was infused for 3 h. (b) Intracortical infusions of MDL 11,939 blocked DOI-stimulated DA release. MDL 11,939 (1.0 μ M) was infused for 3 h. Times of intracranial drug infusions are indicated by the lines. The 5-HT₂ agonist DOI (2.5 mg/kg s.c.) or vehicle was injected at time 30 min as indicated by the arrow. Data are the means \pm SEMs. * p < 0.05 relative to pre-drug baseline.

(see M100907 + DOI group; p > 0.05, n = 8). Neither vehicle injections (n = 6) nor intracortical treatment with M100907 alone (n = 5) altered VTA dialysate glutamate concentrations. Average basal levels of glutamate were $0.45 \text{ ng}/20 \mu\text{l} \pm 0.03$, n = 26. Thus, systemic DOI treatment increased VTA dialysate glutamate levels in the same group of animals that had shown DOI-induced DA increases above in the mPFC (see DOI group, Figure 5a vs b). Similarly, this effect was reversed by intracortical M100907 cotreatment.

DISCUSSION

The main finding of the present work is that blockade of prefrontocortical 5-HT_{2A} receptors blocks the evoked




Figure 3 Stress increases 5-HT and DA release in the medial prefrontal cortex. Rats were gently handled for 20 min at the time indicated by the hatched bar. Data are the means \pm SEMs. * p < 0.05 relative to pre-drug baseline.

release of DA from the mesocortical pathway. Importantly, 5-HT_{2A} receptors could be implicated in regulating both pharmacologically and physiologically induced DA release. Intracortical perfusion with two different 5-HT_{2A} antagonists, MDL 11,939 and M100907, attenuated DA efflux induced by the systemic administration of the 5-HT₂ agonist DOI. These antagonists, as well as a third, SR 46349B, either did not alter basal dialysate DA or produced a small decrease, indicating that cortical 5-HT_{2A} receptors do not generally modulate basal DA release in the mPFC. Rather, either the pharmacological stimulation of 5-HT_{2A} receptors or their physiological stimulation by endogenous 5-HT may act to boost cortical DA levels. One situation in which this may occur is during stress. Stress is known to increase cortical 5-HT (Kawahara *et al*, 1993) and DA release (eg Abercrombie *et al*, 1989). Stress increased both neurochemicals in the present work (see Figure 3). Furthermore, antagonism of cortical 5-HT_{2A} receptors blocked stress-induced DA efflux. In contrast, local inverse agonism of 5-HT_{2B/C} receptors did not. Thus, endogenous 5-HT, released during stress, appears to act on prefrontocortical 5-HT_{2A} receptors to potentiate DA release.

These findings agree with, and build upon, recently published data. Gobert and Millan (1999) demonstrated that systemic administration of M100907 blocked DOI-induced DA release in the PFC. However, as mentioned previously, due to the robust pharmacology of M100907, it is not possible from these studies to unequivocally invoke 5-HT_{2A} receptors in this response. Nonetheless, De Deurwaerdere and Spampinato (1999) observed that treatment with the 5-HT_{2A/2C} antagonist SR 46349B also blocked phasic DA release in the nucleus accumbens. Systemic administration of M100907 attenuated fluoxetine-induced increases in cortical DA (Zhang *et al*, 2000). The current study compared the intracortical administration of several 5-HT_{2A} antagonists (M100907, MDL 11,939, risperidone) on phasic DA release in the mPFC. The findings indicate that 5-HT_{2A} receptors regulate phasic mesocortical DA release, and that these receptors are localized, at least in part, to the medial PFC, where they are abundant.




Figure 4 Effects of intracortical administration of 5-HT₂ antagonists on acute stress-induced DA release in the medial prefrontal cortex. (a) Intracortical perfusion with M100907 (10 μM) blocked stress-induced DA efflux. (b) Intracortical perfusion with SB 206553 did not alter stress-induced DA efflux. (c) Intracortical perfusion with risperidone (10 μM, RISP) blocked stress-induced DA efflux. Drugs were infused for 2 h as indicated by the lines. Rats were gently handled for 20 min at the times indicated by the hatched bars. In contrast to intracortical M100907 and SB 206553 pretreatment, risperidone infusion began 40 min before handling stress was applied to be sure that mPFC DA levels were stable under the treatment prior to stress application (see text). Data are the means ± SEMs. **p* < 0.05 relative to pre-drug baseline.

The present work demonstrates that M100907, MDL 11,939, and SR 46349B are all potent 5-HT_{2A} ligands, with an order of potency SR 46349B > M100907 > MDL 11,939.




Figure 5 Effects of systemic DOI and intracortical M100907 on transmitter release in the medial prefrontal cortex and VTA using dual-probe microdialysis. (a) Intracortical infusions of M100907 blocked DOI-stimulated DA release in the medial prefrontal cortex. M100907 (10.0 μM) was infused for 3 h as indicated by the line. DOI (2.5 mg/kg s.c.) or vehicle was injected at time 30 min as indicated by the arrow. (b) Intracortical infusions of M100907 blocked DOI-stimulated glutamate release in the VTA. M100907 (10.0 μM) was infused for 3 h as indicated by the line. DOI (2.5 mg/kg s.c.) or vehicle was injected at time 30 min as indicated by the arrow. Data are the means ± SEMs. **p* < 0.05 relative to pre-drug baseline.

However, SR 46349B is much less selective (only eight-fold selectivity for the 5-HT_{2A} vs the 5-HT_{2C} receptor, see Table 1). Recent work has consistently shown that the systemic administration of M100907 alone does not alter basal mPFC DA levels over a wide range of doses (eg Bonaccorso *et al*, 2002). While moderate doses (1–3 mg/kg) of SR 46349B also did not alter DA in the mPFC or nucleus accumbens, a higher dose (10 mg/kg) increased cortical DA efflux (Bonaccorso *et al*, 2002). This increase may reflect the relative nonselectivity of SR 46349B and may have resulted from actions on 5-HT_{2C} receptors (*K_i* = 8.24, see Table 1). Other work has shown that systemic administration of 5-HT_{2C} antagonists or inverse agonists increases cortical DA efflux (Millan *et al*, 1998; Gobert *et al*, 2000; Pozzi *et al*, 2002). In contrast to systemic administration, we clearly show that intracortical SR 46349B, as well as MDL 11,939 do not alter basal DA levels in the mPFC.

Cortical administration of the 5-HT_{2B/2C} inverse agonist, SB206553, also does not alter basal mPFC DA levels (Alex *et al*, 2005). Thus, increases in cortical DA efflux by systemic application of drugs with 5-HT_{2A} or 5-HT_{2C} antagonist properties is not due to their interaction with cortical 5-HT_{2A} or 5-HT_{2C} receptors. Interestingly, intracortical infusions of M100907 slightly decreases basal DA release (Pehek *et al*, 2001; current work, see Figure 5). The equally potent 5-HT_{2A} antagonist MDL 11,939 did not diminish basal DA levels in this study, and it is even more preferentially selective for the 5-HT_{2A} vs 5-HT_{2C} receptors than M100907 (see Table 1). The moderate affinity of M100907 for the 5-HT_{2C} receptor ($K_i = 88.0$ nM, see Table 1) would not induce this decrease (Alex *et al*, 2005). Possibly the slight change by M100907, and absence by MDL 11,939, is due to their differential α_1 adrenergic receptor-binding properties ($K_i = 128$ vs 1800, respectively; see Table 1). Local antagonism of adrenergic α_1 receptors diminish DA levels in the mPFC (Pan *et al*, 2004). Nonetheless, selective antagonism of cortical 5-HT_{2A} receptors clearly does not increase basal DA efflux in the mPFC.

In contrast to studies employing the administration of 5-HT_{2A} antagonists alone, the combined, systemic administration of D₂ and 5-HT_{2A} receptor antagonists results in a potentiation of DA release (Westerink *et al*, 2001; Liegeois *et al*, 2002). Although the exact mechanism is not clear, evidence has been provided that this effect may be mediated by a combined potent 5-HT_{2A} and relatively weak D₂ receptor effect interacting with 5-HT_{1A} receptor-regulated brain mechanisms (Bonaccorso *et al*, 2002; Ichikawa *et al*, 2001). This potentiation may also result from the actions of drugs on DA cells in the VTA. We have shown that many VTA DA neurons express 5-HT_{2A} receptors (Nocjar *et al*, 2002). It is clear from the present work that selective antagonism of cortical 5-HT_{2A} receptors does not increase basal DA efflux but, in fact, blocks evoked DA release.

Electrophysiological studies indicate that stimulation of mPFC 5-HT_{2A} receptors increases pyramidal cell activity (Aghajanian and Marek, 1997), suggesting that 5-HT_{2A} receptor agonism may increase the activity of corticocortical glutamatergic projection neurons. Indeed, systemic administration of the 5-HT₂ agonist DOI increased glutamate efflux in the VTA in the present study. Infusions of M100907, directly into the mPFC, blocked this increase, implying that the relevant 5-HT_{2A} receptors were localized in the mPFC. In the same animals, treatment with DOI increased mPFC DA efflux and this was also blocked by intracortical administration of M100907. Thus, systemic DOI increased glutamate in the VTA and DA in the mPFC, in the same animals, through a cortical 5-HT_{2A} receptor mechanism. Interestingly, Artigas and co-workers recently found that either the systemic or intracortical administration of DOI increased the activity of VTA mesocortical DA neurons (Bortolozzi *et al*, 2004). This was blocked by the intracortical administration of 5-HT_{2A} antagonists. Thus, both studies suggest that regulation of mesocortical DA by cortical 5-HT_{2A} receptors may involve a polysynaptic neural circuit. A subset of cortical pyramidal neurons synapse on mesocortical DA cell bodies in the VTA (Sesack and Pickel, 1992; Carr and Sesack, 2000), and the majority of 5-HT_{2A} receptors in the mPFC are localized to the apical dendrites of pyramidal cells (Willins *et al*, 1987; Jakab and

Goldman-Rakic, 1998; Miner *et al*, 2003). Other potential circuits would be indirect projections from the mPFC to the VTA. A role for presynaptic regulation of DA release also cannot be discounted. 5-HT_{2A} receptors have been recently found in the mPFC on presynaptic axons and varicosities that expressed morphological features of monoamine axons (Miner *et al*, 2003). We have previously shown that intracortical infusions of M100907 block local K⁺-stimulated DA release in the mPFC (Pehek *et al*, 2001). Since high K⁺ stimulates transmitter release by depolarizing nerve terminals, it is possible that M100907 may have acted to block DA efflux through actions on 5-HT_{2A} receptors localized presynaptically on DA terminals.

The present work demonstrates that mPFC 5-HT_{2A} receptors do not generally modulate basal cortical DA release. Rather, endogenous 5-HT, released phasically, appears to activate cortical 5-HT_{2A} receptors to subsequently stimulate DA release. Acute stress increases the release of cortical 5-HT (Figure 3; Kawahara *et al*, 1993) and DA (eg Abercrombie *et al*, 1989; Kawahara *et al*, 1999). The current findings demonstrate that blockade of cortical 5-HT_{2A} receptors attenuates mild stress-induced DA release: this is the first evidence that cortical 5-HT_{2A} antagonism blocks physiologically induced DA efflux in the mPFC. This effect did not involve other 5-HT₂ family receptors, since the 5-HT_{2B/2C} inverse agonist SB 206553 did not alter stress-induced cortical DA. Thus, physiologically induced phasic release of DA in the mPFC depends, at least partially, on activation of local 5-HT_{2A} receptors in the area. Interestingly, behavioral deficits in prepulse inhibition, which are known to be regulated in the mPFC, are reversed by M100907 in animals with genetically elevated synaptic levels of DA (Barr *et al*, 2004; Gainetdinov *et al*, 1999). Thus, it has been suggested that 5-HT_{2A} antagonists may be useful in the treatment of conditions characterized by chronic, elevated dopaminergic tone (Gainetdinov *et al*, 1999). Indeed, a recent large-scale placebo-controlled clinical trial with SR46349B demonstrated its ability to treat schizophrenia (Meltzer *et al*, 2004), and we have shown in this study that it has a high affinity for the 5-HT_{2A} receptor. M100907 has a similar affinity for the 5-HT_{2A} receptor as SR46349B (see Table 1), but was minimally effective in treating two schizophrenia patients in a recent report (Talvik-Lotfi *et al*, 2000), although the authors stress that efficacy shown with a sample of two is clearly not conclusive (Talvik-Lotfi *et al*, 2000). Nonetheless, SR46349B, in contrast to M100907, also has high affinity for the 5-HT_{2C} receptor (see Table 1). Many atypical antipsychotic drugs that bind to the 5-HT_{2A} receptor also show moderate to potent 5-HT_{2C} receptor-binding properties (Roth *et al*, 2004), suggesting that efficacy may involve some combination of effect between these receptors. However, our current findings with SB206553 indicate that intracortical 5-HT_{2C} antagonism, without 5-HT_{2A} antagonism, does not alter phasic DA release in the mPFC.

In the present study, intracortical infusions of the atypical antipsychotic drug risperidone also blocked stress-induced DA release. This effect may have been mediated, at least partially, by blockade of local 5-HT_{2A} receptors, since cortical M100907 induced a similar effect. It is unknown whether cortical risperidone infusion, as used in the current work, would increase 5-HT levels in the mPFC as has been

shown with systemic risperidone treatment (Ichikawa *et al*, 1998; Hertel *et al*, 1996; although see Zhang *et al*, 2000; Ojima *et al*, 2004). If so, other non-5-HT₂ serotonin receptors could be implicated in this stress effect of risperidone. 5-HT_{2A} and 5-HT_{1A} receptors locally modulate 5-HT in the mPFC (Martin-Ruiz *et al*, 2001). However, risperidone is a 5-HT_{2A} antagonist and cortical 5-HT_{2A} antagonist treatment does not increase 5-HT locally (Martin-Ruiz *et al*, 2001). Also, risperidone has low affinity for the 5-HT_{1A} receptor ($K_i = 250$ nM; <http://kidb.case.edu/pdsp.php>).

Like other atypical antipsychotic drugs, risperidone has a robust pharmacology, binding to many biogenic amine receptors (Roth *et al*, 2004), including 5-HT_{2A}-serotonin and D₂-dopamine receptors. Indeed, this combination of properties may explain why the systemic administration of higher doses (1 mg/kg) of risperidone produces increases in mPFC DA efflux (Hertel *et al*, 1996; Kuroki *et al*, 1999; Zhang *et al*, 2000). Lower doses of risperidone (0.004 mg/kg), like M100907, do not alter cortical DA efflux (Westerink *et al*, 1998, 2001). These lower doses may result in more selective effects on 5-HT_{2A} receptors and are closer to the range employed clinically. Studies suggest that risperidone may not increase prefrontal DA release clinically in schizophrenia patients (Miller *et al*, 2001; Molina *et al*, 2003).

The possibility exists that 5-HT_{2A} receptor blockade may have clinical utility in the treatment of conditions linked to a phasic hyperactivity of DA systems. Preclinical studies indicate that supranormal DA activity in the mPFC may be maladaptive. DA released during stress (Arnsten and Goldman-Rakic, 1998) or supranormal D₁ receptor stimulation (Zahrt *et al*, 1997; Goldman-Rakic *et al*, 2000) both impair cognitive function. Phasic PFC DA function appears to be magnified in schizophrenia patients (Laruelle *et al*, 1999) and basal cortical DA function possibly diminished (see Molina *et al*, 2005a,b). Indeed, PFC D₁ receptors are upregulated in these patients (Abi-Dargham *et al*, 2002). Phasic release of DA by stress or drugs under these conditions would induce a magnified effect. Indeed, schizophrenic patients are more sensitive to stress and more sensitive to the psychotic and cognitive impairing effects of amphetamine (Abi-Dargham *et al*, 1998; Breier *et al*, 1997; Laruelle *et al*, 1999). Stress is also known to exacerbate schizophrenic symptoms. Thus, the clinical efficacy of risperidone, and other atypical antipsychotic drugs, which block 5-HT_{2A} receptors, could be at least partially due to their ability to diminish the effect of stress on mPFC function in these patients.

However, it should be noted that a large body of evidence indicates that reduced DA tone in the mPFC is associated with impaired cognition (Jentsch and Roth, 1999). Indeed, the cognitive deficits and negative symptoms of schizophrenia have been associated with basal dopaminergic hypoactivity in the PFC (Weinberger, 1987). Therefore, it is likely that an optimal balance of DA tone in the PFC is essential for normal cortical function (Williams and Goldman-Rakic, 1995). Highly efficacious antipsychotic drugs may have receptor-binding properties that increase basal mesocortical DA release while decreasing evoked DA release. In fact, it has been proposed that schizophrenia may be related to deficits in both tonic and phasic DA

release (Grace, 1991). Interestingly, intracortical administration of 5-HT_{2A} or 5-HT_{2C} receptor antagonists do not alter basal mPFC DA (Pehek *et al*, 2001; Alex *et al*, 2005; current work). However, systemic administration of selective 5-HT_{2C} antagonists increase basal DA levels in the mPFC (see Di Matteo *et al*, 2001), while the selective 5-HT_{2A} receptor antagonist M100907 does not (eg Bonaccorso *et al*, 2002). In contrast, intracortical M100907 blocks phasic DA release in the mPFC, while selective 5-HT_{2C} antagonists do not (Pehek *et al*, 2001, current work). Thus, the binding affinity of atypical antipsychotic treatments to both of these receptors may be particularly important (Roth *et al*, 2004; Meltzer *et al*, 2003). Indeed, their efficacy could be due to their ability to increase *basal* cortical DA levels through 5-HT_{2C} receptors located outside the PFC, and to their ability to diminish *phasic* cortical DA levels through their interaction with 5-HT_{2A} receptors located within the PFC. An important area of future work would be to determine whether other atypical antipsychotic drugs that increase basal mPFC DA levels when administered systemically to animals (Meltzer *et al*, 2003) would also locally diminish mPFC DA release induced phasically by stress. Indeed, recent PET studies suggest that the highly efficacious drug clozapine may increase *basal* DA function in the PFC of schizophrenia patients (Cohen *et al*, 1997; Lahti *et al*, 2003; Molina *et al*, 2005a; Potkin *et al*, 2003). However, one exposure to amphetamine, which *phasically* increases cortical DA function in these patients, worsens their psychosis (Laruelle *et al*, 1999).

In summary, intracortical administration of two 5-HT_{2A} antagonists blocked mesocortical DA release induced by the systemic administration of the 5-HT₂ agonist DOI. Treatment with DOI also increased glutamate efflux in the VTA and DA release in the mPFC of the same animals; effects blocked by cortical 5-HT_{2A} receptor antagonism. These results indicate that stimulation of cortical 5-HT_{2A} receptors facilitates mesocortical DA release. They suggest that this effect may be mediated by the activation of corticotegmental glutamatergic projections that synapse on mesocortical DA neurons, although further work must be carried out to test this hypothesis. Importantly, blockade of cortical 5-HT_{2A} receptors or local perfusion with the antipsychotic drug risperidone also attenuated mild stress-induced DA release in the PFC. Thus, cortical 5-HT_{2A} receptors modulate the evoked release of DA in the mPFC, including that which is induced physiologically by stress. It is well known that schizophrenia is exacerbated by stress. Thus, the clinical efficacy of risperidone and other atypical antipsychotic treatments may be due, at least partially, to their ability to diminish stress-induced effects in the mPFC.

ACKNOWLEDGEMENTS

This work was supported by a MERIT Review award from the Department of Veterans Affairs, MH 52220 from NIH, and a grant from Janssen Extramural Research Foundation to EAP. BLR was supported by KO2MH01366, MH57635, MH61887, and the NIMH Psychoactive Drug Screening Program. We thank Marion Merrell Dow, Janssen Pharmaceutica, SmithKline and Beecham, and Sanofi Recherche for their generous contributions of drugs. We also thank Bobbi

Kirkbride and Indrani Sivagnanar for their excellent technical assistance.

REFERENCES

- Abercrombie ED, Keefe KA, DiFrischia DS, Zigmond MJ (1989). Differential effect of stress on *in vivo* dopamine release in striatum, nucleus accumbens, and medial prefrontal cortex. *J Neurochem* 52: 1655–1658.
- Abi-Dargham A, Gil R, Krystal J, Baldwin RM, Seibyl JP, Bowers M et al (1998). Increased striatal dopamine transmission in schizophrenia: confirmation in a second cohort. *Am J Psychiatry* 155: 761–767.
- Abi-Dargham A, Mawlawi O, Lombardo I, Gill R, Martinez D, Huang Y et al (2002). Prefrontal dopamine D1 receptors and working memory in schizophrenia. *J Neurosci* 22: 3708–3719.
- Aghajanian G, Marek G (1997). Serotonin induces excitatory postsynaptic potentials in apical dendrites of neocortical pyramidal cells. *Neuropharmacology* 36: 589–599.
- Alex KD, Yavarian GJ, McFarlane HG, Pluto CP, Pehek EA (2005). Modulation of dopamine release by striatal 5-HT_{2C} receptors. *Synapse* 55: 242–251.
- Arnsten AFT, Goldman-Rakic PS (1998). Noise stress impairs prefrontal cortical cognitive function in monkeys. *Arch Gen Psychiatry* 55: 362–368.
- Azmitia EC, Segal M (1978). An autoradiographic analysis of the differential ascending projections of the dorsal and median raphe nuclei in the rat. *J Compr Neurol* 179: 641–668.
- Barr AM, Lehmann-Masten V, Paulus M, Gainetdinov RR, Caron MG, Geyer MA (2004). The selective serotonin-2A receptor antagonist M100907 reverses behavioral deficits in dopamine transporter knockout mice. *Neuropsychopharmacology* 29: 221–228.
- Beart PM, McDonald D (1982). 5-Hydroxytryptamine and 5-hydroxytryptaminergic–dopaminergic interactions in the ventral tegmental area of rat brain. *J Pharm Pharmacol* 34: 591–593.
- Bonaccorso S, Meltzer H, Li Z, Dai J, Alboszta A, Ichikawa J (2002). SR46349-B, a 5-HT(2A/2C) receptor antagonist, potentiates haloperidol-induced dopamine release in rat medial prefrontal cortex and nucleus accumbens. *Neuropsychopharmacology* 27: 430–441.
- Bortolozzi A, az-Mataix LD, Scorza C, Celada P, Artigas F (2004). Atypical antipsychotics modulate a prefrontal 5-HT_{2A} receptor-mediated activation of the mesocortical dopaminergic pathway. Abstract Viewer/Itinerary Planner Washington, DC: Society for Neuroscience Program No. 45.10.
- Breier A, Su TP, Saunders R, Carson RE, Kolachana BS, de Bartolomeis A et al (1997). Schizophrenia is associated with elevated amphetamine-induced synaptic dopamine concentrations: evidence from a novel positron emission tomography method. *Proc Natl Acad Sci USA* 94: 2569–2574.
- Carr DB, Sesack SR (2000). GABA-containing neurons in the rat ventral tegmental area project to the prefrontal cortex. *Synapse* 38: 114–123.
- Cohen RM, Nordahl TE, Semple WE, Andreason P, Litman RE, Pickar D (1997). The brain metabolic patterns of clozapine- and fluphenazine-treated patients with schizophrenia during a continuous performance task. *Arch Gen Psychiatry* 54: 481–486.
- Davies MA, Compton-Toth BA, Hufeisen SJ, Meltzer HY, Roth BL (2005). The highly efficacious actions of *N*-desmethylozapine at muscarinic receptors are unique and not a common property of either typical or atypical antipsychotic drugs: is M(1) agonism a pre-requisite for mimicking clozapine's actions? *Psychopharmacology (Berl)* 178: 451–460.
- De Deurwaerdere P, Spampinato U (1999). Role of serotonin2A and serotonin2B/2C receptor subtypes in the control of accumbal and striatal dopamine release elicited *in vivo* by dorsal raphe nucleus electrical stimulation. *J Neurochem* 73: 1033–1042.
- Di Matteo V, De Blasi A, Di Giulio C, Esposito E (2001). Role of 5-HT_{2C} receptors in the control of central dopamine function. *Trends Pharmacol Sci* 22: 229–232.
- Doherty MD, Pickel VM (2000). Ultrastructural localization of the serotonin2A receptor in dopaminergic neurons in the ventral tegmental area. *Brain Res* 864: 176–185.
- Donzanti BA, Yamamoto BK (1988). An improved and rapid HPLC-ED method for the isocratic separation of amino acid neurotransmitters from brain tissue and microdialysis perfusates. *Life Sci* 43: 913–922.
- Feenstra MGP, Botterblom MHA (1996). Rapid sampling of extracellular dopamine in the rat prefrontal cortex during food consumption, handling and exposure to novelty. *Brain Res* 742: 17–24.
- Gainetdinov RR, Jones SR, Caron MG (1999). Functional hyperdopaminergia in dopamine transporter knock-out mice. *Biol Psychiatry* 46: 303–311.
- Gobert A, Millan MJ (1999). Serotonin (5-HT)_{2A} receptor activation enhances dialysate levels of dopamine and noradrenaline, but not 5-HT, in the rat frontal cortex of freely-moving rats. *Neuropharmacology* 38: 315–317.
- Gobert A, Rivet J, Lejeune F, Newman-Tancredi A, Adhumeau-Auclair A, Nicolas J et al (2000). Serotonin2C receptors tonically suppress the activity of mesocortical dopaminergic and adrenergic, but not serotonergic, pathways: a combined dialysis and electrophysiological analysis in the rat. *Synapse* 36: 205–221.
- Goldman-Rakic PS, Muly III EC, Williams GV (2000). D1 receptors in prefrontal cells and circuits. *Brain Res Rev* 31: 295–301.
- Gomeza J, Mary S, Brabet I, Parmentier M, Restituito S, Bockaert J et al (1996). Coupling of metabotropic glutamate receptors 2 and 4 to G alpha 15, G alpha 16, and chimeric G alpha q/i proteins: characterization of new antagonists. *Mol Pharmacol* 50: 923–930.
- Grace AA (1991). Phasic versus tonic dopamine release and the modulation of dopamine system responsivity: a hypothesis for the etiology of schizophrenia. *Neuroscience* 41: 1–24.
- Hertel P, Nomikos GG, Iurlo M, Svensson TH (1996). Risperidone: regional effects *in vivo* on release and metabolism of dopamine and serotonin in the rat brain. *Psychopharmacology* 124: 74–86.
- Ichikawa J, Ishii H, Bonaccorso S, Fowler WL, O'Laughlin IA, Meltzer HY (2001). 5-HT_{2A} and D₂ receptor blockade increases cortical DA release via 5-HT_{1A} receptor activation: a possible mechanism of atypical antipsychotic-induced cortical dopamine release. *J Neurochem* 76: 1521–1531.
- Ichikawa J, Kuroki T, Dai J, Meltzer HY (1998). Effect of antipsychotic drugs on extracellular serotonin levels in rat medial prefrontal cortex and nucleus accumbens. *Eur J Pharmacol* 351: 163–171.
- Inglis FM, Moghaddam B (1999). Dopaminergic innervation of the amygdala is highly responsive to stress. *J Neurochem* 72: 1088–1094.
- Jakab RL, Goldman-Rakic PS (1998). 5-Hydroxytryptamine(2A) serotonin receptors in the primate cerebral cortex: possible site of action of hallucinogenic and antipsychotic drugs in pyramidal cell apical dendrites. *Proc Natl Acad Sci USA* 95: 735–740.
- Jentsch JD, Roth RH (1999). The neuropsychopharmacology of phencyclidine: from NMDA receptor hypofunction to the dopamine hypothesis of schizophrenia. *Neuropsychopharmacology* 20: 201–225.
- Kawahara H, Yshida M, Hideyasu Y, Masakatsu N, Masatoshi T (1993). Psychological stress increases serotonin release in the rat amygdala and prefrontal cortex assessed by *in vivo* microdialysis. *Neurosci Lett* 162: 81–84.
- Kawahara Y, Kawahara H, Westerink BHC (1999). Comparison of effects of hypotension and handling stress on the release of noradrenaline and dopamine in the locus coeruleus and medial

- prefrontal cortex of the rat. *Naunyn-Schmiedeberg's Arch Pharmacol* **360**: 42–49.
- Kozikowski A, Steensma D, Araldi GL, Tuckmantel W, Wang S, Pshenichkin S et al (1998). Synthesis and biology of the conformationally restricted ACPD analogue, 2-aminobicyclo[2.1.1]hexane-2,5-dicarboxylic acid-I, a potent mGluR agonist. *J Med Chem* **41**: 1641–1650.
- Kuroki T, Meltzer HY, Ichikawa J (1999). Effects of antipsychotic drugs on extracellular dopamine levels in rat medial prefrontal cortex and nucleus accumbens. *J Pharmacol Exp Ther* **288**: 774–781.
- Lahti AC, Holcomb HH, Weiler MA, Medoff DR, Tamminga CA (2003). Functional effects of antipsychotic drugs: comparing clozapine with haloperidol. *Biol Psychiatry* **53**: 601–608.
- Laruelle M, Abi-Dargham A, Gil R, Kegeles L, Innis R (1999). Increased dopamine transmission in schizophrenia: relationship to illness phases. *Biol Psychiatry* **46**: 56–72.
- Liegeois J-F, Ichikawa J, Meltzer HY (2002). 5-HT_{2A} receptor antagonism potentiates haloperidol-induced dopamine release in rat medial prefrontal cortex and inhibits that in the nucleus accumbens in a dose-dependent manner. *Brain Res* **947**: 157–165.
- Martin-Ruiz R, Puig VM, Celada P, Shapiro DA, Roth BL, Mengod G et al (2001). Control of serotonin function in medial prefrontal cortex by serotonin-2A receptors through a glutamate-dependent mechanism. *J Neurosci* **21**: 9856–9866.
- Meltzer HY (1989). Clinical studies on the mechanism of action of clozapine: the dopamine-serotonin hypothesis of schizophrenia. *Psychopharmacology* **99**: S18–S27.
- Meltzer HY, Arvanitis L, Bauer D, Rein W (2004). Placebo-controlled evaluation of four novel compounds for the treatment of schizophrenia and schizoaffective disorder. *Am J Psychiatry* **161**: 975–984.
- Meltzer HY, Li Z, Kaneda Y, Ichikawa J (2003). Serotonin receptors: their key role in drugs to treat schizophrenia. *Prog Neuro-Psychopharm Biol Psychiatry* **27**: 1159–1172.
- Millan MJ, Dekeyne A, Gobert A (1998). Serotonin (5-HT)_{2C} receptors tonically inhibit dopamine (DA) and noradrenaline (NA), but not 5-HT, release in the frontal cortex *in vivo*. *Neuropharmacology* **37**: 953–955.
- Miller DD, Andreasen NC, O'Leary DS, Watkins GL, Boles Ponto LL, Hichawa RD (2001). Comparison of the effects of risperidone and haloperidol on regional cerebral blood flow in schizophrenia. *Biol Psychiatry* **49**: 704–715.
- Miner LAH, Backstrom JR, Sanders-Bush E, Sesack (2003). Ultrastructural localization of serotonin_{2A} receptors in the middle layers of the rat prelimbic prefrontal cortex. *Neuroscience* **116**: 107–117.
- Moghaddam B, Bunney BS (1990). Acute effects of typical and atypical antipsychotic drugs on the release of dopamine from prefrontal cortex, nucleus accumbens, and striatum of the rat: an *In Vivo* Microdialysis Study. *J Neurochem* **54**: 1755–1760.
- Molina V, Gispert JD, Reig S, Sanz J, Pascau J, Santos A et al (2005a). Cerebral metabolic changes induced by clozapine in schizophrenia and related to clinical improvement. *Psychopharmacology* **178**: 17–26.
- Molina V, Gispert JD, Reig S, Sanz J, Pascau J, Santos A et al (2003). Cerebral metabolism and risperidone treatment in schizophrenia. *Schizophr Res* **60**: 1–7.
- Molina V, Sanz J, Sarramea F, Benito C, Palomo T (2005b). Prefrontal atrophy in first episodes of schizophrenia associated with limbic metabolic hyperactivity. *J Psychiatr Res* **39**: 117–127.
- Nocjar C, Roth BL, Pehek EA (2002). Localization of 5-HT_{2A} receptors on dopamine cells in subnuclei of the midbrain A10 cell group. *Neuroscience* **111**: 163–176.
- Obrenovitch TP, Urenjak J, Richards DA, Ueda Y, Curzon G, Symon L (1993). Extracellular neuroactive amino acids in the rat striatum during ischaemia: comparison between penumbral conditions and ischaemia with sustained anoxic depolarisation. *J Neurochem* **61**: 178–186.
- Ojima T, Ito C, Sakurai E, Sakurai E, Watanabe T, Yanai K (2004). Effects of serotonin-dopamine antagonists on prepulse inhibition and neurotransmitter contents in the rat cortex. *Neurosci Lett* **366**: 130–134.
- Pan WHT, Yang S-Y, Lin S-K (2004). Neurochemical interaction between dopaminergic and noradrenergic neurons in the medial prefrontal cortex. *Synapse* **53**: 44–52.
- Paxinos G, Watson C. (1998). *The Rat Brain in Stereotaxic Coordinates*. Academic Press: New York.
- Pehek EA, McFarlane EA, Maguschak K, Price B, Pluto CP (2001). M100,907, a selective 5-HT_{2A} antagonist, attenuates dopamine release in the rat medial prefrontal cortex. *Brain Res* **888**: 51–59.
- Pehek EA, Yamamoto BK (1994). Differential effects of locally administered clozapine and haloperidol on dopamine efflux in the rat prefrontal cortex and caudate-putamen. *J Neurochem* **63**: 2118–2124.
- Potkin SG, Basile VS, Jin Y, Masellis M, Badri F, Keator D et al (2003). D1 receptor alleles predict PET metabolic correlates of clinical response to clozapine. *Mol Psychiatry* **8**: 109–113.
- Pozzi L, Accancia S, Ceglia I, Invernizzi RW, Smamanin R (2002). Stimulation of 5-hydroxytryptamine (5HT_{2C}) receptors in the ventrosegmental area inhibits stress-induced but not basal dopamine release in the rat prefrontal cortex. *J Neurochem* **82**: 93–100.
- Roth B, Hamblin M, Ciaranello R (1991). Developmental regulation of 5-HT₂ and 5-HT_{1c} mRNA and receptor levels. *Brain Res Dev Brain Res* **58**: 51–58.
- Roth B, Meltzer H, Khan N (1998). Binding of typical and atypical antipsychotic drugs to multiple neurotransmitter receptors. *Adv Pharmacol* **42**: 482–485.
- Roth BL, Baner K, Westkaemper R, Siebert D, Rice KC, Steinberg S et al (2002). Salvinorin A: a potent naturally occurring nonnitrogenous kappa opioid selective agonist. *Proc Natl Acad Sci USA* **99**: 11934–11939.
- Roth BL, Ernsberger P, Steinberg S, Rao S, Rauser L, Savage J et al (2001). The *in vitro* pharmacology of the β -adrenergic receptor pet ligand (s)-fluorocarazolol reveals high affinity for cloned β -adrenergic receptors and moderate affinity for the human 5-HT_{1A} receptor. *Psychopharmacology* **157**: 111–114.
- Roth BL, Sheffler DJ, Kroeze WK (2004). Magic shotguns versus magic bullets: selectively non-selective drugs for mood disorders and schizophrenia. *Natl Rev Drug Discov* **3**: 353–359.
- Rothman RB, Baumann MH, Savage JE, Rauser L, McBride A, Hufeisen SJ et al (2000). Evidence for possible involvement of 5-HT_{2B} receptors in the cardiac valvulopathy associated with fenfluramine and other serotonergic medications. *Circulation* **102**: 2836–2841.
- Sesack SR, Pickel VM (1992). Prefrontal cortical efferents in the rat synapse on Unlabeled neuronal targets of catecholamine terminals in the nucleus accumbens septi and on dopamine neurons in the ventral tegmental area. *J Comp Neurol* **320**: 145–160.
- Shapiro DA, Kristiansen K, Weiner DM, Kroeze WK, Roth BL (2002). Evidence for a model of agonist-induced activation of 5-hydroxytryptamine 2A serotonin receptors that involves the disruption of a strong ionic interaction between helices 3 and 6. *J Biol Chem* **277**: 11441–11449.
- Shi Q, Savage JE, Hufeisen SJ, Rauser L, Grajkowska E, Ernsberger P et al (2003). L-homocysteine sulfinic acid and other acidic homocysteine derivatives are potent and selective metabotropic glutamate receptor agonists. *J Pharmacol Exp Ther* **305**: 131–142.
- Sorg B, Kalivas P (1993). Effects of cocaine and footshock stress on extracellular dopamine levels in the medial prefrontal cortex. *Neuroscience* **53**: 695–703.
- Talvik-Lotfi M, Nyberg S, Nordstrom A-L, Ito H, Halldin C, Brunner F et al (2000). High 5-HT_{2A} receptor occupancy in

- M100907-treated schizophrenic patients. *Psychopharmacology* **148**: 400–403.
- Thierry AM, Tassin JP, Blanc G, Glowinski J (1976). Selective activation of the mesocortical DA system by stress. *Nature* **263**: 242–244.
- Weinberger DR (1987). Implications of normal brain development for the pathogenesis of schizophrenia. *Arch Gen Psychiatry* **44**: 660–669.
- Westerink BHC, de Boer P, de Vries JB, Kruse CG, Long SK (1998). Antipsychotic drugs induce similar effects on the release of dopamine and noradrenaline in the medial prefrontal cortex of the rat brain. *Eur J Pharm* **361**: 27–33.
- Westerink BHC, Kawahara Y, De Boer P, Geels C, De Vries JB, Wikstrom HV et al (2001). Antipsychotic drugs classified by their effects on the release of dopamine and noradrenaline in the prefrontal cortex and striatum. *Eur J Pharmacol* **412**: 127–138.
- Williams G, Goldman-Rakic P (1995). Modulation of memory fields by dopamine D1 receptors in prefrontal cortex. *Nature (London)* **376**: 572–575.
- Willins DL, Deutch AY, Roth BL (1997). Serotonin 5-HT_{2A} receptors are expressed on pyramidal cells and interneurons in the rat cortex. *Synapse* **27**: 79–82.
- Wroblewska B, Wroblewski JT, Pshenichkin S, Surin A, Sullivan SE, Neale JH (1997). N-acetylaspartylglutamate selectively activates mGluR3 Receptors in transfected cells. *J Neurochem* **69**: 174–181.
- Xue C-J, Ng JP, Li Y, Wolf ME (1996). Acute and repeated systemic amphetamine administration: effects on extracellular glutamate, aspartate, and serine levels in rat ventral tegmental area and nucleus accumbens. *J Neurochem* **67**: 352–363.
- Yamamoto BK, Pehek EA (1990). A neurochemical heterogeneity of the rat striatum as measured by *in vivo* electrochemistry and microdialysis. *Brain Res* **506**: 236–242.
- Zahrt J, Taylor JR, Mathew RG, Arnsten AFT (1997). Supranormal stimulation of D1 dopamine receptors in the rodent prefrontal cortex impairs spatial working memory performance. *J Neurosci* **17**: 8528–8535.
- Zhang W, Perry KW, Wong DT, Potts BD, Bao J, Tollefson GD et al (2000). Synergistic effects of olanzapine and other antipsychotic agents in combination with fluoxetine on norepinephrine and dopamine release in rat prefrontal cortex. *Neuropsychopharmacology* **23**: 250–262.