

# Effect of Ginseng Saponins on Enhanced Dopaminergic Transmission and Locomotor Hyperactivity Induced by Nicotine

Sang Eun Kim<sup>\*1</sup>, Insop Shim<sup>2</sup>, June-Key Chung<sup>1</sup> and Myung Chul Lee<sup>1</sup>

<sup>1</sup>Department of Nuclear Medicine, Seoul National University College of Medicine, Seoul, Korea; <sup>2</sup>Department of Integrative Medicine, College of Medicine, The Catholic University of Korea, Seoul, Korea

Several studies have shown that behavioral hyperactivity induced by psychomotor stimulants is prevented by ginseng saponins. In an attempt to investigate whether the effect of ginseng saponins is through their inhibitory action on the enhanced dopaminergic transmission by psychomotor stimulants, we examined the effects of ginseng total saponin (GTS) presynaptically on nicotine-induced dopamine (DA) release in the striatum of freely moving rats using *in vivo* microdialysis technique and postsynaptically on the *in vitro* and *in vivo* binding of [<sup>3</sup>H]raclopride to DA D<sub>2</sub> receptors. Also, we examined the effects of GTS on nicotine-induced locomotor hyperactivity and on nicotine-induced Fos protein expression in the nucleus accumbens and striatum. Systemic pretreatment with GTS (100 and 400 mg/kg, intraperitoneally (i.p.)) resulted in a dose-dependent inhibition of locomotor hyperactivity induced by nicotine. GTS decreased nicotine-induced DA release in the striatum in a dose-dependent manner. However, GTS had no effects on resting levels of locomotor activity and extracellular DA in the striatum. GTS inhibited the *in vitro* binding of [<sup>3</sup>H]raclopride to rat striatal membranes with an IC<sub>50</sub> of 5.14 ± 1.09 μM. High doses of GTS (400 and 800 mg/kg, i.p.) resulted in decreases in the *in vivo* binding of [<sup>3</sup>H]raclopride in the striatum. GTS decreased nicotine-induced Fos protein expression in the nucleus accumbens and striatum, reflecting the inhibition by GTS of nicotine-induced enhancement of dopaminergic transmission. The results of the present study suggest that GTS acts not only on dopaminergic neurons directly or indirectly to prevent nicotine-induced DA release but also postsynaptically by binding to DA D<sub>2</sub> receptors. This may explain the blocking effect of GTS on behavioral activation induced by nicotine and conceivably by other psychostimulants. Our data raise the possibility that GTS, by attenuating nicotine-induced enhancement of dopaminergic transmission, may prove to be a useful therapeutic agent for nicotine addiction and warrant further investigation on its effect on nicotine's rewarding property.

*Neuropsychopharmacology* (2006) **31**, 1714–1721. doi:10.1038/sj.npp.1300945; published online 26 October 2005

**Keywords:** ginseng total saponin; nicotine; dopamine release; dopamine D<sub>2</sub> receptor; Fos protein; locomotor activity

## INTRODUCTION

Evidence suggests that nicotine, the neuroactive compound of tobacco, has common neurochemical as well as behavioral properties with other addictive drugs such as cocaine and amphetamine (Henningfield and Heishman, 1995; Pontieri *et al*, 1996). It is well documented that nicotine stimulates dopamine (DA) release in the nucleus accumbens and striatum (Imperato *et al*, 1986; Damsma *et al*, 1989; Toth *et al*, 1992; Benwell and Balfour, 1994;

Nisell *et al*, 1994a; Pontieri *et al*, 1996; Mirza *et al*, 1996; Marshall *et al*, 1997). This property has been related to the addictive and behavioral properties of nicotine (Benwell and Balfour, 1992; Corrigan *et al*, 1992; Di Chiara and Imperato, 1988).

Ginseng, the root of *Panax ginseng* CA Meyer, has been used widely as a herbal medicine across the world. Many investigators have reported its chemical and pharmacological properties and saponin appears to be responsible for most of the pharmacological effects of ginseng. Behavioral studies have suggested that ginseng saponins or ginsenosides may act on the central dopaminergic system. Pretreatment with ginseng saponins or ginseng extract was able to block the development of behavioral sensitization induced by cocaine (Kim *et al*, 1995a), methamphetamine (Tokuyama *et al*, 1992), and morphine (Kim *et al*, 1995b). The conditioned place preference induced by these drugs was also antagonized by ginseng saponins (Kim *et al*,

\*Correspondence: Dr SE Kim, Department of Nuclear Medicine, Seoul National University Bundang Hospital, 300 Gumi-dong, Bundang-gu, Seongnam-si, Gyeonggi-do, 463-707, Korea, Tel: +82 31 787 7671, Fax: +82 31 787 4018, E-mail: kse@snu.ac.kr

Received 18 March 2005; revised 24 August 2005; accepted 15 September 2005

Online publication: 20 September 2005 at <http://www.acnp.org/citations/Npp092005050190/default.pdf>

1996a,b). Also, it has been shown that ginseng saponins inhibit nicotine-induced hyperlocomotion, behavioral sensitization, and conditioned place preference (Kim and Kim, 1999). However, the neurochemical mechanisms underlying the behavioral effects of ginseng saponins are not well understood.

Convergent evidence suggests that drugs of abuse exert their locomotor stimulating and rewarding effects by increasing dopaminergic transmission. Therefore, it seems possible that the behavioral effects of ginseng saponins reflect their blockade of enhanced dopaminergic transmission by those drugs. In an attempt to examine this hypothesis, we investigated the following: (1) the effect of ginseng total saponin (GTS) on nicotine-induced locomotor hyperactivity; (2) the effect of GTS presynaptically on nicotine-induced DA release in the striatum of freely moving rats using *in vivo* microdialysis technique; (3) the effect of GTS postsynaptically on the *in vitro* and *in vivo* binding of [<sup>3</sup>H]raclopride to DA D<sub>2</sub> receptors; and (4) the effect of GTS on nicotine-induced Fos protein expression in the nucleus accumbens and striatum.

## MATERIALS AND METHODS

### Animals

Male Sprague–Dawley rats, weighing 280–320 g, were used for *in vivo* microdialysis, behavioral, *in vitro* receptor binding, and immunohistochemical studies. Male ICR mice (25–30 g) were used for *in vivo* binding studies. The animals were group-housed in light-, temperature-, and humidity-controlled animal quarters with food and water available *ad libitum*. All procedures involving animal use were in strict accordance with the NIH Guide for the Care and Use of Laboratory Animals and were approved by the Seoul National University Hospital Animal Care and Use Committee.

### Drugs and Chemicals

(–)-Nicotine hydrogen tartrate was purchased from Sigma Chemical Co. (St Louis, MO). GTS was a gift from Korea Ginseng and Tobacco Research Institute (Daejeon, Korea). GTS (a characterized saponin mixture quantitatively containing nine major glycosides known as ginsenosides (Rb<sub>1</sub> 20.14%, Rb<sub>2</sub> 10.19%, Rc 11.34%, Rd 4.63%, Re 12.27%, Rf 3.01%, Rg<sub>1</sub> 16.44%, Rg<sub>2</sub> 2.01%, and Rg<sub>3</sub> 2.64%) and other minor ginsenosides and components (17.33%), according to an HPLC method of Ko *et al* (1992), from *P. ginseng*) was extracted and purified using the method of Ando *et al* (1971). [<sup>3</sup>H]raclopride (79.3 Ci/mmol) was purchased from New England Nuclear (Boston, MA).

### Surgery and Microdialysis Procedures

Rats were anesthetized with sodium pentobarbital (50 mg/kg, intraperitoneally (i.p.)). Using aseptic techniques, a guide cannula (CMA/12, CMA Microdialysis, Solna, Sweden) aimed to terminate in the dorsal striatum (AP 1.0, L 3.2 from bregma; H 3.0 from dura) (Paxinos and Watson, 1986) was stereotactically implanted and attached to the skull

using skull screws and dental cement. The cannula was then closed with a tight-fitting stainless-steel obturator. The microdialysis was performed in freely moving rats. Following 3-day recovery, a 4 mm microdialysis probe (CMA/12, CMA Microdialysis) connected via a dual liquid swivel to a syringe pump was inserted into the guide cannula and perfused with an artificial cerebrospinal fluid (145 mM NaCl, 2.7 mM KCl, 1.2 mM CaCl<sub>2</sub>, 1.0 mM MgCl<sub>2</sub>, 2.0 mM Na<sub>2</sub>HPO<sub>4</sub>; pH 7.4) at a constant rate of 1.5 µl/min. Dialysate samples were collected during 20 min sampling intervals via outlet tubing connected to a microfraction collector (CMA Microdialysis). The position of the probes was verified by histological examination at the end of experiments.

### Analytical Procedures

The dialysate (injection volume 30 µl) was assayed for DA using HPLC coupled to an ESA Coulochem II 5200A electrochemical detection system with an oxidation potential of +320 mV. The detector was equipped with a high-performance analytical cell (ESA model 5014), which is tailored for use in microdialysis applications. The mobile phase was composed of 75 mM monobasic sodium phosphate, 0.1 mM EDTA, 1.4 mM octanesulfonic acid and 10% acetonitrile, and adjusted to pH 3.2 with HPLC grade phosphoric acid. The separation of monoamine metabolites was performed on a Waters Nova-Pak C-18 column (4 µm, 150 × 3.9 mm). The flow rate of the system was 1.0 ml/min. DA in dialysates was expressed as a percentage of three baseline samples collected immediately before nicotine treatment.

### Drug Treatments

Nicotine was dissolved in the artificial cerebrospinal fluid to the concentration of 10 mM (expressed as free base) and infused for 60 min locally into the striatum through the dialysis probe. GTS (100 and 400 mg/kg, dissolved in saline) was injected i.p. 60 min before nicotine was administered.

### Measurement of Locomotor Activity

The effect of GTS on nicotine-induced behavioral hyperactivity was evaluated in rats by measuring locomotor activity using a stabilimeter interfaced to a personal computer system. The stabilimeter, in which a loudspeaker is used both as platform and transducer, was constructed based on the design described by Parreno *et al* (1985). In our preliminary studies, this equipment provided a good reproducibility and a high sensitivity for measuring locomotor activity of small animals. Each animal was placed on the stabilimeter and after a habituation period of 60 min followed by the measurement of baseline activity for 60 min, nicotine (10 mM) was infused for 60 min locally into the striatum through the dialysis probe. Control animals were implanted with the dialysis probe and infused with the artificial cerebrospinal fluid. Locomotor activity was measured during 20 min intervals for a total of 180 min after the administration of nicotine. GTS (100 and 400 mg/kg) was injected i.p. 60 min prior to nicotine.

### *In Vitro* Inhibition Experiments

Inhibition by GTS of the *in vitro* binding of [ $^3\text{H}$ ]raclopride to striatal DA  $\text{D}_2$  receptors was investigated. Rats were killed by rapid decapitation. The striata were quickly dissected, frozen on a block of dry ice, and stored at  $-70^\circ\text{C}$  until needed. The binding of [ $^3\text{H}$ ]raclopride to rat striatal membranes was conducted as described by Lidow *et al* (1989). Briefly, tissues were homogenized in ice-cold 50 mM Tris · HCl buffer (pH 7.4) and centrifuged at 45 000 *g* for 10 min. The resulting pellet was resuspended in 20 volumes of buffer, recentrifuged, and resuspended at 10 mg wet weight/ml of incubation buffer (50 mM Tris · HCl buffer (pH 7.4) containing 120 mM NaCl, 5 mM KCl, 2 mM  $\text{CaCl}_2$ , and 1 mM  $\text{MgCl}_2$ ). Inhibition assays of [ $^3\text{H}$ ]raclopride were performed using a fixed concentration of [ $^3\text{H}$ ]raclopride (4.3 nM) and increasing concentrations of GTS (10 pM to 10 mM). The incubations were conducted in a final volume of 250  $\mu\text{l}$  for 30 min at 20–22°C. The incubation was terminated by rapid filtration through Whatman GF/B filters previously soaked in 0.05% polyethylenimine using a Brandel MP-48LT filtering manifold (Gaithersburg, MD). The radioactivity trapped in the filters was measured in a liquid scintillation counter (Tri-Carb 2500 TR, Packard Instrument Co., Meriden, CT) after addition of 5 ml Aquassure scintillator (Packard). Nonspecific binding was defined using 1  $\mu\text{M}$  (+)-butaclamol.

### *In Vivo* Binding Experiments

The effect of GTS on the *in vivo* binding of [ $^3\text{H}$ ]raclopride was investigated by pretreating mice with different doses of GTS (50–800 mg/kg, i.p.) 60 min before injection of [ $^3\text{H}$ ]raclopride. [ $^3\text{H}$ ]raclopride (1  $\mu\text{Ci}$  in 0.2 ml saline) was injected intravenously into a tail vein, and 15 min later, the animals were killed by cervical dislocation. Different brain regions were immediately dissected on ice, weighed and placed into glass vials. After digestion of the tissues with Soluene-350 (Packard), 10 ml of Formula 989 scintillator (Packard) was added and the samples counted in a liquid scintillation counter.

### Fos Immunohistochemistry

The effect of GTS on nicotine-induced Fos protein expression in dopaminergic target areas was examined. Rats received subcutaneous (s.c.) injection of nicotine (0.4 mg/kg; expressed as free base) 2 h before anesthesia (ketamine 80 mg/kg, i.p. plus xylazine 10 mg/kg, i.p.). GTS (100 mg/kg) was administered i.p. 60 min before nicotine injection. Rats were transcardially perfused with ice-cold 0.1 M phosphate buffered saline (PBS), pH 7.4 followed by 4% paraformaldehyde in 0.1 M PBS. The brains were removed and post-fixed with the same fixative for 2 h at room temperature and immersed in a 20% sucrose solution overnight at 4°C. The frozen tissue was cut into 30  $\mu\text{m}$  thick sections on a cryostat. Coronal sections were rinsed in PBS (2  $\times$  5 min) and placed in 0.3%  $\text{H}_2\text{O}_2$  in methanol for 15 min and then washed in PBS (3  $\times$  5 min). The sections were preincubated in a 2% normal goat serum (DAKO, Glostrup, Denmark) in PBS-T (0.1% Triton X-100 in PBS) for 40 min to block nonspecific staining and then incubated with the

primary antibody (rabbit polyclonal Fos antibody, Oncogene Research, Cambridge, MA) diluted in 1:1000 for 48 h at 4°C. The sections were washed in PBS-T (3  $\times$  5 min) and incubated for 1 h with biotinylated goat anti-rabbit secondary antibody (Vector Laboratories, Burlingame, CA). After washing with PBS (3  $\times$  5 min), the sections were incubated for 1 h with PBS-T containing avidin-biotinylated horseradish peroxidase complex (1:100, Vector Laboratories). The sections were washed in PBS (3  $\times$  5 min) and then rinsed for 10 min in 0.175 M sodium acetate buffer (pH 7.5). The immunoreactive complex was revealed using diaminobenzidine (DAB) and nickel sulfate ( $\text{NiSO}_4$ ) (0.2 mg/ml DAB, 25 mg/ml  $\text{NiSO}_4$  and 0.83  $\mu\text{l}/\text{ml}$  3%  $\text{H}_2\text{O}_2$  in 0.175 M sodium acetate) for 4 min. Sections were washed with PBS twice and mounted on silane-coated slides, air-dried, dehydrated through graded ethanols into xylene, and coverslipped with Permount (Fisher Scientific, Pittsburgh, PA). No Fos-like immunoreactivity has been observed when the primary antibody was omitted. The number of Fos-positive nuclei were counted by using the IBAS image analysis system (Zeiss, Germany) within a squared field area of 210  $\times$  210  $\mu\text{m}$  for the nucleus accumbens and 420  $\times$  420  $\mu\text{m}$  for the striatum using  $\times$  200 magnifications. The mean value of the three to six sections was calculated from each selected region.

### Statistical Analysis

Unless otherwise stated, values are expressed as mean  $\pm$  SEM. Data were analyzed by one-way analysis of variance (ANOVA) or two-way ANOVA (drug treatment  $\times$  time) with repeated measures on the time factor. The origin of significant effects was further examined by *post hoc* comparisons using the Bonferroni technique.


## RESULTS

### Effect of GTS on Nicotine-Induced Locomotor Hyperactivity


Figure 1 shows the effect of GTS on nicotine-induced hyperlocomotion. Local infusion of nicotine (10 mM) into the striatum via the dialysis probe increased the locomotor activity (maximal response = 589% of controls). This effect was attenuated by systemic pretreatment with GTS. An ANOVA indicated a significant effect of drug treatments ( $F_{3,12} = 8.7$ ,  $p < 0.005$ ). *Post hoc* comparisons revealed that GTS attenuated the nicotine-induced hyperlocomotion in a dose-dependent manner. GTS (100 and 400 mg/kg, i.p.) decreased the nicotine-induced maximal locomotor activity by 37% ( $p < 0.01$ ) and 58% ( $p < 0.005$ ), respectively. However, GTS (100 and 400 mg/kg, i.p.) had no effect on resting levels of locomotor activity until 180 min after injection ( $F_{2,9} = 0.7$ ,  $p = \text{NS}$ ) (Figure 3).

### Effect of GTS on Nicotine-Induced DA Release

Figure 2 shows the effect of GTS on nicotine-induced changes in extracellular DA in the striatum. Local infusion of nicotine (10 mM) into the striatum produced an increase in extracellular DA in the striatum (maximal response =  $500 \pm 78\%$  of basal levels). This effect was




**Figure 1** Effect of GTS on nicotine-induced hyperlocomotion. Nicotine (10 mM) was infused intrastrially (i.s.) for 60 min beginning at time 0. GTS (100 and 400 mg/kg) was administered i.p. 60 min before nicotine infusion. Results are means  $\pm$  SEM of four independent experiments. \* $p$  < 0.05, \*\* $p$  < 0.005, \*\*\* $p$  < 0.001 vs the control group; # $p$  < 0.05, ## $p$  < 0.01, ### $p$  < 0.005 vs the saline plus nicotine group.




**Figure 2** Effect of GTS on nicotine-induced changes in extracellular DA in the striatum. Nicotine (10 mM) was infused intrastrially (i.s.) for 60 min beginning at time 0. GTS (100 and 400 mg/kg) was administered i.p. 60 min before nicotine infusion. Results are expressed as a percentage of three baseline samples and are means  $\pm$  SEM of 4–6 independent experiments. \* $p$  < 0.05, \*\* $p$  < 0.005, \*\*\* $p$  < 0.001 vs the control group; # $p$  < 0.05, ## $p$  < 0.005, ### $p$  < 0.001 vs the saline plus nicotine group.


attenuated by GTS. An ANOVA indicated a significant effect of drug treatments ( $F_{3,16} = 9.2$ ,  $p$  < 0.001). *Post hoc* comparisons revealed that GTS attenuated the nicotine-induced DA release in a dose-dependent manner. GTS (100 and 400 mg/kg, i.p.) inhibited the nicotine-induced maximal DA response by 58% ( $p$  < 0.005) and 76% ( $p$  < 0.001), respectively. However, GTS (100 and 400 mg/kg, i.p.) had no effect on resting levels of extracellular DA in the striatum until 180 min after injection ( $F_{2,11} = 0.9$ ,  $p$  = NS) (Figure 3).




**Figure 3** Effect of GTS on resting levels of locomotor activity and extracellular DA in the striatum. GTS (100 and 400 mg/kg) was administered i.p. at time 0. There were no significant differences in resting levels of locomotor activity and extracellular DA between control and GTS-treated animals. Results are means  $\pm$  SEM of 4–5 independent experiments.



**Figure 4** Inhibition of specific [<sup>3</sup>H]raclopride binding by GTS in rat striatal membranes. Presented are the results of a representative experiment; points are the means of triplicate determinations.



**Figure 5** Effect of GTS on the *in vivo* binding of [<sup>3</sup>H]raclopride in brain regions. GTS (50–800 mg/kg) was administered i.p. 90 min before intravenous injection of [<sup>3</sup>H]raclopride. *In vivo* [<sup>3</sup>H]raclopride binding was measured 15 min after radiotracer injection. Results are expressed as ratios of specific-to-nonspecific binding (region-to-cerebellar ratio - 1) and are means  $\pm$  SD;  $n$  = 4 for each group. \* $p$  < 0.0001.



**Figure 6** Effect of GTS on nicotine-induced Fos protein expression in the nucleus accumbens and striatum, with schematic drawings indicating areas in which Fos-positive nuclei were counted. GTS (100 mg/kg) was administered i.p. 60 min before nicotine injection (0.4 mg/kg, s.c.). Results are means  $\pm$  SEM of four independent experiments. \* $p < 0.05$  vs the saline plus saline group, # $p < 0.05$  vs the saline plus nicotine group.

#### Effect of GTS on the *In Vitro* Binding of [ $^3$ H]raclopride

GTS inhibited the binding of [ $^3$ H]raclopride to striatal membranes with an  $IC_{50}$  of  $5.14 \pm 1.09 \mu M$  (mean  $\pm$  SEM of five independent experiments) (Figure 4).

#### Effect of GTS on the *In Vivo* Binding of [ $^3$ H]raclopride

Figure 5 shows the effect of GTS on the *in vivo* binding of [ $^3$ H]raclopride in different brain regions. Doses of 50–200 mg/kg GTS resulted in 11–15%, but not significant decreases in the *in vivo* binding of [ $^3$ H]raclopride in the striatum. Increasing the dose of GTS to 400 and 800 mg/kg led to abrupt decreases in striatal [ $^3$ H]raclopride binding by 55% ( $p < 0.0001$ ) and 53% ( $p < 0.0001$ ), respectively. Regional analysis of the binding data indicated that the maximal effect of GTS on the *in vivo* binding of [ $^3$ H]raclopride occurred in the striatum, a region rich in DA  $D_2$  receptors.

#### Effect of GTS on Nicotine-Induced Fos Protein Expression

Figure 6 shows the effect of GTS on nicotine-induced Fos protein expression in dopaminergic target areas. Nicotine (0.4 mg/kg, s.c.) increased the number of Fos-positive nuclei in the nucleus accumbens and striatum by 89% ( $p < 0.05$ ) and 73% ( $p < 0.05$ ), respectively. GTS (100 mg/kg, i.p.) decreased the nicotine-induced Fos protein expression in the nucleus accumbens and striatum by 32% ( $p < 0.05$ ) and 36% ( $p < 0.05$ ), respectively. GTS did not influence resting levels of Fos-positive nuclei in the nucleus accumbens and striatum (see Figure 6).

#### DISCUSSION

This study demonstrated that systemic pretreatment with GTS inhibits nicotine-induced hyperlocomotion and striatal DA release in a dose-dependent manner. As behavioral hyperactivity produced by nicotine is associated with an increase in DA release in the striatum as well as in the nucleus accumbens (Shim *et al*, 2001), the inhibitory action of GTS on nicotine-induced DA release in the striatum explains, at least in part, its blocking effect on behavioral activation produced by nicotine. We also found that GTS binds, with a modest affinity, to DA  $D_2$  receptors both *in vitro* and *in vivo*. Because GTS had no effect on resting levels of extracellular DA, the inhibition by GTS of *in vivo* binding of [ $^3$ H]raclopride is not due to changes in synaptic DA levels. We do not know the function of GTS at DA  $D_2$  receptor sites. In the present study, the dose-dependent effect of GTS against intrastriatal infusion of nicotine was greater for the reduction of locomotor activity (57% greater reduction with GTS 400 mg/kg compared with GTS 100 mg/kg) than for the reduction of striatal DA release (31% greater reduction with GTS 400 mg/kg compared with GTS 100 mg/kg). Although it is not certain whether the magnitude of locomotor activity changes proportionally to that of DA release, this finding suggests that mechanisms other than DA release inhibition may be involved in the behavioral effect of high-dose GTS. This idea is supported by our finding that high doses of GTS (400 and 800 mg/kg) inhibited the *in vivo* binding of [ $^3$ H]raclopride to DA  $D_2$  receptors in the striatum. Also, it has been reported that GTS showed an antidopaminergic action at postsynaptic DA receptor sites by inhibiting apomorphine-induced climbing behavior (Kim *et al*, 1996a). In addition, high-dose ginseng

extract inhibited adenylate cyclase activity in rat brain tissues (Petkov, 1978). Taken together, it is likely that GTS may act on DA D<sub>2</sub> receptors as an antagonist. The results of the present study suggest that GTS acts not only on dopaminergic neurons directly or indirectly to prevent nicotine-induced DA release but also postsynaptically by binding to DA D<sub>2</sub> receptors. This may explain the blocking effect of GTS on behavioral activation induced by nicotine.

As nicotine increases DA release by stimulating nicotinic acetylcholine receptors (nAChRs) located in the terminal and somatodendritic regions of the dopaminergic neurons (Marshall *et al*, 1997; Mifsud *et al*, 1989; Wonnacott *et al*, 1990; Nisell *et al*, 1994a,b), it is possible that GTS inhibits DA release by acting on nAChRs in the dopaminergic neurons. Supporting this idea, it has been reported that ginseng saponins reduced acetylcholine-evoked Na<sup>+</sup> influx and catecholamine secretion in bovine adrenal chromaffin cells, but did not affect catecholamine secretion induced by high K<sup>+</sup> (an activator of voltage-sensitive Ca<sup>2+</sup> channels) or veratridine (an activator of voltage-sensitive Na<sup>+</sup> channels) (Tachikawa *et al*, 1995). Also, it has been shown that GTS had no effect on DA release induced by intrastriatal infusion of high K<sup>+</sup> solution (Shim *et al*, 2000). These data suggest that GTS blocks the nAChRs or the receptor-operated Na<sup>+</sup> channels (but not voltage-sensitive Na<sup>+</sup> and Ca<sup>2+</sup> channels), inhibit Na<sup>+</sup> influx through the channels and consequently reduce both Ca<sup>2+</sup> influx and DA release. In our preliminary studies, GTS did not affect the binding of [<sup>3</sup>H]nicotine to rat striatal membranes. This suggests that GTS may act on the receptor-operated ion channels rather than nAChRs or on the nAChR subtypes not labeled by [<sup>3</sup>H]nicotine. However, there is possibility that GTS acts on GABAergic and glutamatergic neurons, receptors of the neurotransmitters, or nAChRs located in these neurons, because these neurons interact with dopaminergic neurons in the mesoaccumbens and nigrostriatal systems to modulate DA release (Mansvelder and McGehee, 2002; Smolders *et al*, 1995). Indeed, several studies provided evidence that ginsenosides may regulate the GABA<sub>A</sub> receptor. For example, ginsenosides differentially regulated [<sup>3</sup>H]flunitrazepam and [<sup>3</sup>H]muscimol binding to the GABA<sub>A</sub> receptor in a rat brain membrane fraction (Kimura *et al*, 1994). Prolonged infusion with ginsenoside Rc but not with ginsenoside Rg<sub>1</sub> into rat brain elevated [<sup>3</sup>H]muscimol binding to the GABA<sub>A</sub> receptor in a region-specific manner (Kim *et al*, 2001). In addition, Tsang *et al* (1985) reported that ginseng saponins inhibited uptake of GABA in rat brain synaptosomes. Also, it has been shown that ginsenosides attenuate glutamate-induced injury of neuronal and non-neuronal cells in the brain (Abe *et al*, 1994; Seong *et al*, 1995; Kim *et al*, 2002).

Ginseng saponins were able to attenuate behavioral activation induced not only by nicotine but also by other psychostimulants, such as cocaine (Kim *et al*, 1995a), methamphetamine (Tokuyama *et al*, 1992), and morphine (Kim *et al*, 1995b). Interestingly, we found that GTS also inhibited cocaine-induced increase in extracellular DA without affecting [<sup>3</sup>H]WIN 35 428 binding to the DA transporter in rat striatum (data not shown). In other words, GTS attenuated the behavioral and extracellular DA increasing effects of psychostimulants that behave with different mechanisms (ie, increasing DA release (nicotine)

vs blocking DA transporters (cocaine)). This suggests that GTS may act on the final common pathways of the negative feedback loop for the regulation of dopaminergic neurotransmission.

Drug treatments affecting dopaminergic transmission modulate the expression of Fos protein encoded by *c-fos* proto-oncogene in the striatum (Graybiel *et al*, 1990; Robertson *et al*, 1990; Herrera and Robertson, 1996; Moratalla *et al*, 1996). Acute nicotine elevates Fos expression in various brain regions, including dopaminergic target areas (Ren and Sagar, 1992; Matta *et al*, 1993; Pang *et al*, 1993; Kiba and Jayaraman, 1994; Panagis *et al*, 1996; Salminen *et al*, 1996; Valentine *et al*, 1996). In the present study, GTS decreased nicotine-induced Fos protein expression in the nucleus accumbens and striatum. This reflects the inhibition of nicotine-induced enhancement of dopaminergic transmission by GTS. From our results, the effect of GTS may be due in large measure to its inhibitory effect on DA release because DA D<sub>1</sub> receptor, but not D<sub>2</sub> receptor, mediates nicotine-induced Fos expression in the striatum and nucleus accumbens (Kiba and Jayaraman, 1994; Nisell *et al*, 1997; Svenningsson and Le Moine, 2002). In addition, GTS may exert an inhibitory effect on DA D<sub>1</sub> receptors, which should be investigated by further studies. We found that GTS did not affect resting levels of Fos expression. This is in line with the finding that GTS had no effect on resting levels of extracellular DA.

In this study, for the microdialysis and locomotor activity experiments, nicotine was perfused via the microdialysis probe in a concentration of 10 mM at a rate of 1.5 µl/min (ie, in a dose of 15 nmol/min). Previous microdialysis studies have shown that local perfusion of 1–10 mM (1.5–40 nmol/min) nicotine elicited a reliable and consistent increase in DA content of the striatal dialysate samples (Marshall *et al*, 1997; Lecca *et al*, 2000; Shim *et al*, 2001). Lecca *et al* (2000) reported that perfusion of 0.2 nmol/min nicotine failed to increase DA content, while perfusion of 2 or 10 nmol/min nicotine significantly enhanced DA content in striatal dialysate samples. Also, in our pilot studies, 5 or 10 mM nicotine produced a stable and reliable increase in both DA release and locomotor activity. Thus, we used 10 mM nicotine in our experiments.

A major limitation of this study is that more direct measures of nicotine reinforcement, such as nicotine self-administration and nicotine-induced conditioned place preference, were not employed to assess the effects of GTS. The effects of GTS on those measures of nicotine reinforcement should be examined before drawing definitive conclusions on the potential effects of GTS on nicotine intake. To our knowledge, there is no clinical trial evidence or anecdotal report on the relationship between ginseng use and decreased tobacco consumption. Obviously, clinical studies for the elucidation of the efficacy of ginseng in smoking cessation treatment should be performed in the future.

In conclusion, the results of the present study suggest that GTS acts not only on dopaminergic neurons directly or indirectly to prevent nicotine-induced DA release, but also postsynaptically by binding to DA D<sub>2</sub> receptors. This may explain the blocking effect of GTS on behavioral activation induced by nicotine and conceivably by other psychostimulants. Our data raise the possibility that GTS, by

attenuating nicotine-induced enhancement of dopaminergic transmission, may prove to be a useful therapeutic agent for nicotine addiction and warrant further investigation on its effect on nicotine's rewarding property.

## ACKNOWLEDGEMENTS

This study was supported by grants from the Brain Research Center of the 21st Century Frontier Research Program funded by the Ministry of Science and Technology of Republic of Korea (M103KV010016-04K2201-01630) and the Biomedical Brain Research Center, Ministry of Health and Welfare, Republic of Korea (01-PJ8-PG6-01NE01-0003).

## REFERENCES

- Abe K, Cho SI, Kitagawa I, Nishiyama N, Saito H (1994). Differential effects of ginsenoside Rb1 and malonylginsenoside Rb1 on long-term potentiation in the dentate gyrus of rats. *Brain Res* 649: 7–11.
- Ando T, Tanaka O, Shibata S (1971). Chemical studies on the saponins and sapogenins of ginseng and related crude drugs. *Syoyakugaku Zasshi* 25: 28–32.
- Benwell MEM, Balfour DJK (1992). The effects of acute and repeated nicotine treatment on nucleus accumbens dopamine and locomotor activity. *Br J Pharmacol* 105: 849–856.
- Benwell MEM, Balfour DJK (1994). Comparison of the effects of constant nicotine infusion on nucleus accumbens and striatal dopamine responses to acute nicotine. In: Clarke PBS, Quik M, Thurau K, Adlkofer F (eds). *The Effects of Nicotine on Biological Systems II*. Birkhauser Verlag: Basel, p 66.
- Corrigall WA, Franklin KB, Coen KM, Clarke PB (1992). The mesolimbic dopamine system is implicated in the reinforcing effects of nicotine. *Psychopharmacology* 107: 285–289.
- Damsma G, Day J, Fibiger HC (1989). Lack of tolerance to nicotine-induced dopamine release in the nucleus accumbens. *Eur J Pharmacol* 168: 363–368.
- Di Chiara G, Imperato A (1988). Drugs abused by humans preferentially increase synaptic dopamine concentrations in the mesolimbic system of freely moving rats. *Proc Natl Acad Sci USA* 85: 5274–5278.
- Graybiel AM, Moratalla R, Robertson HA (1990). Amphetamine and cocaine induce drug-specific activation of the *c-fos* gene in striosome-matrix and limbic subdivisions of the striatum. *Proc Natl Acad Sci USA* 87: 6912–6916.
- Henningfield JE, Heishman SJ (1995). The addictive role of nicotine in tobacco use. *Psychopharmacology* 117: 11–13.
- Herrera DG, Robertson HA (1996). Activation of *c-fos* in the brain. *Prog Neurobiol* 50: 83–107.
- Imperato A, Mulas A, Di Chiara G (1986). Nicotine preferentially stimulates dopamine release in the limbic system of freely moving rats. *Eur J Pharmacol* 132: 337–338.
- Kiba H, Jayaraman A (1994). Nicotine induced *c-fos* expression in the striatum is mediated mostly by dopamine D1-receptor and is dependent on NMDA stimulation. *Mol Brain Res* 23: 1–13.
- Kim HS, Hwang SL, Nah SY, Oh S (2001). Changes of [<sup>3</sup>H]MK-801, [<sup>3</sup>H]muscimol and [<sup>3</sup>H]flunitrazepam binding in rat brain by the prolonged ventricular infusion of ginsenoside Rc and Rg1. *Pharmacol Res* 43: 473–479.
- Kim HS, Jang CG, Oh KW, Seong YH, Rheu HM, Cho DH et al (1996a). Effect of ginseng total saponin on cocaine-induced hyperactivity and conditioned place preference in mice. *Pharmacol Biochem Behav* 53: 185–190.
- Kim HS, Jang CG, Park WK, Oh KW, Rheu HM, Cho DH et al (1996b). Blockade by ginseng total saponin of methamphetamine-induced hyperactivity and conditioned place preference in mice. *Gen Pharmacol* 27: 199–204.
- Kim HS, Kang JG, Oh KW (1995a). Inhibition by ginseng total saponin of the development of morphine reverse tolerance and dopamine receptor supersensitivity in mice. *Gen Pharmacol* 26: 1071–1076.
- Kim HS, Kang JG, Seong YH, Nam KY, Oh KW (1995b). Blockade by ginseng total saponin of the development of cocaine induced reverse tolerance and dopamine receptor supersensitivity in mice. *Pharmacol Biochem Behav* 50: 23–27.
- Kim HS, Kim KS (1999). Inhibitory effects of ginseng total saponin on nicotine-induced hyperactivity, reverse tolerance and dopamine receptor supersensitivity. *Behav Brain Res* 103: 55–61.
- Kim S, Ahn K, Oh TH, Nah SY, Rhim H (2002). Inhibitory effect of ginsenosides on NMDA receptor-mediated signals in rat hippocampal neurons. *Biochem Biophys Res Commun* 296: 247–254.
- Kimura T, Saunders PA, Kim HS, Rheu HM, Oh KW, Ho IK (1994). Interactions of ginsenosides with ligand-bindings of GABA<sub>A</sub> and GABA<sub>B</sub> receptors. *Gen Pharmacol* 25: 193–199.
- Ko SR, Kim SC, Choi KJ (1992). Extract yields and saponin contents of red ginseng extracts prepared with various concentrations of ethanol. *Kor J Pharmacogn* 23: 24–28.
- Lecca D, Shim I, Costa E, Javard JI (2000). Striatal application of nicotine, but not of lobeline, attenuates dopamine release in freely moving rats. *Neuropharmacology* 39: 88–98.
- Lidow MS, Goldman-Rakic PS, Rakic P, Innis RB (1989). Dopamine D<sub>2</sub> receptors in the cerebral cortex: distribution and pharmacological characterization with [<sup>3</sup>H]raclopride. *Proc Natl Acad Sci USA* 86: 6412–6416.
- Mansvelder HD, McGehee DS (2002). Cellular and synaptic mechanisms of nicotine addiction. *J Neurobiol* 53: 606–617.
- Marshall DL, Redfern PH, Wonnacott S (1997). Presynaptic nicotine modulation of dopamine release in the three ascending pathway studied by *in vivo* microdialysis: comparison of naive and chronic nicotine-treated rats. *J Neurochem* 68: 1511–1519.
- Matta SG, Foster CA, Sharp BM (1993). Nicotine stimulates the expression of cFos protein in the parvocellular paraventricular nucleus and brainstem catecholaminergic regions. *Endocrinology* 132: 2149–2156.
- Misud JC, Hernandez L, Hoebel BG (1989). Nicotine infused into the nucleus accumbens increases synaptic dopamine as measured by *in vivo* microdialysis. *Brain Res* 478: 365–367.
- Mirza NR, Pei Q, Stolerian IP, Zetterstrom TS (1996). The nicotinic receptor agonists (–)-nicotine and isoarecolone differ in their effects on dopamine release in the nucleus accumbens. *Eur J Pharmacol* 295: 207–210.
- Moratalla R, Elibol B, Vallejo M, Graybiel AM (1996). Network-level changes in expression of inducible Fos-Jun proteins in the striatum during chronic cocaine treatment and withdrawal. *Neuron* 17: 147–156.
- Nisell M, Nomikos GG, Svensson TH (1994a). Systemic nicotine-induced dopamine release in the rat nucleus accumbens is regulated by nicotinic receptors in the ventral tegmental area. *Synapse* 16: 36–44.
- Nisell M, Nomikos GG, Chergui K, Grillner P, Svensson TH (1997). Chronic nicotine enhances basal and nicotine-induced Fos immunoreactivity preferentially in the medial prefrontal cortex of the rat. *Neuropsychopharmacology* 17: 151–161.
- Nisell M, Nomikos GG, Svensson TH (1994b). Infusion of nicotine in the ventral tegmental area or the nucleus accumbens of the rat differentially affects accumbal dopamine release. *Pharmacol Toxicol* 75: 348–352.
- Panagis G, Nisell M, Nomikos GG, Chergui K, Svensson TH (1996). Nicotine injections into the ventral tegmental area increase locomotion and Fos-like immunostaining in the nucleus accumbens of the rat. *Brain Res* 730: 133–142.

- Pang Y, Kiba H, Jayaraman A (1993). Acute nicotine injections induce *c-fos* mostly in non-dopaminergic neurons of the midbrain of the rat. *Mol Brain Res* 20: 162–170.
- Parreno A, Saraza ML, Subero C (1985). A new stabilimeter for small laboratory animals. *Physiol Behav* 34: 475–478.
- Paxinos G, Watson C (1986). *The rat brain in stereotaxic coordinates*, 2nd ed. Academic Press: New York.
- Petkov VW (1978). Effects of ginseng on the brain biogenic monoamines and 3',5'-AMP system: experiments on rats. *Arzneim-Forsch* 28: 388–393.
- Pontieri FE, Francesco E, Tanda G, Francesco O, Di Chiara G (1996). Effects of nicotine on the nucleus accumbens and similarity to those of addictive drugs. *Nature* 382: 255–257.
- Ren T, Sagar SM (1992). Induction of *c-fos* immunostaining in the rat brain after the systemic administration of nicotine. *Brain Res Bull* 29: 589–597.
- Robertson GS, Vincent SR, Fibinger HC (1990). Striatonigral projection neurons contain D1 dopamine receptor-activated *c-fos*. *Brain Res* 523: 288–290.
- Salminen O, Lahtinen S, Ahtee L (1996). Expression of Fos protein in various rat brain areas following acute nicotine and diazepam. *Pharmacol Biochem Behav* 54: 241–248.
- Seong YH, Shin CS, Kim HS, Baba A (1995). Inhibitory effect of ginseng total saponins on glutamate-induced swelling of cultured astrocytes. *Biol Pharm Bull* 18: 1776–1778.
- Shim I, Javaid JI, Kim SE (2000). Effect of ginseng total saponin on extracellular dopamine release elicited by local infusion of nicotine into the striatum of freely moving rats. *Planta Med* 66: 705–708.
- Shim I, Javaid JI, Wirtshafter D, Jang SY, Shin KH, Lee HJ et al (2001). Nicotine-induced behavioral sensitization is associated with extracellular dopamine release and expression of c-Fos in the striatum and nucleus accumbens of the rat. *Behav Brain Res* 121: 137–147.
- Smolders I, De Klippel N, Sarre S, Ebinger G, Michotte Y (1995). Tonic GABA-ergic modulation of striatal dopamine release studied by *in vivo* microdialysis in the freely moving rat. *Eur J Pharmacol* 284: 83–91.
- Svenningsson P, Le Moine C (2002). Dopamine D1/5 receptor stimulation induces *c-fos* expression in the subthalamic nucleus: possible involvement of local D5 receptors. *Eur J Neurosci* 15: 133–142.
- Tachikawa E, Kudo K, Kashimoto T, Takahashi E (1995). Ginseng saponins reduce acetylcholine-evoked Na<sup>+</sup> influx and catecholamine secretion in bovine adrenal chromaffin cells. *J Pharmacol Exp Ther* 273: 629–636.
- Tokuyama S, Oh KW, Kim HS, Takahashi M, Kaneto H (1992). Blockade by ginseng extract of the development of reverse tolerance to the ambulation-accelerating effect of methamphetamine in mice. *Jpn J Pharmacol* 59: 423–425.
- Toth E, Sershen H, Hashim A, Vizi ES, Lajtha A (1992). Effect of nicotine on extracellular levels of neurotransmitters assessed by microdialysis in various brain regions: role of glutamic acid. *Neurochem Res* 17: 265–271.
- Tsang D, Yeung HW, Tso WW, Peck H (1985). Ginseng saponins: influence on neurotransmitter uptake in rat brain synaptosomes. *Planta Med* 3: 221–224.
- Valentine JD, Matta SG, Sharp BM (1996). Nicotine-induced cFos expression in the hypothalamic paraventricular nucleus is dependent on brainstem effects: correlations with cFos in catecholaminergic and noncatecholaminergic neurons in the nucleus tractus solitarius. *Endocrinology* 137: 622–630.
- Wonnacott S, Drasdo A, Sanderson E, Rowell P (1990). Presynaptic nicotinic receptors and modulation of transmitter release. In: Bock G, Marsh J (eds). *The Biology of Nicotine Dependence*. John Wiley & Sons: Chichester. pp 87–101.