

Parametric and Regional Maps of Free Serotonin 5HT_{1A} Receptor Sites in Human Brain as Function of Age in Healthy Humans

Mette Møller^{*1,2}, Steen Jakobsen² and Albert Gjedde^{1,2}

¹Center of Functionally Integrative Neuroscience, Aarhus University, Aarhus, Denmark; ²PET-Centre, Aarhus University Hospital, Aarhus, Denmark

Serotonin 5HT_{1A}-binding sites can be detected in living human brain with the positron-emitting antagonist [¹¹C]WAY-100635. Previous measurements of the availability of [¹¹C]WAY-binding sites in normal aging are equivocal, in part because of the greatly variable binding of this ligand. To test the null hypothesis that the binding potential (pB) of 5HT_{1A} sites remains constant with age; 19 healthy volunteers aged 23–73 years (8 women, 11 men) underwent positron emission tomography. To determine pBs, we applied a novel tissue reference method of analysis, Estimation of Reversible Ligand Binding and Receptor Density (ERLiBiRD) (Gjedde, 2003; Rosa-Neto *et al*, 2004), which extrapolates measures of specific binding to an estimated steady-state. We compared these estimates in the two age groups with results obtained with the conventional Logan Plot and Simplified Reference Tissue Method (SRTM) applied to both regions of interest-based as parametric analyses. The regional distribution of specific binding of free sites [¹¹C]WAY-100635 was similar to that reported in previous studies, with the highest pBs in limbic structures and the raphe nuclei. Although the results of the three methods differed, pBs in the elderly subjects consistently were lower than those of young subjects. Thus, the correlation between pB and age applied to regions-of-interest revealed significant decline of pB at the rate of 3 or 4% per decade, and a 10% decline of the global mean 5HT_{1A} receptor-pB in elderly relative to young subjects. The results demonstrate that the number of available 5HT_{1A}-binding sites declines with age. *Neuropsychopharmacology* (2007) **32**, 1707–1714; doi:10.1038/sj.npp.1301310; published online 24 January 2007

Keywords: PET; serotonin 5HT_{1A} receptors; [¹¹C]WAY-100635; brain; aging

INTRODUCTION

Serotonin 5HT_{1A} receptors play a dual role in the modulation of serotonin neurotransmission, acting as presynaptic autoreceptors on serotonergic neurons, and as postsynaptic receptors throughout the cerebral cortex. *In vitro* autoradiography reveals presynaptic serotonin 5HT_{1A} receptors on the soma of serotonergic neurons in the raphe nuclei, where they regulate the firing of these neurons (Arango *et al*, 2001; Hall *et al*, 1997). Most binding sites are located postsynaptically on the axon hillock or soma of cortical pyramidal neurons (Amargos-Bosch *et al*, 2004; Azmitia *et al*, 1996). The density of cortical binding is heterogeneous with the highest binding in limbic structures such as hippocampus and cingulate gyrus, and the lowest binding in occipital lobe, thalamus, and basal ganglia.

Free serotonin 5HT_{1A}-binding sites have been mapped in PET studies of the living human brain with the positron-

emitting antagonist [¹¹C]WAY-100635 ([¹¹C]WAY) (Pike *et al*, 1995; Wilson *et al*, 1998). Subtle abnormalities in the availability of [¹¹C]WAY-binding sites were reported in clinical PET studies of personality differences (Borg *et al*, 2003), depression (Drevets *et al*, 1999; Bhagwagar *et al*, 2004; Sargent *et al*, 2000), schizophrenia (Tauscher *et al*, 2002), juvenile myoclon epilepsy (JME) (Meschaks *et al*, 2005), bulimia nervosa (Tiihonen *et al*, 2004), ALS (Turner *et al*, 2005), and Parkinson's disease (PD) (Doder *et al*, 2003). Together, these studies show that [¹¹C]WAY binding, as an index of serotonin 5HT_{1A} receptor availability, is correlated with scores of personality, increased in specific cortical regions of patients with schizophrenia and bulimia nervosa, globally decreased in patients with depression and ALS, and focally decreased in patients with JME and in the raphe nucleus of PD patients. Thus, PET images of the [¹¹C]WAY-binding potential (pB) in brain demonstrate disease-specific patterns of disturbed serotonergic neurotransmission.

However, the results of these studies must be weighed against the possible, but not yet proven effect of normal aging on receptor density. The evidence for an effect of normal aging on the availability of [¹¹C]WAY-binding sites is equivocal. Thus, normal aging showed significant decline

*Correspondence: Dr M. Møller, PET-Centre, Aarhus University Hospital, 8000 Aarhus C, Denmark, Tel: +45 8949 4408, Fax: +45 8949 4400, E-mail: moller@pet.auh.dk
Received 21 July 2006; revised 10 November 2006; accepted 15 November 2006

of cortical [¹¹C]WAY binding in one study of rhesus monkeys (Tsukada *et al*, 2001) and three studies of humans (Tauscher *et al*, 2001; Cidis Meltzer *et al*, 2001; Bhagwagar *et al*, 2004), where as no correlation with age was reported in two other studies of humans (Parsey *et al*, 2002; Rabiner *et al*, 2002). In addition, the findings may reflect problems with kinetic analysis of binding due to the very rapid metabolism of the tracer in the circulation (Gunn *et al*, 1998; Osman *et al*, 1998) and the reported high inter-subject variability in [¹¹C]WAY binding, as demonstrated in a large cohort of healthy male subjects. The study found no correlation between *pB* and demographics, physiological or behavioral variables or levels of endogenous serotonin (Rabiner *et al*, 2002).

The present study tested the null hypothesis of no effect of normal aging on the results of non-invasive assays of [¹¹C]WAY binding in brain of healthy young and elderly normal subjects. To this end, we applied standard tissue reference methods to regions of interest (ROIs) and image voxels to obtain parametric estimates in the form of *pB* of [¹¹C]WAY. In addition, we compared the results to the results from the tissue reference method ERLiBiRD (Gjedde, 2003; Rosa-Neto *et al*, 2004). This method obtains estimates of extrapolation of specific uptakes ratios towards a predicted steady-state.

SUBJECTS AND METHODS

Recruitment

This study was approved by the Research Ethics Committee of Aarhus County. Nineteen healthy volunteers aged 23–73 years, divided in two groups with an average age in the elderly group of 12 subjects of 62.5 ± 6.8 years (five women and seven men), and an average age in the young group of seven subjects of 25.1 ± 2.0 years (three women, four men), gave written informed consent to the study. Exclusion criteria included cardiovascular disease and any history of neurological and psychiatric disease. All subjects were physically fit, free of prescribed medication, and did not meet the criteria for depression according to DSM IV.

Radiochemistry

[¹¹C-*carbonyl*]WAY was synthesized from the cyclotron-generated precursor according to the method of McCarron *et al* (1996). In brief, ¹¹CO₂ was collected in a stainless steel cryotrap, warmed, and then flushed with nitrogen through a small coil of 1/16" o.d. polypropylene tubing, which had been earlier flushed with a solution of cyclohexylmagnesium chloride (500 µl of 0.5 M in THF). The [¹¹C]-labeled Grignard adduct, [¹¹C]cyclohexanecarbonyl chloride, was eluted with a solution of thionyl chloride (10 µl in 400 µl THF) into a septum-sealed vial (2 ml) containing WAY-100634 (2 mg), 50 µl THF and triethylamine (60 µl) under a nitrogen atmosphere. The reaction proceeded in seven min at 85°C. The crude product was diluted with 500 µl of HPLC eluent, and purified by reversed phase HPLC on a Ultracarb 7 ODS 30 column (250 × 10 mm; Phenomenex Ltd.), eluting with sterile ethanol/70 mM NaH₂PO₄ (52:48) at 6 ml/min. The fraction containing [¹¹C-*carbonyl*]WAY-100635 (retention time ca. 10 min) was collected, evaporated to near-

dryness at 90°C under vacuum, and then reformulated in sterile saline (10 ml) before passage through a sterile 0.22 µm filter into a sterile vial. Mean specific radioactivity was 42 GBq/µmol (SD 32).

PET and Data Analysis

Subjects were positioned in the ECAT EXACT HR47 tomograph (CTI/Siemens, Knoxville, TN, USA). After a 15 min attenuation scan, 60 min emission recordings in 3-D mode consisting of 22 frames were initiated after intravenous bolus injection of [¹¹C]WAY (150–430 MBq, mean 270, SD 111). High resolution T1-weighted MR scans were made with a 1.5 or 3 T magnet (GE Sigma Systems, Milwaukee, USA). The summed emission recordings were automatically co-registered to the individual MRI scans using published methods (Talairach and Tournoux, 1988; Collins *et al*, 1994; Iversen *et al*, 2006). In brief, individual MR-images were co-registered to a common stereotaxic space (Montreal Neurological Institute) using a 12 parameter affine rigid body transformation; registrations were automatic in the young group, but manual in the elderly group because of inadequate registration of potentially atrophied brains. After calculation of the final PET-Talairach transformation matrix, dynamic emission recordings were resampled into the common coordinates and, using the program DISPLAY, templates of ROIs were drawn bilaterally on the average MR image: hippocampus, and the insular, cingulate and ventral medial prefrontal cortices. Owing to central and cortical atrophy in the elderly, each of the eight regions was applied to the individual resampled MR scan and adjusted to fit. The raphe nuclei were not visible on MR images and were outlined on the individual PET images in each subject. A ROI for the cerebellum (15 cm³) was drawn manually on the MR anatomical template, excluding the vermis and carefully avoiding transverse sinuses and the inferior occipital lobe. Finally, time-radioactivity curves were extracted from the dynamic PET image based on the ROI templates of all 10 regions. We assumed that *pBs* depend on both age and ROI. The linear relation between age, ROI and *pB* were assessed using linear regression for each of the binding potential estimates obtained with conventional methods Logan and SRTM and the method of ERLiBiRD.

Kinetic Analysis

5HT_{1A} binding was quantified using ROI-based and voxel-wise parametric analysis (Table 1). The estimated quantity is the *pB*, which is a measure of the binding of serotonin to a class of receptors, relative to its concentration, and relative to the ratio between the affinity of serotonin and the radioligand. The class of receptors to which the measure applies is determined by the selectivity of the radioligand. In this case, the selectivity of the radioligand is strongly in favor of the serotonin 5HT_{1A} receptors. The $pB = (B_{\max} - B)/K_d$ is a measure of the number of unoccupied receptor sites relative to the affinity towards a ligand.

Parametric images of the *pB* were obtained in three different ways, including the tissue reference method of Logan (Logan, 2000), excluding the first 4 min of data, the multi-linear version of simplified tissue reference region

Table 1 Methods Applied to Imaging Data to Obtain Parametric and Regions-of-Interest (ROI) Maps of Binding Potentials

Methods of analysis	Linear regression	Nonlinear regression
Region-based (ROI)	Reference Logan Plot (Logan, 2000)	ERLiBiRD (Gjedde, 2003) & SRTM (Lammertsma and Hume, 1996)
Voxel-based	Logan & LSRTM (SRTM version of Zhou <i>et al</i> , 2003)	ERLiBiRD (Gjedde, 2003)

method (Zhou *et al*, 2003) and ERLiBiRD (Rosa-Neto *et al*, 2004), excluding the first minute of data.

Using cerebellum as reference region, average uptakes in ROIs were also analyzed by the Logan *et al*, SRTM (Gunn *et al*, 1998; Lammertsma and Hume, 1996) and ERLiBiRD methods (Rosa-Neto *et al*, 2004). The ERLiBiRD method fits the following Equation

$$\rho^* = 1 + p_B(1 - e^{-\alpha\Theta}) + (R_1 - 1)e^{-\alpha\Theta}$$

to paired calculations of the ratio of areas-under-the-curves (AUC ratio = ρ^*) of voxels or regions of binding and reference, and the modified time variable Θ'' equal to the ratio

$$\Theta = \left[\frac{\int_0^T \int_0^u m_{\text{ref}}^* dt du}{\int_0^T m_{\text{ref}}^* dt} \right]$$

where m_{ref}^* is the time activity function of the reference region.

RESULTS

In both elderly and young subjects, the cerebellum time-activity curve, normalized to its peak value, had the same general form consistent with flow-limited clearance from the brain tissue (Figure 1). There was no difference between the patterns of washout from the cerebellum in the two groups of subjects. Figure 2 shows the average parametric image of [¹¹C]WAY binding in groups of young and elderly subjects, calculated by the methods of Logan *et al*, Zhou *et al* (multi-linear SRTM) and ERLiBiRD. The highest p_B s were determined in limbic cortical structures, frontal and temporal neocortex, and in the raphé nuclei, irrespective of the method used for calculating p_B . p_B images obtained with ERLiBiRD globally were 25% higher than by the two other methods.

The mean magnitudes of p_B by the ROI analysis in nine brain regions in the two groups are listed in Table 2. Also ROI-based analyses globally gave lower p_B values with the Logan method than with the ERLiBiRD and SRTM methods. The correlations among the mean p_B estimates of the several methods were consistently high (SRTM vs ERLiBiRD: $r^2 = 0.90$, SRTM vs Logan: $r^2 = 0.73$) (Figure 3). Irrespective of method, the highest p_B s were seen in insula and hippocampus. Values of [¹¹C]WAY binding consistently were either high or low across the regions of all subjects, including the raphé nuclei.

A significant decrease in p_B with age was demonstrated in the ROIs with postsynaptic-binding sites with a reduction in p_B per decade of 0.14 in SRTM ($p = 0.02$), 0.17 in ERLiBiRD ($p < 0.001$) and 0.13 in Logan ($p \leq 0.001$), when adjusting for the significant effect of region (data not shown), thus




Figure 1 Time-activity curves recorded in cerebellum in the 19 healthy subjects of this study (red = elderly group, blue = young group). Abscissa: time (min). Ordinate: activity normalized to peak activity.

averaging a significant decrease of p_B of 3–4% per decade. Figure 4 shows the correlation between p_B and age in five regions-of-interest for the three methods of calculation. A decline in p_B with age was demonstrated in all five regions measured by the three methods except in raphe calculated by the Logan method. A significant reduction was seen in hippocampus and insula using the methods of ERLiBiRD and Logan, the reduction in MPFC was only significant in Logan method. No method yielded a significant decrease in cingulate or the raphe region. SRTM showed a decline with age in all regions, but did not reach significance probably due to its lower precision compared to the Logan method. The ROI-based analysis also revealed a 10% decrease of the global mean magnitude of 5HT_{1A} receptor magnitude of p_B in elderly subjects relative to young healthy subjects. T -test for equal p_B in young and old were significant in all three methods ($p = 0.02$ in SRTM, $p = 0.003$ in ERLiBiRD, $p = 0.004$ in Logan) (Table 2).

DISCUSSION

5HT_{1A} receptors reside in all the cortical layers, but most densely so in the superficial layers (Burnet *et al*, 1995). Agonists of 5HT_{1A} receptors evoke hyperpolarization of pyramidal neurons, unlike agonists of 5HT_{2A} receptors, which cause depolarization (Amargos-Bosch *et al*, 2004; Aghajanian and Marek, 1997; Cruz *et al*, 2004). Therefore, 5HT_{1A} receptors are believed to fundamentally modulate cortical excitation.




Figure 2 Parametric maps of pB of radioligand [^{11}C]WAY in selected sections of brain tissue determined by three different methods, multilinear simplified reference tissue regression method LSRTM (Zhou *et al*, 2003), non-linear regression method ERLiBiRD (Gjedde, 2003), and tissue reference version of Logan Plot method (Logan, 2000). Top three rows show sagittal views, second-to-last row transaxial view, and bottom row coronal view. Left column shows young adults, right column aged adults. Color bars shown to right of columns show color scale for LSRTM and Logan methods above and color scale for ERLiBiRD method below, respectively.

PET studies of several classes of neuroreceptors, for example; dopamine D₁ (Wang *et al*, 1998) and D₂ (Farde *et al*, 1995; Antonini and Leenders, 1993), histamine H₁ (Yanai *et al*, 1992) and serotonin 5HT_{2A} (Adams *et al*, 2004; Meltzer *et al*, 1998; Sheline *et al*, 2002) demonstrate age-dependent decline of the availability of receptors of the order of 10% per decade. Transporters of dopamine (van Dyck *et al*, 2002) and serotonin (Yamamoto *et al*, 2002) also tend to decline with age. In contrast, ambiguous findings have made it difficult to confirm a similar decline of serotonin 5HT_{1A} receptors. Thus, the results reported in the literature are inconclusive based on small sample sizes and biased by the lack of elderly participants, high intersubject variability in the range of 16–24% (Rabiner *et al*, 2002), and the fact that in no study did the researchers follow the same subjects with age. One study demonstrated an association with age in young and aged monkeys with decrease of binding to receptors in temporal and frontal cortices in the elderly monkeys, but less displacement of [^{11}C]WAY binding after injection of the agonist 8-OH-DPAT, suggest-

ing impaired affinity in the aged monkeys (Tsukada *et al*, 2001). One study of humans subjects aged 22–53 years found a 10% decline per decade of pB in the whole brain, except in mesiotemporal cortex (Tauscher *et al*, 2001), and one other study of a wider age range of human subjects (21–80 years) found significant decrease in receptor-rich areas of the brain, but only in men. A more recent study on previously depressed subjects reported a decrease with age in 5HT_{1A} receptor binding compared to controls in both raphé nuclei and postsynaptic regions (Bhagwagar *et al*, 2004). The decrease was more pronounced in the group of depressed subjects, but their ages were not specified. Two other studies found no association between age and pB : They included a large database of 61 males with a narrow age-span (24–53 years), and a smaller study of both genders (20–70 years) with higher pB in women than in men regardless of age (Parsey *et al*, 2002; Rabiner *et al*, 2002). Selected studies of WAY-100635 binding are listed in Table 3. In general, the present change is more modest than reported in the two previous studies showing a significant correlation with age, and reported in studies of the effects of diverse pathologies. The majority of studies listed in Table 3 used the ‘gold standard’ SRTM method but a few used the Logan plot. The magnitude of pB values determined in the present study are consistent with those reported in the literature. The definition of the pB as an index of unoccupied receptors makes a direct comparison with total receptor numbers somewhat tenuous, as reasons for decline may include changes of bound serotonin as well as changes of receptor affinity. Therefore, it is possible that the variability of past results is related to fluctuations of bound serotonin or receptor affinity in addition to receptor numbers.

The agonist 8-OH-[^3H]DPAT is most often used in postmortem studies of 5HT_{1A} receptors. The radioligand binds to high-affinity receptors coupled to G-proteins only and hence reveals only half of the binding of the antagonist [^3H]WAY-100635. Direct comparison of receptor density *in vitro* and *in vivo* studies is not feasible with [^{11}C]WAY (Burnet *et al*, 1997) of the receptor density, although the distribution of the binding of [^{11}C]WAY *in vivo* studies generally fits with the postmortem findings (Hall *et al*, 1997).

As cerebellum is almost devoid of 5HT_{1A} receptors (Hall *et al*, 1997), uptake of the tracer in this region is often used as a surrogate for an arterial input function in the absence of blood sampling. In some studies with [^{11}C]WAY (Rabiner *et al*, 2002; Doder *et al*, 2003; Turner *et al*, 2005), as many as 10% of the subjects were excluded because of abnormally high cerebellar time-activity curves potentially caused by specific binding in cerebellum, incorrect segmentation of cerebellum, or variable properties of the tracer. One of the known properties of [^{11}C]WAY is the very rapid metabolism in the circulation (Gunn *et al*, 1998; Osman *et al*, 1998), which may violate the underlying assumption of continuous exchange of brain tissue with circulation and thereby may induce biases to the results of previously applied kinetic models.

The issue of use of the cerebellum as reference biased by binding of the tracer to receptors in different subdivisions of the cerebellum has been addressed in the literature in various ways, most of them based on the use of arterial

Table 2 Mean Magnitude of Binding Potentials (pB) in Nine Brain Regions in Elderly and Young Healthy Volunteers, Respectively

Method	1	2	3	4	5	6	7	8	9	Mean pB
<i>Mean pB values in 9 regions in 7 young subjects</i>										
SRTM	4.73	4.45	6.50	6.78	5.09	5.41	6.43	6.25	3.52	5.46 ^a /5.70 ^b
	±0.66	0.66	1.19	1.35	0.63	1.55	1.07	1.20	1.01	1.11/0.90
ERLiBiRD	4.46	4.07	5.76	5.53	4.58	4.65	5.90	5.65	3.06	4.85/5.08
SD	0.71	0.62	1.08	0.99	0.87	0.86	1.07	1.05	0.63	0.94/0.71
LOGAN	2.82	2.82	3.64	3.47	3.34	3.37	3.48	3.37	1.73	3.12/3.29
SD	0.50	0.52	0.78	0.74	0.77	0.51	0.60	0.57	0.37	0.59/0.30
<i>Mean pB values in 9 regions in 12 elderly subjects</i>										
SRTM	4.37	4.77	5.77	5.24	4.84	4.82	5.69	5.75	3.29	4.95/5.16
SD	0.98	1.41	2.63	1.21	1.13	1.58	1.49	1.23	0.64	0.8/0.54
ERLiBiRD	4.01	4.07	4.43	4.39	4.06	4.21	4.68	4.74	2.89	4.16/4.32
SD	1.01	1.13	1.36	1.21	0.92	1.53	1.21	0.97	0.80	0.55/0.28
LOGAN	2.58	2.72	3.02	3.06	2.99	2.76	2.94	3.08	1.87	2.78/2.89
SD	0.37	0.55	0.72	0.67	0.50	0.63	0.46	0.37	0.22	0.38/0.18

1. Left cingulate. 2. Right cingulate. 3. Left hippocampus (postsynaptic). 4. Right hippocampus. 5. Left medial prefrontal cortex. 6. Right medial prefrontal cortex. 7. Left insula. 8. Right insula. 9. Raphe nuclei. ^aAll 9 regions. ^bRegion 1–8.




Figure 3 Correlation among methods used to obtain pB values in anatomically defined ROIs in 19 healthy subjects. Abscissae: values of pB obtained with method SRTM. Ordinates: Values of pB obtained with methods ERLiBiRD and Logan, left and right graph, respectively. Correlation coefficients are $r^2 = 0.90$ and 0.73, left and right graph, respectively.

samples to determine binding in cerebellum on the basis of an input function. This approach is seriously limited because of the uncertainty of tracer and tracer metabolite concentrations in arterial blood. An alternative has been to examine pBs in other parts of the brain with the use of cerebellar reference tissues segmented into gray and white matter. In the present study, the cerebellar reference was limited to white matter, as described recently (Hirvonen *et al*, 2007).

In the study, both parametric maps and ROI-based analysis were performed to estimate the binding of [¹¹C]WAY to 5HT_{1A} receptors using linear and non-linear regression methods. As reported in previous studies, pB values calculated by non-linear methods were higher, but less precise than the values calculated by linear models. The ROI-based results showed an agreement between the gold standard SRTM, and the model-independent method ERLiBiRD with values ranging from 4 to 6. In the parametric

maps the multi-linear SRTM (Zhou *et al*, 2003 version) gave low values of 2–3.5 in line with the Logan method results, whereas ERLiBiRD findings were about 25% higher.

The point of the study was to avoid the use of arterial sampling because of the uncertainty associated with the rapid tracer metabolism in arterial blood. Of the three methods, the SRTM conventionally is considered the gold standard because dependent and independent variables are fully separated. However, the method is sensitive to low cerebellar concentrations towards the end of the circulation time, and hence was supplemented with integral methods, which tend to lower statistical noise. Of these, the Logan method still uses the actual concentration of the tracer in ROI in the denominator of both axes and hence retains a residual sensitivity to noise in regions of low specific binding. Only the ERLiBiRD method is based exclusively on the use of integrals. Thus, the Logan method has a known bias in the direction of low pBs, which we confirm in the




Figure 4 Correlation among measures of pB and age in five regions-of-interest: (a) Hippocampus, (b) Cingulate, (c) Medial prefrontal cortex, (d) Insula, and (e) Raphe nuclei. The slopes found in the three methods SRTM (red), ERLiBiRD (green), and Logan (blue) are listed in the panel including p -values. Abscissae: Age (years). Ordinates: pB values (ratio).

present study. For these reasons, we find that the ERLiBiRD has the best balance of precision and accuracy.

Limitations

A potential bias induced by differences in gender specific effects of age cannot be excluded and may magnify or dilute the effects of age in a group of both genders. Owing to the small sample size of this study, it was not possible to

demonstrate a difference in pB between women and men. In the literature, only two studies (Parsey *et al*, 2002; Cidis Meltzer *et al*, 2001) addressed the issue of gender. In the first study, women had higher pB s than men, but also higher volumes of distribution in cerebellum. In the second study, women had a non-significant trend towards higher pB than men, whereas an age-related decline in pB was seen only in men. Thus, the literature is limited and inconclusive on the subject of the influences of age and gender on the pB of WAY.

Table 3 Human Studies of 5HT_{1A} Receptors with WAY-100635

Author/journal	Sample size	Change in pB
Bhagwagar <i>et al</i> (2004) <i>Mol Psychiatry</i>	18 depressed patients/14 controls (males)	17% decrease in patients compared to controls
Cidis Meltzer <i>et al</i> (2001) <i>Brain Res</i>	21 controls (10 males, 11 females)	Decrease in pB with age (men only), % not listed
Doder <i>et al</i> (2003) <i>Neurology</i>	23 PD patients (16 males, 7 females)/8 controls (6 males, 2 females)	27% decrease in raphe of patients compared to controls
Parsey <i>et al</i> (2002) <i>Brain Res</i>	25 controls (12 females, 13 males)	No change in pB with age
Drevets <i>et al</i> (1999) <i>Biol Psychiatry</i>	12 depressed patients (7 females, 5 males)/8 control (4 females, 4 males)	42% decrease in raphe and 27% decrease in mesial temporal cortex (MTC) in patients compared to controls
Rabiner <i>et al</i> (2002) <i>Neuroimage</i>	61 male controls	No change in pB with age
Sargent <i>et al</i> (2000) <i>Arch Gen Psychiatry</i>	25 depressed patients/(22 males, 3 females) 18 controls (17 males, 1 female)	11–12% decrease in patients compared to controls
Tauscher <i>et al</i> (2002) <i>Arch Gen Psychiatry</i>	14 schizophrenic patients (6 females, 8 males)/14 controls (8 females, 6 males)	20% increase in pB in left MTC, 13 % in pB in right MTC in patients compared to controls
Tauscher <i>et al</i> (2001) <i>Neuropsychopharmacology</i>	19 controls (8 females, 11 males)	10% decrease per decade
Turner <i>et al</i> (2005) <i>Brain</i>	21 ALS patients (4 females, 17 males)/19 controls (5 females, 14 males)	21% decrease in patients compared to controls
Meschaks <i>et al</i> (2005) <i>Archives of Neurology</i>	11 patients with juvenile myoclonic epilepsy (6 females, 5 males)/11 controls (6 females, 5 males)	20–28% decrease in hippocampus, dorsolateral prefrontal cortex, and raphe in patients compared to controls
Tiihonen <i>et al</i> (2004) <i>Biological Psychiatry</i>	8 female bulimic patients/10 female controls	Increase in pB in patients compared to controls, % not listed

Varying degrees of atrophy were observed in the elderly brains, but we did not specifically correct for atrophy. This means that an overestimation of the decline in pB with age is possible. One study observed the same age-related decline with and without correction for atrophy (Cidis Meltzer *et al*, 2001). On the other hand, it is known that the currently available methods for correction may induce other biases due to overestimation of cortical thickness and masking of a true decline of the total number of receptors because the correction addresses concentrations of receptors in the tissue rather than their total numbers.

The findings confirm a significant correlation of the pB with age regardless of the chosen model, but the issue of whether the decrease reflects a reduced receptor affinity, decline in density because of loss of receptors due to neuronal loss or atrophy or caused by methodological biases awaits resolution in future studies.

ACKNOWLEDGEMENTS

We thank OL Munk, PhD, for his kind assistance with the regression analysis and the use of the regression program PIP. The authors have no involvement, financial or otherwise, that might potentially bias the presented work. The work was supported by center-of-excellence grants from the National Science Foundation of Denmark to the Center of Functionally Integrative Neuroscience, Aarhus University, and by a fellowship from Aarhus University to MM.

REFERENCES

Adams KH, Pinborg LH, Svarer C, Hasselbalch SG, Holm S, Haugbol S *et al* (2004). A database of [(18)F]-altanserin binding

to 5-HT(2A) receptors in normal volunteers: normative data and relationship to physiological and demographic variables. *Neuroimage* **21**: 1105–1113.

Aghajanian GK, Marek GJ (1997). Serotonin induces excitatory postsynaptic potentials in apical dendrites of neocortical pyramidal cells. *Neuropharmacology* **36**: 589–599.

Amargos-Bosch M, Bortolozzi A, Puig MV, Serrats J, Adell A, Celada P *et al* (2004). Co-expression and *in vivo* interaction of serotonin1A and serotonin2A receptors in pyramidal neurons of prefrontal cortex. *Cereb Cortex* **14**: 281–299.

Antonini A, Leenders KL (1993). Dopamine D2 receptors in normal human brain: effect of age measured by positron emission tomography (PET) and [¹¹C]-raclopride. *Ann N Y Acad Sci* **695**: 81–85.

Arango V, Underwood MD, Boldrini M, Tamir H, Kassir SA, Hsiung S *et al* (2001). Serotonin 1A receptors, serotonin transporter binding and serotonin transporter mRNA expression in the brainstem of depressed suicide victims. *Neuropsychopharmacology* **25**: 892–903.

Azmitia EC, Gannon PJ, Kheck NM, Whitaker-Azmitia PM (1996). Cellular localization of the 5-HT1A receptor in primate brain neurons and glial cells. *Neuropsychopharmacology* **14**: 35–46.

Bhagwagar Z, Rabiner EA, Sargent PA, Grasby PM, Cowen PJ (2004). Persistent reduction in brain serotonin1A receptor binding in recovered depressed men measured by positron emission tomography with [¹¹C]WAY-100635. *Mol Psychiatry* **9**: 386–392.

Borg J, Andree B, Soderstrom H, Farde L (2003). The serotonin system and spiritual experiences. *Am J Psychiatry* **160**: 1965–1969.

Burnet PW, Eastwood SL, Harrison PJ (1997). [³H]WAY-100635 for 5-HT1A receptor autoradiography in human brain: a comparison with [³H]8-OH-DPAT and demonstration of increased binding in the frontal cortex in schizophrenia. *Neurochem Int* **30**: 565–574.

Burnet PW, Eastwood SL, Lacey K, Harrison PJ (1995). The distribution of 5-HT1A and 5-HT2A receptor mRNA in human brain. *Brain Res* **676**: 157–168.

- Cidris Meltzer C, Drevets WC, Price JC, Mathis CA, Lopresti B, Greer PJ *et al* (2001). Gender-specific aging effects on the serotonin 1A receptor. *Brain Res* **895**: 9–17.
- Collins DL, Neelin P, Peters TM, Evans AC (1994). Automatic 3D intersubject registration of MR volumetric data in standardized Talairach space. *J Comput Assist Tomogr* **18**: 192–205.
- Cruz DA, Eggan SM, Azmitia EC, Lewis DA (2004). Serotonin1A receptors at the axon initial segment of prefrontal pyramidal neurons in schizophrenia. *Am J Psychiatry* **161**: 739–742.
- Doder M, Rabiner EA, Turjanski N, Lees AJ, Brooks DJ (2003). Tremor in Parkinson's disease and serotonergic dysfunction: an ¹¹C-WAY 100635 PET study. *Neurology* **60**: 601–605.
- Drevets WC, Frank E, Price JC, Kupfer DJ, Holt D, Greer PJ *et al* (1999). PET imaging of serotonin 1A receptor binding in depression. *Biol Psychiatry* **46**: 1375–1387.
- Farde L, Hall H, Pauli S, Halldin C (1995). Variability in D2-dopamine receptor density and affinity: a PET study with [¹¹C]raclopride in man. *Synapse* **20**: 200–208.
- Gjedde A (2003). Modelling metabolite and tracer kinetics. In: Feinendezen LE, Shreeve WW, Eckelman WC, Bahk Y-W, Wagner Jr HW (eds) *Molecular Nuclear Medicine*. Springer-Verlag: Berlin, Heidelberg, pp 121–169.
- Gunn RN, Sargent PA, Bench CJ, Rabiner EA, Osman S, Pike VW *et al* (1998). Tracer kinetic modeling of the 5-HT_{1A} receptor ligand [carbonyl-¹¹C]WAY-100635 for PET. *Neuroimage* **8**: 426–440.
- Hall H, Lundkvist C, Halldin C, Farde L, Pike VW, McCarron JA *et al* (1997). Autoradiographic localization of 5-HT_{1A} receptors in the post-mortem human brain using [³H]WAY-100635 and [¹¹C]way-100635. *Brain Res* **745**: 96–108.
- Hirvonen J, Kajander J, Allonen T, Oikonen V, Nagren K, Hietala J (2007). Measurement of serotonin 5-HT_{1A} receptor binding using positron emission tomography and [carbonyl-(11)C]WAY-100635—considerations on the validity of cerebellum as a reference region. *J Cereb Blood Flow Metab* **27**: 185–195.
- Iversen P, Hansen DA, Bender D, Rodell A, Munk OL, Cumming P *et al* (2006). Peripheral benzodiazepine receptors in the brain of cirrhosis patients with manifest hepatic encephalopathy. *Eur J Nucl Med Mol Imag* **33**: 810–816.
- Lammertsma AA, Hume SP (1996). Simplified reference tissue model for PET receptor studies. *Neuroimage* **4**: 153–158.
- Logan J (2000). Graphical analysis of PET data applied to reversible and irreversible tracers. *Nucl Med Biol* **27**: 661–670.
- McCarron J, Turton J, Pike V, Poole K (1996). Remotely controlled production of the 5-HT_{1A} receptor radioligand, [carbonyl-¹¹C]-WAY-100635, via ¹¹C-carboxylation of an immobilized Grignard reagent. *J Labelled Compd Radiopharmacol* **38**: 941–953.
- Meltzer CC, Smith G, Price JC, Reynolds III CF, Mathis CA, Greer P *et al* (1998). Reduced binding of [18F]altanserin to serotonin type 2A receptors in aging: persistence of effect after partial volume correction. *Brain Res* **813**: 167–171.
- Meschaks A, Lindstrom P, Halldin C, Farde L, Savic I (2005). Regional reductions in serotonin 1A receptor binding in juvenile myoclonic epilepsy. *Arch Neurol* **62**: 946–950.
- Osman S, Lundkvist C, Pike VW, Halldin C, McCarron JA, Swahn CG *et al* (1998). Characterisation of the appearance of radioactive metabolites in monkey and human plasma from the 5-HT_{1A} receptor radioligand, [carbonyl-¹¹C]WAY-100635—explanation of high signal contrast in PET and an aid to biomathematical modelling. *Nucl Med Biol* **25**: 215–223.
- Parsey RV, Oquendo MA, Simpson NR, Ogden RT, Van Heertum R, Arango V *et al* (2002). Effects of sex, age, and aggressive traits in man on brain serotonin 5-HT_{1A} receptor binding potential measured by PET using [C-11]WAY-100635. *Brain Res* **954**: 173–182.
- Pike VW, McCarron JA, Lammertsma AA, Hume SP, Poole K, Grasby PM *et al* (1995). First delineation of 5-HT_{1A} receptors in human brain with PET and [¹¹C]WAY-100635. *Eur J Pharmacol* **283**: R1–R3.
- Rabiner EA, Messa C, Sargent PA, Husted-Kjaer K, Montgomery A, Lawrence AD *et al* (2002). A database of [(11)C]WAY-100635 binding to 5-HT_{1A} receptors in normal male volunteers: normative data and relationship to methodological, demographic, physiological, and behavioral variables. *Neuroimage* **15**: 620–632.
- Rosa-Neto P, Gjedde A, Olsen AK, Jensen SB, Munk OL, Watanabe H *et al* (2004). MDMA-evoked changes in [(11)C]raclopride and [(11)C]NMSP binding in living pig brain. *Synapse* **53**: 222–233.
- Sargent PA, Kjaer KH, Bench CJ, Rabiner EA, Messa C, Meyer J *et al* (2000). Brain serotonin1A receptor binding measured by positron emission tomography with [¹¹C]WAY-100635: effects of depression and antidepressant treatment. *Arch Gen Psychiatry* **57**: 174–180.
- Sheline YI, Mintun MA, Moerlein SM, Snyder AZ (2002). Greater loss of 5-HT(2A) receptors in midlife than in late life. *Am J Psychiatry* **159**: 430–435.
- Talairach J, Tournoux P (1988). *Co-planar Stereotactic atlas of the Human Brain: 3-Dimensional Proportional System: An Approach to Cerebral Imaging*. Thieme: Stuttgart, New York.
- Tauscher J, Kapur S, Verhoeff NP, Hussey DF, Daskalakis ZJ, Tauscher-Wisniewski S *et al* (2002). Brain serotonin 5-HT_{1A} receptor binding in schizophrenia measured by positron emission tomography and [¹¹C]WAY-100635. *Arch Gen Psychiatry* **59**: 514–520.
- Tauscher J, Verhoeff NP, Christensen BK, Hussey D, Meyer JH, Kecojecic A *et al* (2001). Serotonin 5-HT_{1A} receptor binding potential declines with age as measured by [¹¹C]WAY-100635 and PET. *Neuropsychopharmacology* **24**: 522–530.
- Tiihonen J, Keski-Rahkonen A, Loppinen M, Muhonen M, Kajander J, Allonen T *et al* (2004). Brain serotonin 1A receptor binding in bulimia nervosa. *Biol Psychiatry* **55**: 871–873.
- Tsukada H, Kakiuchi T, Nishiyama S, Ohba H, Harada N (2001). Effects of aging on 5-HT_{1A} receptors and their functional response to 5-HT(1a) agonist in the living brain: PET study with [carbonyl-(11)C]WAY-100635 in conscious monkeys. *Synapse* **42**: 242–251.
- Turner MR, Rabiner EA, Hammers A, Al Chalabi A, Grasby PM, Shaw CE *et al* (2005). [¹¹C]-WAY100635 PET demonstrates marked 5-HT_{1A} receptor changes in sporadic ALS. *Brain* **128**: 896–905.
- van Dyck CH, Seibyl JP, Malison RT, Laruelle M, Zoghbi SS, Baldwin RM *et al* (2002). Age-related decline in dopamine transporters: analysis of striatal subregions, nonlinear effects, and hemispheric asymmetries. *Am J Geriatr Psychiatry* **10**: 36–43.
- Wang Y, Chan GL, Holden JE, Dobko T, Mak E, Schulzer M *et al* (1998). Age-dependent decline of dopamine D1 receptors in human brain: a PET study. *Synapse* **30**: 56–61.
- Wilson AA, Inaba T, Fischer N, Dixon LM, Nobrega J, DaSilva JN *et al* (1998). Derivatives of WAY 100635 as potential imaging agents for 5-HT_{1A} receptors: syntheses, radiosyntheses, and *in vitro* and *in vivo* evaluation. *Nucl Med Biol* **25**: 769–776.
- Yamamoto M, Suhara T, Okubo Y, Ichimiya T, Sudo Y, Inoue M *et al* (2002). Age-related decline of serotonin transporters in living human brain of healthy males. *Life Sci* **71**: 751–757.
- Yanai K, Watanabe T, Meguro K, Yokoyama H, Sato I, Sasano H *et al* (1992). Age-dependent decrease in histamine H1 receptor in human brains revealed by PET. *Neuroreport* **3**: 433–436.
- Zhou Y, Endres CJ, Brasic JR, Huang SC, Wong DF (2003). Linear regression with spatial constraint to generate parametric images of ligand-receptor dynamic PET studies with a simplified reference tissue model. *Neuroimage* **18**: 975–989.