

Commentary

The Importance of Animal Models of Decision Making, Gambling, and Related Behaviors: Implications for Translational Research in Addiction

Marc N Potenza^{*,1}

¹Departments of Psychiatry and Child Study Center, Yale University School of Medicine, Connecticut Mental Health Center, New Haven, CT, USA

Neuropsychopharmacology (2009) 34, 2623–2624; doi:10.1038/npp.2009.152

Addictions are arguably the most costly medical disorders, with annual costs to the US society estimated to exceed \$500 billion dollars and to be larger than those associated with stroke, heart disease, and diabetes combined (Uhl and Grow, 2004). Despite the immense costs associated with addictions, relatively few effective treatments exist, particularly for specific addictions such as cocaine dependence for which no pharmacotherapies are currently approved by the US Food and Drug Administration. As such, there exists a significant need for more research into the etiologies and neurobiologies of addictions such that more effective treatment strategies may be developed for these currently costly disorders. Given that addictions, similar to other psychiatric disorders, are complex processes for which etiologies likely involve multiple genetic and environmental determinants that interact in a dynamic manner, using multiple approaches and disciplines in an integrative manner will be important to make substantial inroads into our understanding of addictions and to translate this knowledge into prevention and treatment advances.

Addictions have been conceptualized as disorders of misdirected motivation in which one typically hedonic and maladaptive behavior (eg, drug use) takes precedence over other more adaptive ones (eg, performing well at work or attending to familial concerns) (Potenza, 2006). This conceptualization is consistent with descriptions of the core components of addictions that typically include an element of continued engagement in a behavior despite adverse consequences (Potenza, 2006). Consistently, much research over the past decade has involved assessments of decision making in individuals with addictions (Bechara, 2003). Neuroeconomic paradigms have investigated temporal discounting of monetary rewards (as well as drug

rewards) in individuals with addictions and have found that individuals with addictions tend to discount rewards more rapidly than do those without addictions, and that the discounting behaviors are associated with clinically relevant features, such as treatment outcome (Krishnan-Sarin *et al*, 2007). More complex paradigms, such as the Iowa Gambling Task (IGT) that involves risk-reward decision making coupled with the learning of a strategy that is not intuitively obvious, have also demonstrated deficits in individuals with addictions (Bechara, 2003). Moreover, an association between IGT performance and real-life measures of functioning (eg, ability to hold a job) has been reported in addicted individuals, highlighting the real-life relevance of performance on this task (Bechara, 2003). As such, an improved understanding of the neurobiology of decision making holds significant promise for generating clinical advances for the prevention and treatment of addictions.

Precisely how decision making is related to addiction remains relatively poorly understood. That is, individual differences in decision making may relate to genetic and/or early life environmental factors, or their interactions, and may exist before the drug exposure experienced in drug addictions. Alternatively and not mutually exclusively, drug exposure may influence the neural structures and functions underlying decision making. Animal models are important in investigating these questions, and animal data indicate roles for each possibility. For example, in the first instance, individual differences in impulsivity before drug exposure have been linked to drug use behaviors. In a model of cocaine dependence, response impulsivity predicted compulsive drug use (Belin *et al*, 2008), and in investigations of nicotine administration, response impulsivity predicted enhanced motivation to initiate and maintain nicotine self-administration, and impulsive choice on a delayed reward task predicted reinitiation of nicotine use after abstinence (Diergaarde *et al*, 2008). With respect to the second possibility, alcohol exposure during adolescence has been found to lead to risky decision making later in life (Nasrallah *et al*, (in press)), indicating that both individual differences in decision making before substance exposure

*Correspondence: Dr MN Potenza, Departments of Psychiatry and Child Study Center, Yale University School of Medicine, Connecticut Mental Health Center, Room S-104, 34 Park Street, New Haven, CT 06519, USA, Tel: +1 203 9747356, Fax: +1 203 9747366, E-mail: marc.potenza@yale.edu
Received 24 August 2009; accepted 25 August 2009

and substance exposure itself (as well as the timing and duration of such) are important considerations in addictions. These animal studies, that include pre-drug-exposure assessments, controlled experimental conditions, longitudinal characterization, and (in some cases) nondrug-exposed experimental groups, provide important information that could not be obtained in human studies because of the lack of feasibility. As such, they offer a critical complement to human investigations.

The recent investigation by Zeeb *et al* (2009), as well as another recent one by Rivalan *et al* (2009), models the IGT in rats. Both groups use the task to investigate important aspects of decision making: influences on task performance of serotonergic and dopaminergic drugs as well as individual differences in reward sensitivity and anxiogenic risk taking, respectively. Although each study provides important information in its own right, there is significant hope that they represent the initial basis for multiple investigations that deconstruct and probe specific aspects of decision making as related to addiction. As evidenced by other examples mentioned above, such investigations could use carefully controlled longitudinal experimental conditions that are not practical or feasible in human studies. Such investigations could also adopt additional approaches (eg, genetic manipulations, proteomic analyses, electrophysiological recordings, or neurochemical assessments) that are also typically not feasible in people, yet provide important information on neurobiological processes. In addition, careful coordination between preclinical and clinical investigations, such as those that model the same behavior or paradigm (herein, the IGT) across species could facilitate the translation of information in a manner that could help dissect and clarify complex processes underlying addiction and generate clinical advances. Importantly, the availability of animal models of the IGT also holds promise for better understanding gambling behaviors and human conditions such as pathological gambling, for which the lack of availability of appropriate animals models has slowed research into the disorder. As such, these studies hold significant implications of a broad range of neuropsychiatric conditions and thus could contribute to multiple clinical advances that could benefit many individuals, their families, and the society as a whole.

ACKNOWLEDGEMENTS

This study was funded in part by NIH Grant nos R01 DA019039, R01 DA020908, RL1 AA017539, and UL1 DE19586, the NIH Roadmap for Medical Research/Common Fund, the VA VISN1 MIRECC, Women's Health Research at Yale, and a Center of Research Excellence Grant from the National Center for Responsible Gaming. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the National Center for Responsible Gaming or the Institute for Research on Gambling Disorders or any of the other funding agencies.

DISCLOSURE

Dr Potenza reports that he has no conflicts of interest over the past 5 years to report as related to the subject of the report. Dr Potenza has received financial support or compensation for the following: Dr Potenza consults for and is an advisor to Boehringer Ingelheim; has consulted for and has financial interests in Somaxon; has received research support from the National Institutes of Health, Veterans Administration, Mohegan Sun Casino, the National Center for Responsible Gaming and its affiliated Institute for Research on Gambling Disorders, Forest Laboratories, Ortho-McNeil, Oy-Control/Biotie and GlaxoSmithKline pharmaceuticals; has participated in surveys, mailings or telephone consultations related to drug addiction, impulse control disorders or other health topics; has consulted for law offices and the federal public defender's office in issues related to impulse control disorders; has performed grant reviews for the National Institutes of Health and other agencies; has given academic lectures in grand rounds, CME events, and other clinical or scientific venues; has generated books or book chapters for publishers of mental health texts; and provides clinical care in the Connecticut Department of Mental Health and Addiction Services Problem Gambling Services Program.

REFERENCES

- Bechara A (2003). Risky business: emotion, decision-making, and addiction. *J Gambling Stud* 19: 23–51.
- Belin D, Mar AC, Dalley JW, Robbins TW, Everitt BJ (2008). High impulsivity predicts the switch to compulsive cocaine-taking. *Science* 320: 1352–1355.
- Diergaarde L, Pattij T, Poortvliet I, Hogenboom F, de Vries W, Schoffelmeer ANM *et al* (2008). Impulsive choice and impulsive action predict vulnerability to distinct stages of nicotine seeking in rats. *Biol Psychiatry* 63: 301–308.
- Krishnan-Sarin S, Reynolds B, Duhig AM, Smith A, Liss T, McFertridge A *et al* (2007). Behavioral impulsivity predicts treatment outcome in a smoking cessation program for adolescent smokers. *Drug Alc Dep* 88: 79–82.
- Nasrallah NA, Yang TWH, Bernstein IL (in press) Long-term risk preference and suboptimal decision making following adolescent alcohol use. *Proc Natl Acad Sci USA*.
- Potenza MN (2006). Should addictive disorders include non-substance-related conditions? *Addiction* 101(Suppl 1): 142–151.
- Rivalan M, Ahmed SA, Dellu-Hagedorn F (2009). Risk-prone individuals prefer the wrong options on a rat version of the Iowa Gambling Task. *Biol Psychiatry*. [published online ahead of print May 30]; <http://journals.elsevierhealth.com/periodicals/bps/inpress>
- Uhl GR, Grow RW (2004). The burden of complex genetics in brain disorders. *Arch Gen Psychiatry* 61: 223–229.
- Zeeb FD, Robbins TW, Winstanley CA (2009). Serotonergic and dopaminergic modulation of gambling behavior as assessed using a novel rat gambling task. *Neuropsychopharmacol* 34: 2329–2343.

See related article by Zeeb *et al* on page 2329 in NPP, Volume 34, Number 10.