

Erratum

Noradrenergic $\alpha 1$ Receptors as a Novel Target for the Treatment of Nicotine Addiction

Benoit Forget, Carrie Wertheim, Paola Mascia, Abhiram Pushparaj, Steven R. Goldberg and Bernard Le Foll

Neuropsychopharmacology (2010) 35, 2006; doi:10.1038/npp.2010.62

Correction to: Neuropsychopharmacology advance online publication, 31 March 2010; doi:10.1038/npp.2010.42

In this article, the data below Table 2 are incorrect. Table 2 is republished here with the correct data.

Table 2 Response Rates of Rats During the Drug Discrimination Study at Different Nicotine Doses (0–0.4 mg/kg) After Prazosin (0–1 mg/kg) Pretreatment

	Nicotine dose (mg/kg)					
	0	0.01	0.03	0.1	0.3	0.4
Vehicle	2.1 ± 0.3	1.9 ± 0.2	2 ± 0.3	2 ± 0.2	2.2 ± 0.2	2 ± 0.3
Prazosin 0.25 mg/kg	1.8 ± 0.2	1.7 ± 0.3	1.7 ± 0.2	1.9 ± 0.2	2.3 ± 0.2	1.5 ± 0.3
Prazosin 0.5 mg/kg	2 ± 0.2	1.3 ± 0.2	1.7 ± 0.2	1.6 ± 0.2	1.8 ± 0.3	1.7 ± 0.3
Prazosin 1 mg/kg	1.7 ± 0.2	1.7 ± 0.1	2 ± 0.3	1.8 ± 0.2	2 ± 0.2	1.6 ± 0.2
ANOVA $F_{3,3} =$	1.7, NS	1.75, NS	1.95, NS	1.3, NS	1.4, NS	1.3, NS

Data are expressed as rate of responding on levers (in lever presses/sec ± SEM).