

Index

Note: Page numbers of article titles are in **bold face** type.

A

- Abscess, brain, diffusion MRI in, 117
- Adenomas, pituitary, transsphenoidal surgery for.
 See Transsphenoidal pituitary surgery.
- Alignment stem, in stereotactic brain biopsy,
 167–168
- Anesthesia, for transsphenoidal pituitary surgery,
 with intraoperative MRI, 159
- Arachnoid cyst, diffusion MRI in, 117
- Archicerebellum, anatomy of, 77–78, 80
- Astrocytoma, pilocytic, MRI in, at 0.12 T,
 150–151

B

- B1 field, in MRI, creation of, 6–7
- B1 vector, in MRI, creation of, 7–8
- Biopsy, brain. *See* Brain, biopsy of.
- Blood-brain barrier, opening of, to targeted
 therapy, 209
- Boltzmann equation, 4
- Bone, brightness of, in MRI, 10
- Brain
 - abscess of, diffusion MRI in, 117
 - arachnoid cyst of, diffusion MRI in, 117
 - biopsy of, MRI in, **165–172**
 - historical view of, 165–166
 - in burr hole procedures, 191–192
 - prospective stereotaxy in, 167–168
 - results of, 168–171
 - technical advances in, 171
 - technique for, 166–170
 - versus MR spectroscopy, 168–170
 - epidermoid tumors of, diffusion MRI in,
 117
 - tumors of. *See* Brain tumors.
- Brain shift phenomenon, MRI in, 65–75, 136,
 205–206

Brain tumors

- cellularity of, diffusion MRI in, 118, 120
- diffusion MRI in, 116–120
- diffusion tensor MRI in, 121–131
 - for grading, 122–123
 - for presurgical planning, 123–125
 - functional MRI with, 125–127
 - intraoperative utility of, 128–130
 - limitations of, 131
 - therapeutic, 130–131
- grading of, diffusion MRI in, 122–123
- MR spectroscopy in, **101–114**
 - accuracy of, 101
 - applications of, 103–104
 - case study of, 105–106
 - coregistration of metabolic and anatomic
 MRI in, 104–105
 - data processing in, 104, 108, 110
 - principles of, 101–103
 - procedure for, 104–105
 - results of, 105–107, 110–111
 - three-dimensional (at 3T), 107–111
- MRI in
 - at 0.12 T, 147–151
 - grid computing in, 65–75
- staging of, diffusion MRI in, 118
- treatment of, diffusion tensor MRI in, 130–131

Brigham and Women's Hospital Magnetic
 Resonance Therapy Unit, glioma experience
 of, 137–138

Burr hole procedures, MRI in, 191–192

C

- Carr-Purcell-Meiboom-Gill pulse sequence, in
 MRI, 19, 24, 26–27
- Cavitation, for blood-brain barrier opening, to
 targeted therapy, 209
- Cerebellum, functional MRI of, **77–99**
 - active versus passive movement in, 84, 88
 - anatomic considerations in, 77–78, 80
 - clinical perspective in, 95–98
 - cognitive processing in, 92, 94–95, 97

Cerebellum (*continued*)

- cortical topography in, 84, 86
- executed versus imagined movement in, 84, 87
- historical view of, 77, 79, 81, 83
- in topography, 81–84, 86
- individual variations in, 83–85
- language processing in, 92, 94–95, 97
- neurologic circuits and, 77
- projections in, 83
- somatosensory cancellation in, 92
- somatotropic map of, 81–84, 86
- spatial normalization in, 83–85
- voluntary movement frequency in, 87, 89–90
- voluntary movement timing in, 89, 91, 93

Cerebrospinal fluid, MRI of, 16

Chemical selective saturation, in MRI, 34–36

Climbing fibers, cerebellum input from, 77

Cognitive processing, in cerebellum, functional MRI of, 92, 94–95, 97

Computed tomography, intraoperative
in epilepsy, 174
versus intraoperative MRI, 152

Craniopharyngioma, MRI in, at 0.12 T, 147–149

Cyst, arachnoid, diffusion MRI in, 117

D

Diamagnetic materials, in MRI, 5

Diffusion MRI, 115–134

- in brain tumors, 116–120
- tensor, 121–131
- physical basis of, 115–116

Diffusion tensor MRI

- in brain tumors, 121–131
- for grading, 122–123
- for presurgical planning, 123–125
- functional MRI with, 125–127
- intraoperative utility of, 128–130
- limitations of, 131
- therapeutic, 130–131
- physical basis of, 120–121

Dipoles

- energy transition of, 4
- large single magnetic dipole formed from, 7–8
- net vector of, 5–6
- phase coherence of, 4–5
- precession of, 2

“Double doughnut” system, for intraoperative MRI, 202

E

Echo time, in MRI, 45

Enthalpy, in T1 relaxation, 11

Epidermoid tumors, brain, diffusion MRI in, 117

Epilepsy, 173–183

- definition of, 173
- surgery for
 - efficacy of, 173–174
 - history of, 174
- MRI in, **173–183**, 193
 - advantages of, 180
 - current status of, 178–180
 - patient selection for, 176
 - results of, 176–178
 - technology of, 174–175
 - University of Calgary experience in, 180–182
 - treatment of, 173

Exponential decay components, in MRI, 18

F**Fast spin echo technique, in MRI, 54–60**

- clinical applications of, 59–60
- disadvantages of, 58–59
- image contrast in, 56–57
- J coupling in, 60
- T1-weighted, 57–58
- T2-weighted, 58

Fat molecules, in MRI

- brightness of, 10, 14
- chemical saturation of, 36
- J coupling of, 60
- precession of, 13–14

Ferromagnetic materials, in MRI, 5, 15

FLAIR (fluid-attenuated inversion recovery)
pulse sequence, in MRI, 24, 33–34

Fluid-attenuated inversion recovery (FLAIR)
pulse sequence, in MRI, 24, 33–34

Focused ultrasound surgery, MRI-guided,
208–209

Fourier transform, in image reconstruction, in MRI, 46, 52, 54

Free induction decay, in MRI, 7–8, 19, 44

Frequency encoding, in MRI, 45–46

FSE (fast spin echo) technique, in MRI, 54–60

Functional MRI. *See* Magnetic resonance imaging, functional.

G

GEMSS (grid-enabled medical simulation services) project, MRI with grid computing, 65–75

Glioma surgery

diffusion tensor MRI in, 124–125

MR spectroscopy in, 104–107

MRI in, **135–141**, 189, 192–193

advantages of, 135–137

clinical experience of, 137–138

complications of, 138

cortical identification in, 136–137

safety of, 138

tumor resection extent and, 138

versus other modalities, 138–140

Gliomatosis cerebri, diffusion tensor MRI in, 124–125

Gradient echo, in MRI, 49–51

Gradient-recalled acquisition in the steady state (GRASS) pulse sequence, in MRI, 19, 24, 37–38

GRASS (gradient-recalled acquisition in the steady state) pulse sequence, in MRI, 19, 24, 37–38

Grid computing, in MRI, **65–75**

chain execution in, 72–73

description of, 66–68

image processing chain in, 68–72

need for, 65–66

performance of, 73–74

quality of service from, 73

Grid-enabled medical simulation services (GEMSS) project, MRI with grid computing, 65–75

H

Hahn echo, in MRI, 23

Half-Fourier single-shot turbo spin echo

(HASTE) sequences

in brain biopsy, 166

in transsphenoidal surgery, 188–189

HASTE (half-Fourier single-shot turbo spin echo) sequence

in brain biopsy, 188–189

in transsphenoidal surgery, 188–189

Hemorrhage, intracerebral, burr hole procedures for, MRI in, 191–192

Hitachi AIRIS II MRI unit, in transsphenoidal pituitary surgery, 158–159

Hydrogen

precession of, 2

spin density of, 17–18

I

Image processing chain, in MRI, with grid computing, 68–72

data transfer and conversion in, 69

deformation field in, 72

executing over grid, 72–73

intensity adjustment in, 71

intensity nonuniformity correction in, 69–71

linear registration n, 71

nonlinear registration in, 71

overlay of data in, 72

requirements for, 68

security of, 68

steps of, 68

transfer to presentation device in, 72

Image reconstruction, in MRI, 46–47

Intraoperative MRI. *See* Magnetic resonance imaging, intraoperative.

Inversion recovery (IR) pulse sequence, in MRI, 19, 24, 28–34

conventional, 28–30

fluid-attenuated (FLAIR), 24, 33–34

short time (STIR), 19, 24, 32–33

with short repetition time, 31–32

Ischemia, cerebral, intraoperative detection of, MRI in, 130

J

J coupling, in MRI, 60

Joubert syndrome, cerebellar vermis

underdevelopment in, functional MRI in, 96–98

L

Language processing, in cerebellum, functional MRI of, 92, 94–95, 97

Larmour frequency, in MRI, 2, 9–10

Larsell classification, of cerebellar anatomic assignments, 79–81

Limited flip angle imaging, in MRI, 38, 40

Lymphoma, brain, diffusion MRI in, 118, 120, 125

M

Magnetic fields, in T2 relaxation, 12–14

Magnetic resonance imaging

functional

diffusion tensor MRI with, in brain tumors, 125–127

in neuronavigation, 195–196

of cerebellum, **77–99**

intraoperative, **185–200**

benefits of, 204–207

clinical experience with, 189–194

development of, 185–186

diffusion. *See* Diffusion MRI; Diffusion tensor MRI.

“double doughnut” system for, 202

equipment for, 186–189

for neuronavigation, 189

future of, **201–213**

grid computing in, **65–75**

head holder for, 188

in brain biopsy, **165–172**

in brain tumors, **101–114**

in burr hole procedure, 191–192

in epilepsy, **173–183**, 193

in glioma surgery, **135–141**, 192–193

in neuronavigation. *See* Neuronavigation, MRI in.

in therapy delivery systems, 209–210

in thermal ablation, 208–209

in transsphenoidal pituitary surgery, **155–164**, 190–191

of cerebellum, functional, **77–99**

operating room setup for, 186–189

paradigms of, 202–204

PoleStar 0.12T unit for, **143–154**

types of, 202

unresolved issues in, 207–208

versus intraoperative computed tomography, 152

principles of, **1–63**

basic pulse sequences, 19–24

Carr-Purcell-Meiboom-Gill sequence, 19, 24, 26–27

chemical selective saturation, 34–36

coherence of fast tissues, 16–17

echo time, 45

fast spin echo technique, 54–60

frequency encoding, 45–46

gradient echo, 19, 24, 37–38, 49–51

hydrogen availability, 17–19

image reconstruction, 46–47

inversion recovery, 28–34

limited flip angle imaging, 38, 40

multislice acquisition, 47–48

net vector, 5–6

phase coherence, 4–5

phase encoding, 41–45

precession, 2

prescanning, 43–44

proton density weighting, 23

pulse preparation, 36–37

selective excitation, 41–43

signal formation, 6–8

single-slice mode, 48

spatial encoding, 41

spin echo, single-slice, two-dimensional data acquisition, 41

spin echo pulse. *See* Spin echo (SE) pulse sequence.

T1 relaxation. *See* T1 relaxation.

T1 weighting, 25, 57–58

T2 relaxation. *See* T2 relaxation.

T2 weighting, 25–26, 58

thermal motion, 3–4

three-dimensional data acquisition, 51–54

Magnetic resonance spectroscopy, intraoperative

in brain biopsy

historical view of, 165–166

procedure for, 168

results of, 168–170

versus MRI-guided brain biopsy, 168–170

in brain tumors, **101–114**

Magnetization vector, recovery of, to ground state. *See* T1 relaxation.

Magnetoencephalography, functional MRI with, in brain tumors, 106–107

Meningioma, MRI in, diffusion, 120

Metastasis, to brain, diffusion MRI in, 125

Mossy fibers, cerebellum input from, 77

Movement, cerebellar control of, functional MRI of

active versus passive, 84, 88

executed versus imagined, 84, 87

frequency of, 87, 89–90

somatosensory cancellation in, 92

timing of, 89, 91–93

MRI. *See* Magnetic resonance imaging.

Multislice acquisition, in MRI, 47–48

N

Navigation. *See* Neuronavigation.

Navigus trajectory guide, for brain biopsy, 166

Neocerebellum, anatomy of, 77–79

Net vector, in MRI, 5–6

Neuronavigation, MRI in, 189

equipment for, 189

for anatomic guidance, 194–195

for brain biopsy, 165–172

for brain tumors, 106–107

for epilepsy surgery, 181

functional, 195–196

in PoleStar system, 143, 146

Normalization, of cerebellar individual differences, 83–85

Number of excitations, in MRI, 23

O

Oligoastrocytoma, MR spectroscopy in, 106

Oligodendroglioma, MRI in, at 0.12 T, 147–149

P

Paleocerebellum, anatomy of, 77–80

Phase angle, in MRI, 45

Phase coherence, in MRI, 4–5

establishment of, 6–8

loss of. *See* T2 relaxation.

Phase encoding, in MRI, 44–45

Pilocytic astrocytoma, MRI in, at 0.12 T, 150–151

Pituitary tumors, transsphenoidal surgery for.
See Transsphenoidal pituitary surgery.

Pivot point, in stereotactic brain biopsy, 167

Pixels, in MRI

in three-dimensional data acquisition, 52–54
versus voxels, 41

PoleStar intraoperative MRI (0.12 T), **143–154**

case illustrations of, 147–151

necessity of, 150, 152–153

specifications of, 143–145

stereotactic accuracy of, 145

technique for, 145–147

Posterior vermal split syndrome, in cerebellum,
functional MRI in, 96

Precession, in MRI, 2

Prescanning, in MRI, 43–44

Prospective stereotaxy, in MRI, in brain biopsy,
167–168, 171

Proteins, in MRI, brightness of, 10

Proton(s)

energy transition of, 4–5

precession of, 2

relaxation of. *See* T1 relaxation; T2 relaxation.

Proton density weighting, in MRI, 23

Pulse sequences, in MRI, 19–24

acronyms for, 20–23

phases of, 19

preparation of, 36–37

R

Radiofrequency energy

B1 field created by, 6–7

description of, 5–6

net vector and, 4

Registration, in grid computing, for MRI, 71

Relaxation, in MRI. *See* T1 relaxation; T2
relaxation.

Repetition times (TR), in MRI, 11

alteration of, 25

fast spin echo, 54

in T1 relaxation, 11

short, 47–48

total imaging time and, 31

Resynchronization (spin locking), in MRI,
7–8

S

Seizures, in epilepsy. *See* Epilepsy.

Selective excitation, in MRI, 41–43

Short time inversion recovery (STIR) pulse
sequence, in MRI, 19, 24, 32–33

Signal generation, in MRI, 6–8
net vector in, 5–6

Signal intensity, in MRI, 18

Single-slice mode, in MRI, 48

Single-voxel MR spectroscopy, in brain biopsy,
168

Somatosensory cancellation, cerebellar function
in, functional MRI of, 92

Somatotropy, of cerebellum, functional MRI of, 81–84, 86

Spectroscopy, magnetic resonance. *See* Magnetic resonance spectroscopy.

Speech production, in cerebellum, functional MRI of, 92

SPGR (spoiled grass) pulse sequence, in MRI, 24

Spin echo (SE) pulse sequence, in MRI, 19, 23
fast, 54–60
prescanning in, 43–44
with single-slice, two-dimensional data acquisition, 41

Spin lattice relaxation time, in MRI. *See* T1 relaxation.

Spin locking, in MRI, 7–8

Spin-spin dephasing, in MRI. *See* T2 relaxation.

Spoiled grass (SPGR) pulse sequence, in MRI, 24

SSFP (steady-state free precession) pulse sequence, in MRI, 19

Static magnetic fields, in T2 relaxation, 12–13

Steady-state free precession (SSFP) pulse sequence, in MRI, 19

Stereotaxy, in MRI, 145
prospective, in brain biopsy, 167–168, 171

STIR (short time inversion recovery) pulse sequence, in MRI, 19, 24, 32–33

T

T1 relaxation, in MRI, 8–11, 15
equation for, 18
for selected tissues, 12–14
versus T2 relaxation, 14–15

T1 weighting, in MRI, 25
fast spin echo, 57–58

T2 relaxation, in MRI, 12–14
equation for, 18
for selected tissues, 12–14
static magnetic fields and, 12–13
time-varying magnetic fields and, 13–14
versus T1 relaxation, 14–15

T2 weighting, in MRI, 25–26
fast spin echo, 58

Talairach space, in cerebellum, normalization of, in function MRI, 83–85

Target point, in stereotactic brain biopsy, 167

Thermal ablation, MRI-guided, 208–209

Thermal motion, in MRI, 3–4

3D Slicer, for intraoperative MRI, 202–203

Three-dimensional data acquisition
in MR spectroscopy, 107–111
in MRI, 51–54

Time of repetition (TR), in MRI, 11, 19

Time-varying magnetic fields, in T2 relaxation, 13–14

TR. *See* Repetition times (TR).

Transsphenoidal pituitary surgery, MRI in, **155–164**, 190–191
anesthesia for, 159
drawbacks of, 161–162
half-Fourier single-shot turbo spin echo (HASTE), 188–189
literature review on, 156
low-field Hitachi magnet in, 158–159
microsurgical technique in, 159–160
operating room setup for, 157–158
positioning for, 159
results of, 160–162
technologies for, 156–157
twin operating theater concept in, 157–158

Tumor(s)
brain. *See* Brain tumors; *specific tumors*.
pituitary, transsphenoidal surgery for.
See Transsphenoidal pituitary surgery.

Turbo spectroscopic MRI, in brain biopsy, 168

U

Ultrasound waves, focused, for brain surgery, MRI-guided, 208–209

University of Calgary, epilepsy surgery in, with intraoperative MRI, 180–182

University of Cincinnati, intraoperative MRI experience at, 157–162

V

Vascular protection, intraoperative MRI for, 205

Vectors, in MRI, net, 5–6

Voxels, in MRI
in three-dimensional data acquisition, 52–54
versus pixels, 41

W

Water molecules, in MRI

brightness of, 9–10, 13–14

diffusion tensor, 120–121

precession of, 13–14

White matter

diffusion tensor MRI of, 123–124

MR spectroscopy of, 110