

Isomerization of allylic alcohols to carbonyl compounds by aqueous-biphase rhodium catalysis

Claudio Bianchini,* Andrea Meli and Werner Oberhauser

Istituto per lo Studio della Stereochimica ed Energetica dei Composti di Coordinazione, ISSECC-CNR, 50132 Firenze, Italy. E-mail: bianchin@fi.cnr.it

Received (in Montpellier, France) 18th February 2000, Accepted 21st April 2000


First published as an Advance Article on the web 14th June 2000

Isomerization of allylic and homoallylic alcohols is catalyzed by the zwitterionic Rh(I) complex (sulphos)Rh(cod) in water–*n*-octane to give the corresponding aldehyde or ketone in high yields and chemoselectivity. A π -allyl metal hydride mechanism is proposed on the basis of various independent experiments in both homogeneous and biphasic systems [sulphos = $-\text{O}_3\text{S}(\text{C}_6\text{H}_4)\text{CH}_2\text{C}(\text{CH}_2\text{PPh}_2)_3$].

The isomerization of allylic alcohols to carbonyl compounds is an important reaction that can be accomplished with homogeneous,¹ heterogeneous² and phase-transfer catalysts.³ Serious drawbacks have often been encountered in the application of this reaction to obtain synthetically useful quantities of either aldehydes or ketones, however. Most common drawbacks are: (i) difficult separation of the catalyst from product(s)/reagent(s);^{1,3} (ii) low chemoselectivity, generally due to competitive acetalization and dehydration reactions;^{1,2} (iii) sensitivity to the degree of substitution on the double bond;¹ (iv) possible catalyst deactivation *via* decarbonylation of aldehydes to give stable carbonyl metal compounds.^{1,4}

The present paper constitutes a preliminary account of the first example of effective aqueous-biphase isomerization of allylic alcohols catalyzed by a water-soluble transition metal complex in a chemoselective manner. A comparative, *in situ* and reactor, study in the homogeneous phase has provided valuable mechanistic information, particularly on the pathways to catalyst deactivation. Prior to our work, the isomerization of geraniol to citronellal was attempted in toluene–water using $\text{Ni}(\text{cod})_2\text{-dppbt}$ [tetrasulfonated 1,4-bis-(diphenylphosphino)butane]. Very low conversion (1%) and selectivity (3%) were obtained, however.⁵

The zwitterionic rhodium complex (sulphos)Rh(cod) (**1**) is an efficient catalyst for the hydrogenation and hydroformylation of olefins in water–*n*-octane [sulphos = $-\text{O}_3\text{S}(\text{C}_6\text{H}_4)\text{CH}_2\text{C}(\text{CH}_2\text{PPh}_2)_3$].⁶ We show here that the same catalytic system can successfully be employed for the chemoselective transformation of both allylic and homoallylic alcohols into the corresponding aldehyde or ketone. Selected results of aqueous-biphase and homogeneous reactions (in 1,2-dichloroethane) are reported in Table 1.




In biphasic conditions, 100 equiv. of allylic alcohol was converted to propanal within 1 h at 100 °C (entry 1). By simple phase-separation, the aldehyde product (>99% in *n*-octane) was separated from the catalyst, which was recycled three times to give the following turn-over-frequencies [mol product (mol catalyst)⁻¹ h⁻¹]: 490, 430 and 390. Appreciable deactivation (*ca.* 10%) occurred after each run due to partial incorporation of active rhodium into the dicarbonyl complex

(sulphos)Rh(CO)₂ (**2**), which was independently shown to form upon reaction of **1** with propanal (and aldehydes in general) and to be inactive for allylic alcohol isomerization. In homogeneous phase, where the catalyst and the aldehyde product are in the same fluid system (entry 1), the formation of **2** was apparently much faster (NMR evidence) and the catalyst could not be re-used for a second run (90% formation of **2**).

In both homogeneous and biphasic conditions, 2-methyl-2-propen-1-ol (entry 2) and 3-buten-2-ol (entry 3) were isomerized to 2-methylpropanal and methyl ethyl ketone, respectively. The biphasic reactions gave better results in general. Increasing the steric hindrance at the C-3 carbon atom of the allylic moiety as in 2-buten-1-ol (95 : 5 mixture of *E* and *Z* isomers) (entry 4) or 3-methyl-2-buten-1-ol (entry 5) resulted in a progressive decay of the catalytic performance.¹ No catalytic activity was observed when two methyl groups were introduced in the C-1 position as in 1,1-dimethyl-2-propen-1-ol (entry 6). The homoallylic alcohol 3-buten-1-ol was isomerized by **1** to butanal in both phase systems (entry 7). Like the 2-buten-1-ol isomer, the conversion to aldehyde was higher in the homogeneous phase, however.























As a general feature, neither hydrates of aldehydes¹ nor dehydration products^{1,2} were observed in the biphasic reactions, whereas some homogeneous reactions gave dehydration products in the form of free and coordinated butadiene or isoprene. Actually, the complexes (sulphos)Rh(η^4 -butadiene) (**3**) and (sulphos)Rh(η^4 -isoprene) (**4**) were either detected by *in situ* NMR experiments or isolated as termination products of the homogeneous reactions involving 2-buten-1-ol and 3-buten-1-ol or 3-methyl-2-buten-1-ol, respectively.† The transformation of homoallyl alcohol most likely involves its preliminary isomerization to the allylic isomer 2-buten-1-ol. Indeed, **1** is an efficient olefin isomerization catalyst as shown by the conversion of the cod ligand to free cycloocta-1,3-diene⁷ (¹H NMR, GC-MS) as well as the isomerization of allyl ethyl ether to *cis*-propenyl ethyl ether (entry 8).

The two established pathways for transition metal-catalyzed isomerization of allylic alcohols are the *p*-allyl metal



Scheme 1 Proposed mechanism for the isomerization reaction of allyl alcohol to propanal catalyzed by **1**.

Table 1 Homogeneous and liquid-biphase isomerization of allyl alcohols and related compounds catalyzed by **1**^a

Entry	Substrate	Products	Homogeneous yield(%)	Liquid-biphase yield(%)
1			95	100
2			38	73
3			98	100
4			62	30
5			4	<1
			4	<1
			1	<1
			4	<1
			9	<1
6		none		
7			89	33
8			14	
			<1	
			22	
			31	

^a Reaction conditions: autoclave, 100 mL; catalyst, 0.021 mmol; substrate, 2.1 mmol; 1,2-dichloroethane (homogeneous) or 1 : 1 water–*n*-octane (liquid-biphase), 30 mL; 100 °C; 1 h.

hydride^{1c,3} and the metal hydride addition-elimination mechanisms.^{1a,b} A third variation is the “internal redox” mechanism suggested by Trost and Kulawiec that involves the coordination of the allylic alcohol as bidentate ligand.^{1b,f}

The results from the catalytic reactions and from various independent experiments are consistent with the π -allyl metal hydride mechanism shown in Scheme 1, which is quite similar to that proposed by Bergens and Bosnich for [Rh(diphosphine)(solvent)]⁺ catalyst precursors.^{1c} Commonalities between the latter Rh systems and **1** are: (i) the interception of enol intermediates [Z- and E-CH(CH₃)=CHOH]^c along the isomerization of allylic alcohol to propanal; (ii) the selective incorporation of deuterium in the C-2 position of propanal when allylic alcohol is isomerized in D₂O; (iii) the failure to isomerize 1,1-dimethyl-2-propen-1-ol. Unlike the cationic Rh-diphosphine systems, the zwitterionic Rh-sulphos fragment also converts homoallylic alcohol into butanal due to its peculiar ability to isomerize olefins.⁷

In conclusion, we have shown here that aqueous-biphase catalysis can be successfully employed for the isomerization of both allylic and homoallylic alcohols to carbonyl compounds. The main advantage of the present system over homogeneous systems is the facile catalyst recycling by simple phase separation with no loss of rhodium metal.

Further modification of sulphos-based catalysts (different metals, introduction of stereocenters), as well as extension of the range of substrates and prospects for asymmetric induction⁸ remain exciting avenues to explore for the clean and selective transformation of allylic residues into aldehyde or ketone functional groups by aqueous-biphase catalysis.

Acknowledgements

MURST and CNR (legge 95/95) are gratefully acknowledged for financial support.

Notes and references

† Satisfactory elemental analyses were obtained for compounds **3** and **4**.⁹ Selected spectroscopic data for **3**: ³¹P{¹H} NMR (CD₂Cl₂, 81.01 MHz): 25 °C, δ 12 (br, P_A); –30 °C, δ 14.4 [br d, J(P_MRh) = 100 Hz, P_M], 4.5 [dt, J(P_AP) = 22.1 Hz, J(P_ARh) = 153.2 Hz, P_A]; –70 °C, δ 22 (br, P_Q), 8 (br, P_M), 3.8 [br d, J(P_ARh) = 150 Hz, P_A]. ¹H NMR (CD₂Cl₂, 200.13 MHz, 25 °C): η^4 -butadiene hydrogens δ 5.61 (m, 2H, CH Hs), 2.0–1.8 (m, 4H, CH₂). For **4**: ³¹P{¹H} NMR (CD₂Cl₂, 81.01 MHz): 25 °C, δ 11 (br); –30 °C, δ 13.4 (2nd order m, P_A-P_B), δ 1.2 [ddd, J(P_MP) = 23.8, 15.8 Hz, J(P_MRh) = 151.7 Hz, P_M]. ¹H NMR (CD₂Cl₂, 200.13 MHz, 25 °C): η^4 -isoprene hydrogens δ 5.58 [t, 1H, J(HH) = 6.9 Hz, H₃ Hs], 2.15 (m, 2H, H_{1s} + H_{4s}), 1.81 (s, 3H, Me), 1.52 (m, 2H, H_{1a} + H_{4a}).

- (a) D. V. McGrath and R. H. Grubbs, *Organometallics*, 1994, **13**, 224; (b) B. M. Trost and R. J. Kulawiec, *J. Am. Chem. Soc.*, 1993, **115**, 2027; (c) S. H. Bergens and B. Bosnich, *J. Am. Chem. Soc.*, 1991, **113**, 958; (d) C. Bianchini, E. Farnetti, M. Graziani, M. Peruzzini and A. Polo, *Organometallics*, 1993, **12**, 3753; (e) F. P. Pruchnik, P. Smolenski and K. Wajda-Hermanowicz, *J. Organomet. Chem.*, 1998, **570**, 63; (f) C. Slugovc, E. Rüba, R. Schmid and K. Kirchner, *Organometallics*, 1999, **18**, 4230.
- C. Hoang-Van and O. Zegaoui, *Appl. Catal. A*, 1997, **164**, 91.
- H. Alper and K. Hachem, *J. Org. Chem.*, 1980, **45**, 2269.
- (a) J. P. Collman and L. S. Hegedus, *Principles and Applications of Organotransition Metal Chemistry*, University Science Books, Mill Valley, CA, 1980, ch. 9, p. 496; (b) C. M. Beck, S. E. Rathmill, Y. J. Park, J. Chen, R. H. Crabtree, L. M. Liable-Sands and A. L. Rheingold, *Organometallics*, 1999, **18**, 5311; (c) This work.
- H. Bricout, E. Monflier, J.-F. Carpentier and A. Mortreux, *Eur. J. Inorg. Chem.*, 1998, 1739.
- (a) C. Bianchini, P. Frediani and V. Sernau, *Organometallics*, 1995, **14**, 5458; (b) C. Bianchini, A. Meli, V. Patinec, V. Sernau and F. Vizza, *J. Am. Chem. Soc.*, 1997, **119**, 4945.
- Complex **1** catalyzes the isomerization of ca. 90 equiv. of cycloocta-1,5-diene to cycloocta-1,3-diene in water–MeOH–*n*-octane (1 : 1 : 2; v : v : v) at 100 °C in 1 h.
- C. Botteggi and G. Giacomelli, *Gazz. Chim. Ital.*, 1976, **106**, 1131.
- Compounds **3** and **4** were isolated by column chromatography as yellow–orange solids from the catalytic solutions.