

First structurally characterised metal complex of a 2*H*-benzimidazole derivative. Copper mediated synthesis of 2,2-dimethyl-2*H*-benzimidazole from 1,2-phenylenediamine and acetone

Sanchita Hati,^a Goutam K. Patra,^a Jnan P. Naskar,^a Michael G. B. Drew^b and Dipankar Datta^{*a}

^a Department of Inorganic Chemistry, Indian Association for the Cultivation of Science, Calcutta 700 032, India. E-mail: icdd@mahendra.iacs.res.in

^b Department of Chemistry, University of Reading, Whiteknights, Reading RG6 6AD, UK


Received (in Montpellier, France) 8th September 2000, Accepted 7th November 2000

First published as an Advance Article on the web 4th January 2001

2,2-Dimethyl-2*H*-benzimidazole (**1**) was synthesised by exploiting the reaction of acetone with bis(1,2-phenylenediamine)copper(II) perchlorate. Additionally, the X-ray crystal structure of [Cu(**1**)(PPh₃)₂][ClO₄] is reported.

2*H*-Benzimidazole is a tautomer of 1*H*-benzimidazole and has an *ortho*-quinonoid electronic structure. Our AM1 calculations¹ using the standard MOPAC package (version 1.1) shows that in the gas phase 2*H*-benzimidazole is energetically less stable than 1*H*-benzimidazole by 27.5 kcal mol⁻¹. While 1*H*-benzimidazole is not a rare ligand in transition metal chemistry, nothing is known about the metal binding of 2*H*-benzimidazole. Herein we describe, for the first time, two copper(I) complexes of the 2,2-dimethyl derivative of 2*H*-benzimidazole. The structure of one of them has been determined. We also describe a novel viable synthesis of 2,2-dimethyl-2*H*-benzimidazole (**1**).

Reaction of acetone with bis(ethylenediamine)copper(II) perchlorate gives rise to the cationic copper(II) complex of Curtis' macrocycle **A**.² In an attempt to generate **B**, the 1,2-phenylenediamine (OPDA *i.e.* *ortho*-phenylenediamine) counterpart of **A**, which is not yet known, we have reacted acetone with Cu(OPDA)₂(ClO₄)₂. It is found that this reaction in air at room temperature does not yield **B**; it leads instead to a reddish-brown copper(I) complex of **1** of the formulation Cu(**1**)ClO₄ (**2**), sparingly soluble in acetone, in almost quantitative yield and from the filtrate is obtained the perchloric acid salt^{3,4} of 2,3-dihydro-2,2,4-trimethyl-1*H*-1,5-benzodiazepine (**3**) in *ca.* 35% yield (eqn. 1).



Later we have found that reaction 1 also occurs in the absence of air. Complex **2** is stable in air in the solid state for about two weeks. However, in methanol **2** is stable in air for about 2 h only; on standing in air, the reddish-brown colour of the methanol solution slowly disappears becoming almost colour-

less. The identity of the organic fragment in **2** is corroborated by the X-ray crystal structure (Fig. 1) of [Cu(**1**)(PPh₃)₂][ClO₄] (**4**), which is obtained by the smooth reaction of PPh₃ with **2** at room temperature in a 2 : 1 molar ratio in methanol.

Complex **4** is indefinitely stable in air in the solid state as well as in solution. As revealed by the X-ray crystal structure, the copper(I) centre in **4** has a trigonal NP₂ coordination sphere with the interligand angles very close to 120°. The coordination sphere in most tricoordinate copper(I) complexes is found to be somewhat T-shaped. For an example, see ref. 5. Fig. 1 describes the first structurally characterised metal complex of a 2*H*-benzimidazole. It is noted that the corresponding 1*H*-benzimidazole complex is not known. Our bond valence sum calculations^{6,7} show that the Cu–N bond distance [1.997(3) Å] in **4** is quite close to that expected (1.98 Å) for a symmetric CuN₃⁺ moiety. In this context, we mention that while some 2,2-dialkyl derivatives of 2*H*-imidazole are known,⁸ no metal complexes have so far been reported.

Interestingly, when OPDA is refluxed in acetone, one obtains 2-methyl-2-isobutenylbenzimidazoline.⁹ On the other hand, reaction of acetone with OPDA at room temperature in






Fig. 1 The structure of the cation in **4** with ellipsoids at 30% occupancy. Selected bond distances (Å) and angles (°): Cu1–N1 1.997(3), Cu1–P1 2.287(3), Cu1–P2 2.280(2), N1–C2 1.499(5), C2–N3 1.469(5), N3–C4 1.286(5), C4–C5 1.453(6), C5–C6 1.326(7), C6–C7 1.436(7), C7–C8 1.364(7), C8–C9 1.425(6), C9–N1 1.308(5), C4–C9 1.469(5); N1–Cu1–P1 118.09(12), N1–Cu1–P2 119.83(11), P1–Cu1–P2 120.00(4).





Scheme 1

the presence of an acid gives the acid salt of **3**.^{3,4} Thus, the copper atom clearly plays an important role in the formation of **1** here. Since reaction 1 also occurs in an N_2 atmosphere, we can say that copper(II) acts as a one-electron oxidant in reaction 1 and it is not a case^{10–12} of copper-mediated aerial oxidation.

Demetallation of **2** by aqueous ammonia affords **1** in the free state. The overall synthetic procedure is summarised in Scheme 1. The overall yield of **1** in Scheme 1 is *ca.* 60%. Compound **1** is a light reddish-brown liquid. Its 1H and ^{13}C NMR spectra in a 4 : 1 $CDCl_3$ – CCl_4 mixture (Fig. 2) show that **1** has a C_2 axis, as expected. It is extremely unstable in air; it rapidly turns red and solidifies with decomposition. From the decomposed red solid, crystals of OPDA can be isolated by trituration with hot hexane (65–70 °C fraction from petroleum). Previously, **1** has been synthesised by reacting benzofurazan *N*-oxide with 2-nitropropane under basic catalysis, followed by reduction of the resulting 2,2-dimethyl-2*H*-benzimidazole bis-*N*-oxide with $NaBH_4$.^{13,14} However, this procedure is not straightforward and the overall yield was not reported. Moreover,^{13,14} **1** had not been fully characterised.

Because of “blocked tautomerisation”, 2,2-dimethyl-2*H*-benzimidazole is more stable than 2*H*-benzimidazole towards tautomerisation. Our AM1 calculations show that **1** is energetically less stable than its 1*H* tautomer by only 15.6 kcal mol^{–1}. This indicates, within the framework of the AM1 method, a gain of 11.9 kcal mol^{–1} in stability for **1** towards tautomerisation in comparison with 2*H*-benzimidazole.

Thus, we have found a new and easy synthesis of 2,2-dimethyl-2*H*-benzimidazole (**1**) that exploits the reaction between acetone and bis(1,2-phenylenediamine)copper(II) perchlorate. Though in the free state **1** is very unstable in air, it becomes quite stable in air upon binding copper(I). At present we are engaged in an examination of the generality of Scheme 1 and in developing the transition metal chemistry of 2*H*-benzimidazole.




Fig. 2 (a) 1H and (b) ^{13}C NMR spectra (300 MHz; TMS reference) of **1** in a 4 : 1 $CDCl_3$ – CCl_4 mixture. For the labelling scheme of the C atoms, see Fig. 1.

Experimental

Syntheses

Reaction 1. Silky violet $Cu(OPDA)_2(ClO_4)_2$ (3 g, 6.27 mmol), prepared by a reported procedure,¹⁵ was suspended in 75 ml of purified acetone and stirred for 6 h at room temperature with a $CaCl_2$ (fused) guard tube fitted to the reaction vessel. The reaction mixture gradually turned intense red during stirring. The reddish-brown $Cu(I)ClO_4$ (**2**) that appeared was filtered off, washed thoroughly with acetone and dried *in vacuo* over fused $CaCl_2$; yield: 1.9 g (93%). Anal. calc. for $C_9H_{10}ClCuN_2O_4$: C, 34.94; H, 3.26; N, 9.06; Cu, 20.55%. Found: C, 34.90; H, 3.30; N, 9.01; Cu, 20.49%. FTIR (KBr): ν 1620s,br (C=N), 1135vs, 1100vs, 1070vs (ClO_4) cm^{-1} . Diamagnetic: A_M (CH_3OH) 83 $cm^2 \Omega^{-1} mol^{-1}$ (1 : 1 electrolyte). UV-VIS (nujol): λ_{max} 460, 358, 280 nm. UV-VIS (CH_3OH ; $c = 0.50 \times 10^{-3} mol dm^{-3}$): λ_{max} ($\epsilon/dm^3 mol^{-1} cm^{-1}$) 414 (1200), 352 (2800), 270 (3300), 218 (8100) nm. 1H NMR (300 MHz, $(CD_3)_2SO$, TMS): δ 1.57 (s, methyl, 6H), 7.29–7.36 (br, m, phenyl, 4H). ^{13}C NMR (300 MHz, $(CD_3)_2SO$, TMS): δ 22.71, 104.54, 124.87, 137.06, 159.27.

The filtrate obtained after the isolation of **2** was evaporated on a water bath to obtain a dark mass. It was dissolved in 10 ml of hot water and filtered. To the red filtrate, cooled to room temperature, a saturated aqueous solution of $NaClO_4 \cdot H_2O$ was added dropwise with stirring until a crystalline precipitate started to appear. The reaction mixture was left in air for 1 h. The deposited orange-red diamond-like crystals were filtered off, washed with 5 ml of cold water and dried *in vacuo* over fused $CaCl_2$; yield: 0.6 g (34%). The crystals were characterised as $3 \cdot HClO_4$.^{3,4}

[Cu(I)(PPh₃)₂]ClO₄ (4**).** To a suspension of 0.32 g (1 mmol) of **2** in 15 ml of methanol, 0.52 g (2 mmol) of PPh_3 , dissolved in 20 ml of methanol, was added dropwise with constant stirring to obtain a clear yellowish-orange solution. The reaction mixture was left in air overnight. The yellowish-brown compound **4** that precipitated was filtered off, washed with 5 ml of cold methanol and dried *in vacuo* over fused $CaCl_2$; yield: 0.45 g (54%). An additional crop of crystals could be obtained by adding diethyl ether to the filtrate. Anal. calc. for $C_{45}H_{40}ClCuN_2O_4P_2$: C, 64.80; H, 4.84; N, 3.36; Cu, 7.62%. Found: C, 64.72; H, 4.92; N, 3.36; Cu, 7.67%. FTIR (KBr): ν 1610s,br (C=N), 1150–1075vs,br (ClO_4) cm^{-1} . Diamagnetic: A_M (CH_3OH) 91 $cm^2 \Omega^{-1} mol^{-1}$ (1 : 1 electrolyte). UV-VIS (nujol): λ_{max} 465, 360, 290 nm. UV-VIS (CH_3OH ; $c = 0.05 \times 10^{-3} mol dm^{-3}$): λ_{max} ($\epsilon/dm^3 mol^{-1} cm^{-1}$) 345 (7050), 250 (25 800), 222 (66 900) nm. 1H NMR (300 MHz, $(CD_3)_2SO$, TMS): δ 1.37 (s, methyl, 6H), 7.07–7.50 (br, m, phenyl, 34H). ^{13}C NMR (300 MHz, $(CD_3)_2SO$, TMS): δ 22.05, 104.66, 125.83, 129.27, 129.41, 130.76, 132.42, 132.98, 133.86, 135.73, 159.33.

2,2-Dimethyl-2*H*-benzimidazole (1**).** **2** (3 g, 9.3 mmol) was taken up in 75 ml of chloroform and stirred for 5 min. To the resulting dark brown suspension, 50 ml of 25% aqueous ammonia was added and stirred for 1 h at room temperature, after which it was filtered. The blue aqueous layer of the filtrate was discarded. The light reddish-brown layer of the filtrate was collected and evaporated at room temperature under reduced pressure to *ca.* 5 ml, which was then loaded on a neutral alumina column (25 × 2.5 cm) and eluted with a 1 : 9 diethyl ether–petroleum ether mixture. The yellowish fraction (*ca.* 200 ml) was collected, which upon removal of the solvent at room temperature under reduced pressure yielded **1** as a light reddish-brown liquid. (The whole process was completed within 8 h). Yield: 0.9 g (63%). Anal. calc. for $C_9H_{10}N_2$: C, 73.93; H, 6.90; N, 19.16%. Found: C, 73.91; H, 6.88; N, 19.13%. Density: 0.93 g cm^{-3} . EI-MS: m/z 146.1 (1^+ , 100), 131.1 ($1^+ - CH_3$, 90), 105.1 ($1^+ - C_3H_5$, 28), 90.1

($\text{I}^+ - \text{NC}_6\text{H}_3$, 42), 78.1 (C_6H_6^+ , 12), 63.0 (C_5H_3^+ , 6), 51.0 (C_4H_3^+ , 8), 41.0 (C_3H_5^+ , 15%). UV-VIS (CH_3OH): λ_{max} ($\text{g}/\text{dm}^3 \text{ mol}^{-1} \text{ cm}^{-1}$) 354 (2500), 263 (2800), 217 (6000) nm. ^1H NMR (300 MHz, $\text{CDCl}_3\text{-CCl}_4$, TMS): δ 1.54 (s, methyl, 6H), 6.99–7.02 (br, phenyl, 2H), 7.18–7.21 (br, phenyl, 2H). ^{13}C NMR (300 MHz, $\text{CDCl}_3\text{-CCl}_4$, TMS): δ 21.33, 104.14, 125.69, 134.06, 159.04. All studies on **1** were done with freshly prepared samples.

X-Ray crystallography

Single crystals of **4**, which are yellowish-brown in colour, were obtained by indirect diffusion of diethyl ether into a moderately concentrated dichloromethane solution of the complex. The data were collected with Mo-K α radiation using the MARresearch Image Plate system at 293(2) K. The structure was refined on F^2 using SHELXL-93.¹⁶ Final residuals: $R_1 = 0.0599$ and $wR_2 = 0.1591$ for 5304 reflections with $I \geq 2\sigma(I)$; $R_1 = 0.0907$ and $wR_2 = 0.1775$ for all data.

Crystal data. $\text{C}_{45}\text{H}_{40}\text{ClCuN}_2\text{O}_4\text{P}_2$ (**4**): $M_w = 833.72$, monoclinic, space group $C2/c$, $a = 39.72(4)$, $b = 10.198(13)$, $c = 24.59(3)$ Å, $\beta = 123.78(1)^\circ$, $U = 8280(16)$ Å³, $Z = 8$, $\mu = 0.714 \text{ mm}^{-1}$, 12 812 reflections collected, 7360 independent, $R_{\text{int}} = 0.0289$.

CCDC reference number 440/235. See <http://www.rsc.org/suppdata/nj/b0/b0074101/> for crystallographic files in .cif format.

Acknowledgements

M. G. B. D. thanks EPSRC (UK) and the University of Reading for funds for the Image Plate System. D. D. gratefully

acknowledges the financial support received from the Department of Science and Technology, New Delhi, India.

References

- 1 M. J. S. Dewar, E. G. Zoebisch, E. F. Healy and J. J. P. Stewart, *J. Am. Chem. Soc.*, 1985, **107**, 3902.
- 2 N. F. Curtis, *Coord. Chem. Rev.*, 1968, **3**, 3.
- 3 J. P. Naskar, S. Hati, D. Datta, U. Samanta and P. Chakrabarti, *Z. Kristallogr.*, 1998, **213**, 112.
- 4 U. Samanta, P. Chakrabarti, J. P. Naskar, S. Hati and D. Datta, *Indian J. Chem., Sect. A*, 1999, **38**, 553.
- 5 J. P. Naskar, S. Hati, D. Datta and D. A. Tocher, *Chem. Commun.*, 1997, 1319.
- 6 S. Hati and D. Datta, *J. Chem. Soc., Dalton Trans.*, 1995, 1177.
- 7 J. P. Naskar, S. Hati and D. Datta, *Acta Crystallogr., Sect. B*, 1997, **53**, 885.
- 8 A. R. Katritzki and J. M. Lagowski, in *Comprehensive Heterocyclic Chemistry: The Structure, Reactions, Synthesis and Uses of Heterocyclic Compounds*, ed. K. T. Potts, Pergamon, Oxford, 1984, vol. 5, part 4A, ch. 4.01.
- 9 R. C. Elderfield and J. R. McCarthy, *J. Am. Chem. Soc.*, 1951, **73**, 975.
- 10 C. Jallabert, C. Lapinte and H. Riviere, *J. Mol. Catal.*, 1980, **7**, 127.
- 11 C. Jallabert, C. Lapinte and H. Riviere, *J. Mol. Catal.*, 1982, **14**, 75.
- 12 C. Lapinte, G. Meyer, L. Nadjo and H. Riviere, *Nouv. J. Chim.*, 1984, **8**, 165.
- 13 D. W. S. Latham, O. Meth-Cohn and H. Suschitzky, *J. Chem. Soc., Chem. Commun.*, 1972, 1040.
- 14 D. W. S. Latham, O. Meth-Cohn, H. Suschitzky and J. A. L. Herbert, *J. Chem. Soc., Perkin Trans. 1*, 1977, 470.
- 15 E. J. Duff, *J. Chem. Soc. (A)*, 1968, 434.
- 16 G. M. Sheldrick, SHELXL-93, Program for Crystal Structure Refinement, University of Göttingen, Germany, 1993.