

This article was downloaded by:

On: 26 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713597286>

Synthesis of 2'-Deoxyribonucleoside Derivatives of 1-Deazapurine

G. Cristalli^a; S. Vittori^a; A. Eleuteri^a; R. Volpini^a; E. Camaioni^a; G. Lupidi^a

^a Dipartimento di Scienze Chimiche and Dipartimento di biologia Cellulare, Università di Camerino, Camerino, Italy

To cite this Article Cristalli, G. , Vittori, S. , Eleuteri, A. , Volpini, R. , Camaioni, E. and Lupidi, G.(1994) 'Synthesis of 2'-Deoxyribonucleoside Derivatives of 1-Deazapurine', *Nucleosides, Nucleotides and Nucleic Acids*, 13: 1, 835 — 848

To link to this Article: DOI: 10.1080/15257779408013282

URL: <http://dx.doi.org/10.1080/15257779408013282>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

SYNTHESIS OF 2'-DEOXYRIBONUCLEOSIDE DERIVATIVES OF 1-DEAZAPURINE.

G. Cristalli*, S. Vittori, A. Eleuteri, R. Volpini, E. Camaioni, G. Lupidi.

Dipartimento di Scienze Chimiche and Dipartimento di biologia Cellulare, Università di Camerino, 62032 Camerino, Italy.

Abstract. 5,7-Dichloro-3H-imidazo[4,5-b]pyridine (4) is a versatile base which can be coupled with a variety of sugar moieties and transformed in a series of 7-alkyl(aryl)amino-derivatives by reacting with the corresponding amines. In this paper synthesis, structure elucidation and ADA inhibitory activity of 2'-deoxyribonucleoside derivatives of N⁶-substituted 1-deazaapurines are described.

In our effort to synthesize deaza nucleosides endowed with pharmacological activity, we selected 1-deazapurine derivatives since 1-deazaadenosine (1) has been shown to possess cytotoxic activity,¹ to inhibit adenosine deaminase² and platelet aggregation,³ and to act as an agonist of adenosine receptors.⁴

We have already reported the coupling of 7-nitro-3H-imidazo[4,5-b]pyridine (2)¹ and of 5-chloro-7-nitro-3H-imidazo[4,5-b]pyridine (3)⁵ with ribose and 2-deoxyribose derivatives to obtain 6-amino-¹ and 6-hydroxyamino-1-deazapurine nucleosides.^{6,7}

Our approach is now to use 5,7-dichloro-3H-imidazo[4,5-b]pyridine (4)⁵ as a more versatile base which can be coupled with a

This paper is dedicated to the memory of Professor Roland K. Robins.




Figure 1

variety of sugar moieties and transformed in a series of 7-alkyl(aryl)amino derivatives by reacting with the suitable amines.


We previously reported on the synthesis of N⁶-substituted 1-deazaadenosine^{1,4} and in this paper the synthesis of the corresponding 2'-deoxyribonucleosides is described. The preparation of the corresponding 2',3'-dideoxyribonucleosides and their antiviral activity is reported elsewhere.⁸

CHEMISTRY


The synthesis of 5,7-dichloro-3-(2-deoxy-β-D-*erythro*-pentofuranosyl)-3H-imidazo[4,5-b]pyridine (**8**) was carried out according to Scheme I.

Reaction of freshly prepared 1-chloro-2-deoxy-3,5-di-O-p-toluoyl-α-D-*erythro*-pentofuranose⁹ with the sodium salt of 5,7-dichloro-3H-imidazo[4,5-b]pyridine (**4**),⁵ generated in situ by the treatment with sodium hydride in acetonitrile, gave compounds **5**, **6** and **7**. Deblocking of the mixture with methanolic ammonia yielded the dichloro derivatives **8**, **9** and **10** in 49%, 10% and 2% yield, respectively.

Reaction of 5,7-dichloro-3-(2-deoxy-β-D-*erythro*-pentofuranosyl)-3H-imidazo[4,5-b]pyridine (**8**) with the appropriate amine gave the N⁷-substituted-5-chloro-3-(2-deoxy-β-D-*erythro*-pentofuranosyl)-3H-imidazo[4,5-b]pyridines **13a-e** (Scheme II).



Scheme 1



Scheme 11

Catalytic hydrogenolysis of the chlorine atom in **13a-e** with 10% Pd/C in ethanol and 2N NaOH afforded the corresponding derivatives **14a-e**.

Isomer identification

The site of deoxyribosylation was assigned by UV and ¹H NMR data. Since in the literature¹⁰ were reported the UV spectra of 1-methyl-1H-imidazo[4,5-b]pyridine and 3-methyl-3H-imidazo[4,5-b]pyridine, compounds **8** and **9** were dehalogenated in a hydrogen atmosphere using 10% Pd/C in ethanol and 2N NaOH to give the corresponding imidazo[4,5-b]pyridine derivatives **11** and **12**. The UV spectra of compounds **11** and **12** were essentially indistinguishable and strictly similar to that of 3-methyl-3H-imidazo[4,5-b]pyridine (Table 1). The physical data of compound **11** are also in agreement with that of 9-β-D-2'-deoxyribofuranosyl-1-deazapurine obtained by Betbeder et al.¹¹ by enzymatic synthesis.

Moreover, the UV spectra of compounds **8** and **9** were very similar to that of 5,7-dichloro-3-(β-D-ribofuranosyl)-3H-imidazo[4,5-b]pyridine, previously reported by Cline et al.¹², whereas compound **10** showed a spectral profile different from that observed for these three N-3 derivatives (Table 1).

Table 1. UV spectra of 1-deazapurine derivatives.

U V λ_{\max} nm (ϵ)		
Compd	pH 1.0	pH 12.0
8	291 (14200) 252 (11500)	292 (11600) 266 (9000) 255 (7800) sh
9	290 (14100) 253 (11600)	292 (11400) 265 (9100) 254 (7900) sh
10	291 (11900) sh 286 (12000) 245 (8700) sh 237 (8900)	295 (12600) sh 288 (14500) 262 (10000)
5,7-dichloro-3-(β -D-ribofuranosyl)-3-H-imidazo[4,5-b]pyridine ^{1 2}	287 (9000) 254 (6000)	289 (8900) 259 (5900) 254 (5800) sh
11	282 (8900) 276 (14400) 236 (5500)	288 (7200) 282 (8900) 248 (5800)
12	282 (9000) 275 (10500) 236 (5400)	288 (7300) sh 282 (9200) 248 (5900)
3-Methyl-3H-imidazo[4,5-b] pyridine ^{1 0}	281 (8900) 275 (10600) 236 (4500)	288 (6900) sh 282 (9000) 252 (4500)
1-Methyl-1H-imidazo[4,5-b] pyridine ^{1 0}	289 (7600) sh 282 (8700) 277 (7900) 235 (2000)	288 (7900) sh 282 (9700)

Table 2. N.O.E.-data % of compounds **8**, **9** and **10** upon irradiation of H-1' (DMSO-d₆, 25 °C, 300 MHz)

	H-2'a	H-2'b	H-3'	H-4'	H-2
8	a	3.0	a	1.3	3.0
9	3.0	a	1.0	a	3.4
10	a	3.0	a	1.0	<1

a: no detectable intensity enhancement (< 0.5%)


Furthermore standard criteria for assignment of regioisomers in purine 2'-deoxyribonucleosides are reported by Kazimierczuk et al. and by Hildebrand and Wright.^{13,14} According to these authors, N-9 and N-7 purine 2'-deoxyribonucleosides (N-3 and N-1, respectively, in 1-deazapurines) are differentiated by ¹H NMR by the characteristic downfield chemical shifts for the anomeric H-1' and the purine H-8 resonances (H-2 in 1-deazapurines) of the N-7 (N-1) isomer relative to those of the N-9 (N-3) isomer. According to that, the 1-deaza derivatives **8** and **9** exhibited H-2 signal at δ 8.79 and 8.82 respectively and H-1' signal at δ 6.39 and 6.46 respectively, whereas compound **10** showed a singlet at δ 9.01 (H-2) and a triplet at δ 6.76 (H-1').

Anomer identification

The anomeric configuration of compounds **8**, **9** and **10** was assigned applying n.O.e. difference spectroscopy. Saturation of H-1' of **9** resulted in n.O.e.s of the H-2'a and H-3' signals (3.0 % and 1.0 %, respectively) while there was none at H-4' signal, establishing α-D-configuration (Table I). Saturation of H-1' of **8** and **10** yielded n.O.e.s of the H-2'b and H-4' signals (3.0 % and 1.3 %, respectively) while there was none at H-3' signal, establishing β-D-configuration (Table 2).

Furthermore a stronger n.O.e. effect was observed on H-2 when H-1' was irradiated in compounds **8** and **9** by comparison with **10**, confirming N³-glycosylation for both **5** and **6**. This observation is in

Table 3. ADA inhibitory activity of 1-deazapurine derivatives.

	n	R	R ₁	K _i , μM
	14a	H	H	0.19
	13a	H	Cl	23
	14b	CH ₃	H	0.25
	13b	CH ₃	Cl	4.2
	14c	cC ₅ H ₉	H	100
	13c	cC ₅ H ₉	Cl	>1000
	14d	cC ₆ H ₁₁	H	130
	13d	cC ₆ H ₁₁	Cl	>1000
	14e	CH(CH ₃)CH ₂ C ₆ H ₅	H	150
	13e	CH(CH ₃)CH ₂ C ₆ H ₅	Cl	>1000

agreement with the data reported by Seela et al.¹⁵ in the case of 3-deaza-2'-deoxyadenosine.

On these basis, compounds **8**, **9** and **10** were identified as 5,7-dichloro-3-(2-deoxy-β-D-*erythro*-pentofuranosyl)-3H-imidazo[4,5-b]pyridine, 5,7-dichloro-3-(2-deoxy-α-D-*erythro*-pentofuranosyl)-3H-imidazo[4,5-b]pyridine and 5,7-dichloro-1-(2-deoxy-β-D-*erythro*-pentofuranosyl)-1H-imidazo[4,5-b]pyridine, respectively

BIOLOGICAL EVALUATION

The synthesized nucleosides were evaluated as inhibitors of adenosine deaminase from calf intestine and the results are reported in Table 3.

None of the tested compounds proved to be substrate of ADA and some of them have good inhibitory activity. They showed the same structure activity relationships as the corresponding ribose derivatives.¹⁷

1-Deaza-2'-deoxyadenosine (**14a**) is the most potent in the series ($K_i = 0.19 \mu\text{M}$) and resulted to be more active than 1-deazaadenosine itself ($K_i = 0.66 \mu\text{M}$). The presence of a chlorine atom in position 2 produced a decrease in ADA inhibitory activity. Accordingly, the

introduction of a chlorine atom in the same position of substrates made the compounds more resistant to the enzyme. Substitution on the N⁶ amino group did not produce a decrease in activity only when small groups are introduced (**13b** K_i = 4.2 μM and **14b** K_i = 0.25 μM). Increased lipophilicity and steric hindrance of substituents resulted in derivatives much less active than the parent compound **14a**.

EXPERIMENTAL SECTION

Chemistry.

Melting points were determined with a Buchi apparatus and are uncorrected. ¹H NMR spectra were obtained with a Varian VXR 300 MHz spectrometer. UV spectra were recorded on a Perkin-Elmer Coleman 575 spectrophotometer. TLC were carried out on pre-coated TLC plates with silica gel 60 F-254 (Merck). For column chromatography, silica gel 60 (Merck) was used. Elemental analyses were determined on Carlo Erba model 1106 analyser.

5,7-Dichloro-3-(2-deoxy-3,5-di-O-p-toluoyl-β-D-erythro-pentofuranosyl)-3H-imidazo[4,5-b]pyridine (5),
5,7-dichloro-3-(2-deoxy-3,5-di-O-p-toluoyl-α-D-erythro-pentofuranosyl)-3H-imidazo[4,5-b]pyridine (6),
5,7-dichloro-1-(2-deoxy-3,5-di-O-p-toluoyl-β-D-erythro-pentofuranosyl)-1H-imidazo[4,5-b]pyridine (7).

To a suspension of 1.2 g (6.28 mmol) of 5,7-dichloro-3H-imidazo[4,5-b]pyridine (**4**)⁵ in 30 mL of dry acetonitrile under an atmosphere of dry N₂ was added 300 mg of 60% NaH and the mixture was stirred at room temperature for 30 min. To the ice-cooled mixture was added 3.6 g (9.25 mmol) of freshly prepared 1-chloro-2-deoxy-3,5-di-O-p-toluoyl-D-erythro-pentofuranose⁹ and the suspension was stirred at 0 °C for 3 h. The reaction mixture was filtered through Celite to remove the insoluble material and the filtrate was concentrate to a residue which was used without further purification for the next reaction.

Analytical samples of **5** and **6** were obtained by flash chromatography eluting with C₆H₁₂-AcOEt from (90:10) to (50:50). (**5**): mp 146-148 °C; ¹H NMR (Me₂SO-d₆) δ 2.36 (s, 3H, CH₃), 2.40 (s,

3H, CH₃), 2.80 (m, 1H, H-2'), 3.26 (m, 1H, H-2''), 4.57 (m, 3H, CH₂-5' and H-4'), 5.79 (m, 1H, H-3'), 6.59 (t, 1H, J = 6.9 Hz, H-1'), 7.26 (d, 2H, H-Ph), 7.36 (d, 2H, H-Ph), 7.68 (s, 1H, H-6), 7.78 (d, 2H, H-Ph), 7.95 (d, 2H, H-Ph), 8.80 (s, 1H, H-2). Anal. Calcd. for C₂₇H₂₃Cl₂N₃O₅: C, 60.01; H, 4.29; N, 7.78. Found: C, 60.37; H, 4.10; N, 7.47.

(6): ¹H NMR (Me₂SO-d₆) δ 2.38 (s, 3H, CH₃), 2.41 (s, 3H, CH₃), 3.09 (m, 2H, H-2' and H-2''), 4.55 (m, 2H, CH₂-5'), 5.04 (m, 1H, H-4'), 5.66 (m, 1H, H-3'), 6.70 ("t", 1H, H-1'), 7.28 (d, 2H, H-Ph), 7.38 (d, 2H, H-Ph), 7.60 (d, 2H, H-Ph), 7.67 (s, 1H, H-6), 7.95 (d, 2H, H-Ph), 8.85 (s, 1H, H-2). Anal. Calcd. for C₂₇H₂₃Cl₂N₃O₅: C, 60.01; H, 4.29; N, 7.78. Found: C, 60.32; H, 4.08; N, 7.45.

5,7-Dichloro-3-(2-deoxy-β-D-erythro-pentofuranosyl)-3H-imidazo[4,5-b]pyridine (8),

5,7-dichloro-3-(2-deoxy-α-D-erythro-pentofuranosyl)-3H-imidazo[4,5-b]pyridine (9),

5,7-dichloro-1-(2-deoxy-β-D-erythro-pentofuranosyl)-1H-imidazo[4,5-b]pyridine (10).

To the residue obtained from the above reaction was added 50 mL of methanol saturated at 0 °C with ammonia and the mixture was set aside at room temperature for 36 h. The reaction mixture was evaporated and the residue was flash chromatographed over silica gel eluting with CHCl₃-MeOH-NH₃ (96.5:3:0.5) to give 0.94 g (49%) of **8**, 0.19 g (10%) of **9** and 0.04 g (2%) of **10**. (Yields are based on **4**).

(8): mp 51-53 °C; ¹H NMR (Me₂SO-d₆) δ 2.34 (m, 1H, H-2'), 2.71 (m, 1H, H-2''), 3.55 (m, 2H, CH₂-5'), 3.87 (m, 1H, H-4'), 4.41 (m, 1H, H-3'), 6.39 (t, 1H, J = 6.8 Hz, H-1'), 7.66 (s, 1H, H-6), 8.79 (s, 1H, H-2). Anal. Calcd. for C₁₁H₁₁Cl₂N₃O₃: C, 43.44; H, 3.65; N, 13.82. Found: C, 43.12; H, 3.52; N, 14.11.

(9): ¹H NMR (Me₂SO-d₆) δ 2.44 (m, 1H, H-2'), 2.78 (m, 1H, H-2''), 3.48 (m, 2H, CH₂-5'), 4.19 (m, 1H, H-4'), 4.35 (m, 1H, H-3'), 6.46 (dd, 1H, J = 5.5, 2.0 Hz, H-1'), 7.69 (s, 1H, H-6), 8.82 (s, 1H, H-2). Anal. Calcd. for C₁₁H₁₁Cl₂N₃O₃: C, 43.44; H, 3.65; N, 13.82. Found: C, 43.15; H, 3.51; N, 14.04.

(10): ¹H NMR (Me₂SO-d₆) δ 2.46 (m, 1H, H-2'), 2.67 (m, 1H, H-2''), 3.61 (m, 2H, CH₂-5'), 3.91 (m, 1H, H-4'), 4.41 (m, 1H, H-3'), 6.76 (t, 1H,

$J = 6.1$ Hz, H-1'), 7.66 (s, 1H, H-6), 9.01 (s, 1H, H-2). Anal. Calcd. for $C_{11}H_{11}Cl_2N_3O_3$: C, 43.44; H, 3.65; N, 13.82. Found: C, 43.09; H, 3.48; N, 13.99.

3-(2-deoxy- β -D-*erythro*-pentofuranosyl)-3H-imidazo[4,5-b]pyridine (11).

To 50 mg (0.16 mmol) of **8** in 10 mL of ethanol and 0.5 mL of 2 N NaOH was added 50 mg of 10% Pd/C, and the mixture was shaken with hydrogen at 60 psi for 6 h. The catalyst was removed, and the filtrate concentrated to dryness. The residue was chromatographed on a silica gel column eluting with $CHCl_3$ -MeOH (90:10) to give 25 mg (67%) of **11** as a chromatographically pure solid.

(**11**): 1H NMR (Me_2SO-d_6) δ 2.35 (m, 1H, H-2'), 2.78 (m, 1H, H-2''), 3.60 (m, 2H, CH_2 -5'), 4.91 (m, 1H, H-4'), 4.46 (m, 1H, H-3'), 6.54 (t, 1H, $J = 6.8$ Hz, H-1'), 7.34 (dd, 1H, $J_{6,7} = 7.9$, $J_{6,5} = 4.7$ Hz, H-6), 8.14 (d, 1H, $J_{7,6} = 7.8$ Hz, H-7), 8.39 (d, 1H, $J_{5,6} = 4.7$ Hz, H-5), 8.70 (s, 1H, H-2). Anal. Calcd. for $C_{11}H_{13}N_3O_3$: C, 56.16; H, 5.57; N, 17.86. Found: C, 56.38; H, 5.71; N, 17.57.

3-(2-deoxy- α -D-*erythro*-pentofuranosyl)-3H-imidazo[4,5-b]pyridine (12).

The title compound was prepared following the same procedure as **11**, starting from 50 mg (0.16 mmol) of **9** to obtain 22 mg (58%) of **12** as a chromatographically pure solid.

(**12**): 1H NMR (Me_2SO-d_6) δ 2.43 (m, 1H, H-2'), 2.81 (m, 1H, H-2''), 3.50 (m, 2H, CH_2 -5'), 4.17 (m, 1H, H-4'), 4.36 (m, 1H, H-3'), 6.54 (dd, 1H, $J = 7.7$, 3.1 Hz, H-1'), 7.34 (dd, 1H, $J_{6,7} = 8.1$, $J_{6,5} = 4.7$ Hz, H-6), 8.13 (d, 1H, $J_{7,6} = 8.1$ Hz, H-7), 8.40 (d, 1H, $J_{5,6} = 4.7$ Hz, H-5), 8.73 (s, 1H, H-2). Anal. Calcd. for $C_{11}H_{13}N_3O_3$: C, 56.16; H, 5.57; N, 17.86. Found: C, 56.33; H, 5.69; N, 17.54.

Preparation of N⁷-Substituted 5-chloro-3-(2-deoxy- β -D-*erythro*-pentofuranosyl)-3H-imidazo[4,5-b]pyridines (13a-e).

A mixture of 0.45 g (1.5 mmol) of **8** and 30 mL of the appropriate amine was heated in a steel bomb at 135 °C for 20 h. Compound **13e** was prepared by heating a mixture of 0.45 g (1.5 mmol) of **8** in 30

mL of ethanol and 0.61 g (4.5 mmol) of L-amphetamine in a steel bomb at 135 °C for 60 h. The reaction mixture was evaporated and the residue was chromatographed on a silica gel column eluting with CHCl_3 - C_6H_6 -MeOH (70:20:10) to give **13a-e** as chromatographically pure solids

(**13a**) yield: 46%; mp 227-229 °C; ^1H NMR ($\text{Me}_2\text{SO}-d_6$) δ 2.22 (m, 1H, H-2'), 2.66 (m, 1H, H-2''), 3.54 (m, 2H, CH_2 -5'), 3.84 (m, 1H, H-4'), 4.36 (m, 1H, H-3'), 6.28 (t, 1H, $J = 6.9$ Hz, H-1'), 6.36 (s, 1H, H-6), 6.78 (s, 2H, NH_2), 8.29 (s, 1H, H-2). Anal. Calcd. for $\text{C}_{11}\text{H}_{13}\text{ClN}_4\text{O}_3$: C, 46.41; H, 4.60; N, 19.68. Found: C, 46.12; H, 4.51; N, 19.92.

(**13b**) yield: 51%; mp 129-131 °C; ^1H NMR ($\text{Me}_2\text{SO}-d_6$) δ 2.23 (m, 1H, H-2'), 2.64 (m, 1H, H-2''), 2.87 (d, 3H, NCH_3), 3.54 (m, 2H, CH_2 -5'), 3.83 (m, 1H, H-4'), 4.36 (m, 1H, H-3'), 6.28 (m, 2H, H-1' and H-6), 7.24 (d, 1H, NH), 8.29 (s, 1H, H-2). Anal. Calcd. for $\text{C}_{12}\text{H}_{15}\text{ClN}_4\text{O}_3$: C, 48.25; H, 5.06; N, 18.75. Found: C, 47.95; H, 4.89; N, 18.96.

(**13c**) yield: 75%; mp 196-198 °C; ^1H NMR ($\text{Me}_2\text{SO}-d_6$) δ 1.57 (m, 4H, H cyclopent.), 1.69 (m, 2H, H cyclopent.), 1.97 (m, 2H, H cyclopent.), 2.26 (m, 1H, H-2'), 2.65 (m, 1H, H-2''), 3.54 (m, 2H, CH_2 -5'), 3.84 (m, 1H, H-4'), 4.29 (m, 1H, H-1 cyclopent.), 4.38 (m, 1H, H-3'), 6.29 (t, 1H, $J = 6.9$ Hz, H-1'), 6.36 (s, 1H, H-6), 7.05 (d, 1H, NH), 8.29 (s, 1H, H-2). Anal. Calcd. for $\text{C}_{16}\text{H}_{21}\text{ClN}_4\text{O}_3$: C, 54.47; H, 6.0; N, 15.88. Found: C, 54.71; H, 6.13; N, 15.54.

(**13d**) yield: 46%; mp 112-114 °C; ^1H NMR ($\text{Me}_2\text{SO}-d_6$) δ 1.13 and 1.29 (m, 5H, H cyclohexyl), 1.58, 1.70, and 1.87 (m, 5H, H cyclohexyl), 2.23 (m, 1H, H-2'), 2.63 (m, 1H, H-2''), 3.52 (m, 3H, CH_2 -5' and H-1 cyclohexyl), 3.83 (m, 1H, H-4'), 4.37 (m, 1H, H-3'), 6.28 (t, 1H, $J = 6.9$ Hz, H-1'), 6.38 (s, 1H, H-6), 6.90 (d, 1H, NH), 8.29 (s, 1H, H-2). Anal. Calcd. for $\text{C}_{17}\text{H}_{23}\text{ClN}_4\text{O}_3$: C, 55.66; H, 6.32; N, 15.27. Found: C, 55.92; H, 6.18; N, 15.51.

(**13e**) yield: 50%; mp 260-262 °C; ^1H NMR ($\text{Me}_2\text{SO}-d_6$) δ 1.15 (d, 3H, CH_3 -CH), 2.23 (m, 1H, H-2'), 2.68 (m, 2H, H-2'' and $\text{CH}_2\text{C}_6\text{H}_5$), 2.95 (m, 1H, $\text{CH}_2\text{C}_6\text{H}_5$), 3.73 (m, 2H, CH_2 -5'), 3.84 (m, 1H, H-4'), 4.37 (m, 2H, H-3' and CHNH), 6.28 (pt, 1H, $J = 6.9$ Hz, H-1') 6.36 (s, 1H, H-6), 7.04 (d, 1H, NH), 7.21 (m, 5H, H-Ph), 8.32 (s, 1H, H-2). Anal. Calcd. for $\text{C}_{20}\text{H}_{23}\text{ClN}_4\text{O}_3$: C, 59.63; H, 5.75; N, 13.91. Found: C, 59.98; H, 5.89; N, 13.63.

Preparation of N⁷-Substituted 3-(2-deoxy-β-D-erythro-pentofuranosyl)-3H-imidazo[4,5-b]pyridines (14a-e).

To a solution of 13a-e (0.5 mmol) in 40 mL of ethanol and 1 mL of 2N NaOH was added 0.1 g of 10% Pd/C, and the mixture was shaken with hydrogen at 40 psi for 4 h. The catalyst was removed, and the filtrate concentrated to dryness. The residue was chromatographed on a silica gel column eluting with CHCl₃-C₆H₆-MeOH (70:20:10) to give 14a-e as chromatographically pure solids.

(14a) yield: 65%; mp 205-207 °C (lit.¹⁶ 207-208); ¹H NMR (Me₂SO-d₆) δ 2.22 (m, 1H, H-2'b), 2.75 (m, 1H, H-2'a), 3.54 (m, 2H, CH₂-5'), 3.89 (m, 1H, H-4'), 4.41 (m, 1H, H-3'), 6.37 (m, 2H, H-6 and H-1'), 6.47 (s, 2H, NH₂), 7.76 (d, 1H, J= 5.5 Hz, H-5), 8.29 (s, 1H, H-2). Anal. Calcd. for C₁₁H₁₄N₄O₃: C, 52.79; H, 5.64; N, 22.39. Found: C, 52.94; H, 5.45; N, 22.18.

(14b) yield: 80%; mp 162-164 °C; ¹H NMR (Me₂SO-d₆) δ 2.19 (m, 1H, H-2'), 2.75 (m, 1H, H-2''), 2.86 (d, 3H, NCH₃), 3.54 (m, 2H, CH₂-5'), 3.88 (m, 1H, H-4'), 4.38 (m, 1H, H-3'), 6.27 (d, 1H, J= 5.8 Hz, H-6), 6.36 (pt, 1H, J= 6.9 Hz, H-1'), 6.86 (m, 1H, NH), 7.86 (d, 1H, J= 5.8 Hz, H-5), 8.23 (s, 1H, H-2). Anal. Calcd. for C₁₂H₁₆N₄O₃: C, 54.54; H, 6.10; N, 21.20. Found: C, 54.87; H, 6.31; N, 19.94.

(14c) yield: 55%; mp 74-76 °C; ¹H NMR (Me₂SO-d₆) δ 1.53 (m, 4H, H cyclopent.), 1.68 (m, 2H, H cyclopent.), 1.95 (m, 2H, H cyclopent.), 2.18 (m, 1H, H-2'), 2.74 (m, 1H, H-2''), 3.50 (m, 1H, CH₂-5'), 3.62 (m, 1H, CH₂-5''), 3.87 (m, 1H, H-4'), 4.20 (m, 1H, H-1 cyclopent.), 4.38 (m, 1H, H-3'), 6.34 (d, 1H, J= 5.7 Hz, H-6), 6.36 (t, 1H, J= 7.5 Hz, H-1'), 6.61 (d, 1H, J= 7.5 Hz, NH), 7.81 (d, 1H, J= 5.7 Hz, H-5), 8.23 (s, 1H, H-2). Anal. Calcd. for C₁₆H₂₂N₄O₃: C, 60.39; H, 6.97; N, 17.60. Found: C, 60.80; H, 7.11; N, 17.29.

(14d) yield: 61%; mp 67-69 °C; ¹H NMR (Me₂SO-d₆) δ 1.14 and 1.30 (m, 5H, H cyclohexyl), 1.59, 1.70, and 1.90 (m, 5H, H cyclohexyl), 2.16 (m, 1H, H-2'), 2.74 (m, 1H, H-2''), 3.49 (m, 1H, CH₂-5'), 3.60 (m, 1H, CH₂-5''), 3.77 (m, 1H, H-1 cyclohexyl), 3.87 (m, 1H, H-4'), 4.38 (m, 1H, H-3'), 6.35 (m, 2H, H-1' and H-6), 6.48 (d, 1H, NH), 7.78 (d, 1H, J= 6.0 Hz, H-5), 8.23 (s, 1H, H-2). Anal. Calcd. for C₁₇H₂₄N₄O₃: C, 61.43; H, 7.28; N, 16.85. Found: C, 61.87; H, 7.39; N, 16.50.

(14e) yield: 23%; ¹H NMR (Me₂SO-d₆) δ 1.14 (d, 3H, CH₃-CH), 2.16 (m, 1H, H-2'), 2.71 (m, 2H, H-2'' and CH₂C₆H₅), 2.96 (m, 1H, CH₂C₆H₅), 3.54

(m, 2H, CH₂-5'), 3.87 (m, 1H, H-4'), 4.37 (m, 2H, H-3' and CHNH), 6.35 (m, 2H, H-1' and H-6), 6.60 (d, 1H, NH), 7.19 (m, 5H, H-Ph), 7.78 (d, 1H, J = 5.4 Hz, H-5), 8.24 (s, 1H, H-2). Anal. Calcd. for C₂₀H₂₄N₄O₃: C, 65.20; H, 6.57; N, 15.21. Found: C, 64.98; H, 6.76; N, 14.97.

Enzyme Assay.

The method used for the determination of activity against adenosine deaminase has been described in a preceding paper.²

Acknowledgement.

This work was supported by grants from the Italian Research Council (CNR) "Progetto Finalizzato Chimica Fine II" and from Ministero P. I. (fondi ricerca 60%). We thank M. Brandi, F. Lupidi and Dr. G. Rafaiani for technical assistance.

REFERENCES

- 1) Cristalli, G.; Franchetti, P.; Grifantini, M.; Vittori, S.; Bordoni, T.; Geroni, C. *J. Med. Chem.* **1987**, *30*, 1686.
- 2) Lupidi, G.; Cristalli, G.; Marmocchi, F.; Riva, F.; Grifantini, M. *J. Enzyme Inhib.* **1985**, *1*, 67.
- 3) Antonini, I.; Cristalli, G.; Franchetti, P.; Grifantini, M.; Martelli, S.; Petrelli, F. *J. Pharm. Sci.* **1984**, *73*, 366.
- 4) Cristalli, G.; Franchetti, P.; Grifantini, M.; Vittori, S.; Klotz, K.-N.; Lohse, M. *J. Med. Chem.* **1988**, *31*, 1179.
- 5) Cristalli, G.; Grifantini, M.; Vittori, S.; Balduini, W.; Cattabeni, F. *Nucleosides & Nucleotides* **1985**, *4* (5), 625.
- 6) Cristalli, G.; Vittori, S.; Eleuteri, A.; Grifantini, M.; Volpini, R.; Lupidi, G.; Capolongo, L.; Pesenti, E. *J. Med. Chem.* **1991**, *34*, 2226.
- 7) Cristalli, G.; Vittori, S.; Eleuteri, A.; Grifantini, M.; Lupidi, G.; Capolongo, L.; Pesenti, E. *Nucleosides & Nucleotides* **1991**, *10*(1-3), 253.
- 8) a. Vittori, S.; Camaioni, E.; Eleuteri, A.; Volpini, R.; Lupidi, G.; Cristalli, G. 10th International Round Table Nucleosides, Nucleotides and Their Biological Application, Park city-Utah (USA), 16-20 September 1992. b. Cristalli, G.; Vittori, S.;

- Camaioni, E.; Eleuteri, A.; Volpini, R.; Lupidi, G.; Palù, G. manuscript in preparation.
- 9) Hoffer, M. *Chem. Ber.* **1960**, *93*, 2777.
- 10) Mizuno, Y.; Ikehara, M.; Itoh, T.; Saito, K. *J. Org. Chem.* **1963**, *28*, 1837.
- 11) Betbeder, D.; Hutchinson, D.W.; Richards, A.O'L. *Nucleic Acid Res.* **1989**, *17(11)*, 4217-4222.
- 12) Cline, B.L.; Panzica, R.P.; Townsend, L.B. *J. Heterocyc. Chem.* **1978**, *15*, 839.
- 13) Kazimierczuk, Z.; Cottam, H.B.; Revankar, G.R.; Robins, R.K. *J. Am. Chem. Soc.* **1984**, *106*, 6379-6382.
- 14) Hildebrand, C.; Wright, G.E. *J. Org. Chem.* **1992**, *57*, 1808-1813.
- 15) Seela, F.; Rosemeyer, H.; Fischer, S. *Helv. Chim. Acta* **1990**, *73*, 1602-1611.
- 16) Sugawara, T.; Ishikura, T.; Itoh, T.; Mizuno, Y. *Nucleosides & Nucleotides* **1982**, *1(3)*, 239-251.
- 17) Cristalli, G.; Eleuteri, A.; Vittori, S.; Volpini, R.; Camaioni E.; Lupidi, G. *Drug Dev. Res.* **1993**, *28 (3)*, 253.

Received 7/20/93

Accepted 12/7/93