

This article was downloaded by:

On: 26 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713597286>

### Synthesis and Properties of Some 2'-O-d-Ribofuranosyl-nucleosides

Sergey N. Mikhailov<sup>a</sup>; Andre De Bruyn<sup>b</sup>; Piet Herdewijn<sup>c</sup>

<sup>a</sup> Engelhardt Institute of Molecular Biology, Russian Academy of Sciences, Vavilov str., Moscow, Russia; <sup>b</sup> Laboratory of Organic Chemistry, University of Ghent, Ghent, Belgium; <sup>c</sup> Rega Institute, Katholieke Universiteit Leuven, Leuven, Belgium

**To cite this Article** Mikhailov, Sergey N. , De Bruyn, Andre and Herdewijn, Piet(1995) 'Synthesis and Properties of Some 2'-O-d-Ribofuranosyl-nucleosides', *Nucleosides, Nucleotides and Nucleic Acids*, 14: 3, 481 — 484

**To link to this Article:** DOI: 10.1080/15257779508012410

**URL:** <http://dx.doi.org/10.1080/15257779508012410>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

## SYNTHESIS AND PROPERTIES OF SOME 2'-*O*-D-RIBOFURANOSYL-NUCLEOSIDES

Sergey N. Mikhailov<sup>1\*</sup>, André De Bruyn<sup>2</sup>, Piet Herdewijn<sup>3</sup>


<sup>1</sup>Engelhardt Institute of Molecular Biology, Russian Academy of Sciences, Vavilov str. 32, Moscow, 117984, Russia; <sup>2</sup>Laboratory of Organic Chemistry, University of Ghent, Krijgslaan 281, B-9000 Ghent, Belgium; <sup>3</sup>Rega Institute, Katholieke Universiteit Leuven, Minderbroedersstraat 10, B-3000 Leuven, Belgium.

**ABSTRACT :** A high yield preparation of 9-(2-*O*-β-D-ribofuranosyl)-adenine and its pyrimidine analogues has been achieved and their physico-chemical properties were investigated.

Disaccharide nucleosides are important structural elements of several antibiotics such as amicetin, hikizimycin, thuringiensin, tunicamycin and others. Recently some new derivatives of disaccharide nucleosides have been isolated from natural products and shown to possess interesting biological activities<sup>1</sup>. Disaccharide nucleosides have been prepared either by coupling of suitably blocked disaccharides with nucleic acid bases or by condensation of protected nucleosides with monosaccharide. Most of complex nucleoside antibiotics have been prepared using the first route. This route is rather lengthy and it may be shortened in some cases starting from natural nucleosides. Several attempts are described to use classical methods of glycosidic bond formation with blocked nucleosides as a alcohol component. The yields of these reactions were usually very low (20-30%) due to a formation of several by-products<sup>1</sup>. Till now no general high yielding synthetic procedure to disaccharide nucleosides from natural nucleosides has been developed and their physico-chemical and biological properties have not been studied systematically.

Here we present our results on the preparation of *O*-D-ribofuranosyl-nucleosides starting from readily available partially blocked N-acylribonucleosides<sup>2</sup> **1a-c** and a slight excess of 1-*O*-acetyl-2,3,5-tri-*O*-benzoyl-β-D-ribofuranose in the presence of tin tetrachloride. The glycosylation reaction proceeded stereospecifically with the formation of the β-glycosidic bond. The yields of **2a-c** were in the range of 70% to 80%, which is higher than reported for similar condensation reactions between blocked nucleosides and monosaccharides<sup>1</sup>. The reaction conditions are near the same as earlier used for the preparation of alkyl ribofuranosides<sup>3</sup>. The silyl group was selectively deblocked to yield partially protected **3a-c**, which may be used for further modification.

Deblocking with ammonia in methanol gave free disaccharide nucleosides **4a-c** in high overall yield. It should be mentioned that adenine disaccharide **4a** have been isolated from yeast methionine initiator tRNA<sup>4,5</sup> and prepared previously<sup>6</sup> utilizing nearly the same approach.



The chemical shifts of **4a-b** were identified from a COSY 90 (absolute value mode) experiment. For the nucleoside moiety we followed the connectivities starting from H-1'. The glycosidic protons were assigned by comparison of their chemical shifts with reference compounds. Thus the resonance of the proton at the highest frequency is assigned to that of the nucleoside. For the assignment of the resonances of the  $\beta$ -D-ribofuranosyl moiety we could not start by H-1', since the coupling constant  $J_{1',2'}$  is smaller than the resolution. Therefore we started the following up of the connectivities from the resonances of H-5'a and H-5'b. The NMR data including NOE proved the introduction of the ribose moiety into the 2'-position.

According to  $^1\text{H}$  NMR data the incorporation of a ribose moiety in 2'-position of nucleoside resulted in a small shift of the S  $\leftrightarrow$  N equilibrium towards the S-conformer. The chemical shifts and coupling constants of the pyrimidine disaccharide nucleosides **4b,c** have near the same values as in the case of Urd, Cyt (data are not shown) and isopropyl  $\beta$ -D-ribofuranoside (**5**). The

Table :  $^1\text{H}$  NMR Chemical Shifts and coupling constants

compound →	4a		4b		4c		5
moiety →	Ado	Rib	Cyd	Rib	Urd	Rib	Rib
chemical shifts							
H-8 (H-6)	8.32		7.85		7.87		
H-2 (H-5)	8.19		6.12		5.92		
H-1'	6.12	5.07	6.09	5.15	6.04	5.15	5.10
H-2'	4.80	4.13	4.42	4.19	4.44	4.16	3.98
H-3'	4.55	3.99	4.38	4.20	4.36	4.19	4.16
H-4'	4.29	3.82	4.15	4.00	4.11	3.99	3.98
H-5'a	3.92	3.32	3.91	3.73	3.88	3.75	3.79
H-5'b	3.83	2.75	3.82	3.40	3.79	3.48	3.62
coupling constants							
5,6			7.6		8.1		
1',2'	6.4	<0.5	5.2	<0.5	5.0	<0.5	1.5
2',3'	5.0	4.5	5.4	4.6	5.4	4.8	4.8
3',4'	3.3	7.4	4.7	6.8*	5.1	6.9	6.5
4',5'a	2.6	3.7	3.1	3.5	2.9	3.4	3.5
4',5'b	3.6	6.8	4.5	6.7	4.5	6.6	6.6
5'a,5'b	-13.0	-12.0	-12.7	-12.2	-12.7	-12.1	-12.2

$^1\text{H}$  NMR spectra were recorded on a Bruker AM 360 apparatus operating at 360.031 MHz at 20°C with a pulse angle of 19° and a resolution of 0.33 Hz/point for 10 mg samples in 0.5 ml of  $\text{D}_2\text{O}$ .

\* Coupling constant was measured from the pattern for H-4' (ribose moiety).

anisotropical influence of purine ring in **4a** is much more pronounced not only for sugar protons of the nucleoside moiety but also for the ribose residue, especially for 5'a and 5'b protons. Also NOE constraints were observed between base protons and H-5'a and H-5'b of ribose residue. A detailed study on the conformational aspects of these disaccharide nucleosides is in preparation.

Financial support from INTAS (project 93-1500) and NATO (grant HTECH.CRG 940362) is gratefully acknowledged.

## REFERENCES

1. Lerner, L.M. Synthesis and Properties of Various Disaccharide Nucleosides in: *Chemistry of Nucleosides and Nucleotides* (Townsend L.B. ed.), Plenum Press, New York, **1991**, vol. 2, 27-79.
2. Markiewicz, W.T.; Wiewiorowski, M. in *Nucleic Acid Chemistry. Improved and New Synthetic Procedures, and Techniques* (Townsend L.B., Tipson R.S. eds), Wiley J., New York, **1986**, part 3, 229.
3. Hanessian, S.; Banoub, J. in: *Methods in Carbohydrate Chemistry*, Academic Press, New York, **1980**, 243.
4. Desgres, J.; Keith, G.; Kuo, K.C.; Gehrke, C.W. *Nucleic Acids Res.* **1989**, 17, 865.
5. Wimbach, P.; Crain, P.F.; McCloskey, J.A. *Nucleic Acids Res.* **1994**, 22, 2183.
6. Niewczyk, A.; Krzyzaniak, A.; Barciszewski, J.; Markiewicz, W.T. *Nucleosides and Nucleotides*, **1991**, 10, 635.