

This article was downloaded by:

On: 26 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713597286>

Studies on Reactions of Nucleoside H-Phosphonates with Bifunctional Reagents. Part III. Further Studies on Transesterification of Nucleoside H-Phosphonate Diesters with Amino Alcohols

Michał Sobkowski^a; Jacek Stawiński^b; Anna Sobkowska^a; Adam Kraszewski^a

^a Institute of Bioorganic Chemistry, Polish Academy of Sciences, Poznań, Poland ^b Department of Organic Chemistry, Arrhenius Laboratory, Stockholm University, Stockholm, Sweden

To cite this Article Sobkowski, Michał, Stawiński, Jacek, Sobkowska, Anna and Kraszewski, Adam (1995) 'Studies on Reactions of Nucleoside H-Phosphonates with Bifunctional Reagents. Part III. Further Studies on Transesterification of Nucleoside H-Phosphonate Diesters with Amino Alcohols', *Nucleosides, Nucleotides and Nucleic Acids*, 14: 3, 839 – 842

To link to this Article: DOI: 10.1080/15257779508012484

URL: <http://dx.doi.org/10.1080/15257779508012484>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

**STUDIES ON REACTIONS OF NUCLEOSIDE H-PHOSPHONATES WITH
BIFUNCTIONAL REAGENTS. PART 3.
FURTHER STUDIES ON TRANSESTERIFICATION OF NUCLEOSIDE
H-PHOSPHONATE DIESTERS WITH AMINO ALCOHOLS**

**Michał Sobkowski^a, Jacek Stawiński^b, Anna Sobkowska^a
and Adam Kraszewski^{a*}**

^a Institute of Bioorganic Chemistry, Polish Academy of Sciences,
Noskowskiego 12/14, 61-704 Poznań, Poland.


^b Department of Organic Chemistry, Arrhenius Laboratory, Stockholm University,
S-10691 Stockholm, Sweden.

Abstract. Reactions of 5'-O-(4,4'-dimethoxytrityl)thymidin-3'-yl 2-cyanoethyl phosphonate with alcohols, amines and amino alcohols in pyridine are described. Transesterification was found to be faster than β -elimination of 2-cyanoethyl group.

INTRODUCTION

H-Phosphonate diesters with nucleophiles undergo several characteristic reactions and the most typical are: transesterification, alkylation, and dealkylation (for review see ref. 1). However, in mild basic conditions often used in natural products chemistry simple dialkyl H-phosphonates appear to be relatively stable.

During our studies on reactions of nucleoside H-phosphonates with bifunctional reagents² we have found that nucleoside H-phosphonate diesters could react with amines, alcohols, and amino alcohols giving substitution products at the phosphorus center³. It is noteworthy that these reactions proceed under much milder conditions than those required for simple dialkyl H-phosphonate diesters⁴⁻⁶. In this paper reactivity of 5'-O-(4,4'-dimethoxytrityl)thymidin-3'-yl 2-cyanoethyl phosphonate diester **1**, which under the reaction conditions may undergo transesterification or β -elimination, is discussed.




Scheme 1

RESULTS AND DISCUSSION

Reaction with n-butylamine. Compound **1** when treated with *n*-butylamine within 30 min produced only 5'-O-(4,4'-dimethoxytrityl)thymidin-3'-yl phosphonate monoester **5** as a product of β -elimination of the 2-cyanoethyl group.

Reaction with ethanol and a mixture of ethanol and n-butylamine (Scheme 1). The reaction of **1** with ethanol alone was sluggish and after 23 h the reaction mixture consisted of unreacted **1** (32%), 5'-O-(4,4'-dimethoxytrityl)thymidin-3'-yl ethyl phosphonate **2** (42%), 2-cyanoethyl ethyl phosphonate **3**⁸ (14%), diethyl phosphonate **4** (7%), and some by-products (5%). The same reaction proceeded much faster in the presence of butylamine and **1** after 2 min reacted completely. At this stage the main compounds observed were the products of transesterification **2** and **3** which underwent further substitution, and after 4 h diethyl phosphonate **4** was found as a final product. The H-phosphonate monoesters **5** and **6** were formed as products of a parallel β -elimination of the 2-cyanoethyl group.

Reactions with amino alcohols 7a-d (Scheme 2). The reaction of **1** with 2-aminoethanol **7a** was very fast and the starting H-phosphonate was consumed in less than 1 min yielding 59% of di(2-aminoethyl) phosphonate **10a**, 23% of 2-aminoethyl 2-cyanoethyl phosphonate **9a**⁸, and 18% of **5** (β -elimination of 2-cyanoethyl group). After 4 min **9a** was also converted to **10a** which underwent further transformation to some unidentified compounds. In the case of 3-aminopropan-1-ol **7b**, after 1 min the main product was



Scheme 2

Table 1. The ^{31}P NMR data of the intermediates and the final products.

Compound	Chemical shift ^a (ppm)	$^1J_{\text{PH}}$ (Hz)	$^3J_{\text{PH}}$ (Hz)
1	7.72; 7.74	718.4	2.8 ^b
3 ^g	7.94	693.4	9.3 ^c
6	2.57	606.1	8.4 ^e
9a ^g	8.29	698.9	8.3 ^c
9b ^g	8.37	705.8	d
9c ^g	8.15	d	d
9d ^g	8.16	d	d
11d ^g	2.95	595.1	8.4 ^e

^aSpectra in pyridine with heteronuclear decoupling (0.5% H_3PO_4 in D_2O as an external reference).^bDoublets of quartets. ^cDoublet of quintets. ^dNot resolved due to overlapping. ^eDoublet of triplets.^g ^{31}P NMR data for H-phosphonate diesters of type **2**¹, **8**², and **10**³ have been already published.

3-aminopropyl 5'-O-(4,4'-dimethoxytrityl)thymidin-3'-yl phosphonate **8b** (74%) together with 3-aminopropyl 2-cyanoethyl phosphonate **9b**^g (10%), **5** (12%), and bis(3-aminopropyl) phosphonate **10b** (4%), which became the final product of transesterification after 2 h. For 4-aminobutan-1-ol **7c**, the reaction looked very similar to that of **7b** in the first stage, but compounds **8c** and **10c** were rather unstable and were converted into species which were not further identified. The reaction of **1** with 6-amino-hexan-1-ol **7d** seemed to be slower but it was still fast enough so that **1** could not be registered in time required for recording the first NMR spectra. In this instance also the

products of β -elimination were formed [signals at 2.13 ppm (**5**, 22%) and 2.95 ppm (6-aminoethyl phosphonate **11d**⁸, 14%)] but di(6-aminoethyl) phosphonate **10d** was the major product (64%) of transesterification.

From the above experiments it is clear that **1** is much more susceptible to transesterification than 5'-O-(4,4'-dimethoxytrityl)thymidin-3'-yl ethyl phosphonate or even 5'-O-(4,4'-dimethoxytrityl)thymidine-3'-yl 3'-O-(4,4'-dimethoxytrityl)thymidine-5'-yl phosphonate³. The reaction is chemoselective to high extent with 2-cyanoethoxy as a preferable leaving group. The ratios of the products **8** to **9** at the early stages of the reaction were ~7:1 for amino alcohols **7b-d** and ~6:1 for **2** to **3**, when using the mixture of ethanol and n-butylamine. In neither case traces of phosphoramidates could be detected in the reaction mixtures.

These properties of 2-cyanoethyl nucleoside H-phosphonate diesters should be kept in mind while working with this type of compounds (e.g. during their transformation towards phosphoesters in the presence of active nucleophiles). On the other hand this fast reaction (particularly with 2-aminoethanol) can be employed for rapid dephosphonylation of diesters, e.g., for the recovery of nucleosides as described recently by Brill⁹.

ACKNOWLEDGEMENTS

We are indebted to Prof. Per J. Garegg for his interest, and to the State Committee for Scientific Research, Republic of Poland, the Swedish Natural Science Research Council and the Swedish Research Council for Engineering Sciences for financial support.

REFERENCES AND NOTES

- 1 K. Troev, G. Borisov, *Phosphorus and Sulfur*, 1987, **29**, 129.
- 2 A. Kraszewski, M. Sobkowski and J. Stawiński, *J. Chem. Soc. Perkin Trans. I*, 1993, 1699.
- 3 M. Sobkowski, J. Stawiński, A. Sobkowska and A. Kraszewski, *J. Chem. Soc. Perkin Trans. I*, 1994, 1803.
- 4 E. Müller, *Methoden der organischen Chemie (Houben-Weyl)*, George Thieme Verlag, Stuttgart, 1964, vol. XII/2, p.35.
- 5 K. Troev, E. Tashev and G. Borisov, *Phosphorus and Sulfur*, 1982, **12**, 313.
- 6 E. Tashev, S. Shenkov, K. Troev and G. Borisov, *Phosphor. Sulfur Silicon*, 1991, **56**, 225.
- 7 All reactions proceeded in dry pyridine and were monitored by ³¹P NMR. Concentration of **1** was 0.1 M and the nucleophilic compounds were added in 10-fold molar excess.
- 8 Assignment of ³¹P NMR signal to this compound was made according to its predicted magnetic properties.
- 9 W.K.-D. Brill, *Tetrahedron Lett.*, 1994, **35**, 3041.