

This article was downloaded by:

On: 26 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713597286>

Diazepinone Nucleosides as Inhibitors of Cytidine Deaminase.

G. Cristalli^a; R. Volpini^a; S. Vittori^a; E. Camaioni^a; G. Rafaiani^a; S. Potenza^b; A. Vita^a

^a Dipartimento di Scienze Chimiche, Università di Camerino, Camerino, Italy ^b Dipartimento di Biologia Cellulare, Università di Camerino, Camerino, Italy

To cite this Article Cristalli, G. , Volpini, R. , Vittori, S. , Camaioni, E. , Rafaiani, G. , Potenza, S. and Vita, A.(1996) 'Diazepinone Nucleosides as Inhibitors of Cytidine Deaminase.', *Nucleosides, Nucleotides and Nucleic Acids*, 15: 10, 1567 – 1580

To link to this Article: DOI: 10.1080/07328319608002457

URL: <http://dx.doi.org/10.1080/07328319608002457>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

DIAZEPINONE NUCLEOSIDES AS INHIBITORS OF CYTIDINE DEAMINASE.

G. Cristalli*, R. Volpini, S. Vittori, E. Camaioni, G. Rafaiani, S. Potenza, #A. Vita.

*Dipartimento di Scienze Chimiche and #Dipartimento di Biologia Cellulare,
Università di Camerino, 62032 Camerino, Italy.*

Abstract. The synthesis of 2'-deoxy and 2',3'-dideoxy derivatives of 1-β-D-ribofuranosyl-1,3,4,7-tetrahydro-2H-1,3-diazepin-2-one (**2**) was undertaken in order to find new cytidine deaminase (CDA) inhibitors and potential adjuvants in anticancer chemotherapy. Replacement of ribose by a 2-deoxyribose moiety led to compound **9** that appeared slightly more potent than **2** ($K_i = 2.5 \times 10^{-8}$ M). Remarkably, the corresponding α-2'-deoxynucleoside **10** acted as a very potent inhibitor of human placenta CDA, with a $K_i = 7.5 \times 10^{-8}$ M. Attempt to synthesize the 2',3'-dideoxy derivative of **2** led to N-[4,5-dihydroxy-1-(2-oxo-2,3,4,7-tetrahydro[1,3]diazepin-1-yl)-pentyl]-2,2,2-trifluoroacetamide (**13**), which is devoid of CDA inhibitory activity.

In the last years several experiments have revealed that differences exist between normal and cancer cells in terms of enzyme activities involved in pyrimidine and purine metabolisms.¹⁻⁵

Our interests have been focused on adenosine deaminase (ADA) an enzyme that catalyzes deamination of adenosine and 2'-deoxyadenosine to form inosine and 2'-deoxyinosine,⁶⁻¹¹ and on cytidine deaminase (CDA) which catalyzes the deamination of cytidine and 2'-deoxycytidine to uridine and 2'-deoxyuridine, respectively.¹²⁻¹⁵

Inhibition of these enzymes could result in enhanced antiproliferative and antiviral effects of nucleoside analogs which are substrates of ADA or CDA. As expected, inhibitors of ADA and CDA have been shown to prolong the plasma half-life of these drugs.¹⁶ More recently a tumor-selective approach to chemotherapy has been reported, utilizing analogues of deoxycytidine and the CDA inhibitor tetrahydrouridine (**1**, THU, $K_i = 10^{-7}$ M).¹⁷




FIGURE 1


Moreover, Marquez et al. synthesized a series of β -D-ribofuranosyl cyclic urea nucleosides and demonstrated that the compound with the seven-membered aglycon (2) was the more potent, with a $K_i = 2.5 \times 10^{-8}$ M on the human liver enzyme (FIGURE 1).^{18,19} This behaviour is similar to that observed for the inhibition of ADA: in fact, coformycin (3) and 2'-deoxycoformycin (4) are very potent inhibitors of this enzyme with $K_i = 10^{-11}$ M and $K_i = 10^{-12}$ M, respectively.^{20,21} In addition, we previously reported that the substitution of a ribose by a 2'-deoxyribose resulted in an increase in ADA inhibitory activity also in the 1-deazaadenosine series.⁹

On these bases, the synthesis of 2'-deoxy and 2',3'-dideoxy derivatives of compound 2 was undertaken in order to find new adjuvants in anticancer chemotherapy. Moreover, since these compounds are analogues of natural nucleosides, they will be evaluated with regard to their antitumor and antiviral potential.

CHEMISTRY

The synthesis of 2' deoxynucleosides 9 and 10 was carried out according to SCHEMES 1 and 2.

Activation of the base 1,3,4,7-tetrahydro-2H-1,3-diazepin-2-one (5)¹⁹ was performed by reacting 5 with bis(trimethylsilyl)trifluoroacetamide (BSTFA) at room temperature for 2 h. The crude silylated derivative 5a¹⁹ was then added of freshly prepared 1-chloro-2-deoxy-3,5-di-O-*p*-toluoyl- α -D-erythro-pentofura



SCHEME 1


nose (6)²² in dry acetonitrile in the presence of freshly distilled stannic chloride at room temperature for 24 h to give compounds 7 and 8 in 68% total yield. Deblocking of 7 and 8 with methanolic ammonia at room temperature afforded 1-(2-deoxy- β -D-*erythro*-pentofuranosyl)-1,3,4,7-tetrahydro-2H-1,3-diazepin-2-one (9) and its α anomer 10, in 50% and 40% yield, respectively (SCHEME 2).

The anomeric configuration of compounds 7 and 8 was assigned by applying n.O.e. difference spectroscopy. Saturation of H-1' of 7 resulted in n.O.e.s of the H-2'a and H-4' signals (3.0 % and 2.0 %, respectively) while there was none at H-2'b signal, thus establishing a β -D-configuration. Saturation of H-1' of 8 yielded n.O.e.s of the H-2'b signal (1.3 %) while there was none at H-4' signal, thus establishing an α -D-configuration (TABLE 1).

On these basis, compounds 9 and 10 were identified as β and α anomers, respectively.

The condensation between 5a and 1-methoxy-2,3-dideoxy-5-O-(4-methylbenzoyl)-D-*glycero*-pentofuranose (11)²³ in the presence of trimethylsilyl triflate (TMS-TF) in dry methylene chloride gave a derivative which was afterwards identified as N-[4,5-dihydroxy-5-O-*p*-toluoyl-1-(2-oxo-2,3,4,7-tetrahydro[1,3]diazepin-1-yl)-pentyl]-2,2,2-trifluoroacetamide (12). Deprotection of 12 with methanolic ammonia at room temperature for 24 h gave the corresponding N-[4,5-dihydroxy-1-(2-oxo-2,3,4,7-tetrahydro[1,3]diazepin-1-yl)-pentyl]-2,2,2-trifluoroacetamide (13). (SCHEME 3).

The structure of compounds 12 and 13 was elucidated by means of NMR techniques and chemical reactions. In fact the proton NMR spectrum of 12, taken in DMSO- d_6 , showed, in addition to the expected signals for the dideoxy



R = *p*-toluoyl

SCHEME 2

TABLE 1. N.O.E.-data % of compounds **7** and **8** upon irradiation of H-1' (DMSO-d₆, 25 °C, 300 MHz)

Cpd	H-2'a	H-2'b	H-4'
7	3.0	*	2.0
8	*	1.3	*

*: no detectable intensity enhancement (< 0.5%)

nucleoside, a triplet centered at $\delta = 5.08$ ppm and a doublet at $\delta = 9.62$ ppm, both of them exchangeable with D₂O.

Deblocking in the presence of K₂CO₃ in ethanol led to compound **13** whose ¹H-NMR spectrum presented now three exchangeable protons, two of them overlapped at $\delta = 4.95$ ppm and the other again at $\delta = 9.62$ ppm.


To confirm the presence of an additional hydroxyl group, compound **13** was treated with *p*-nitrobenzoylchloride in dry pyridine at room temperature for 3 hours. This reaction yielded two compounds identified as N-[4,5-dihydroxy-4,5-O-di-*p*-nitrobenzoyl-1-(2-oxo-2,3,4,7-tetrahydro[1,3]diazepin-1-yl)-pentyl]-2,2,2-trifluoroacetamide (**14**) and N-[4,5-dihydroxy-4,5-O-di-*p*-nitrobenzoyl-1-(3-*p*-nitrobenzoyl-2-oxo-2,3,4,7-tetrahydro[1,3]diazepin-1-yl)-pentyl]-2,2,2-trifluoroacetamide (**15**).

Treatment of both **14** and **15** with K₂CO₃ in ethanol led again to compound **13** (SCHEME 3). The ¹H-NMR of compound **14** clearly showed the presence of two overlapped *p*-nitrobenzoyl groups and the disappearance of the multiplet at $\delta = 4.95$ ppm, in comparison with the starting material **13**.

Moreover, the multiplet assigned to H_{4'} in the supposed nucleoside presented a strong downfield shift (from $\delta = 3.65$ ppm to $\delta = 5.50$ ppm).

From these data it was possible to conclude that the sugar was cleaved on the C₁-O bond with formation of a 4,5-dihydroxypentyl chain. Accordingly, the two hydroxyl groups explained the presence of:

- i) signals at $\delta = 5.08$ and at $\delta = 4.95$, both of them exchangeable with D₂O, in the ¹H-NMR spectra of compounds **12** and **13**, respectively;
- ii) two *p*-nitrobenzoyl groups in compound **14**;



SCHEME 3




FIGURE 2

- iii) the strong downfield shift of the multiplet assigned to $H_{4'}$ proton in the 1H -NMR spectrum of compound **14**, due to benzylation of the hydroxyl group directly bond to $C_{4'}$.

To clarify the structure of the substituent on C_1 , IR and ^{13}C -NMR spectra of compound **13** were investigated.

The IR spectrum of compound **13** showed the presence of two carbonyl groups with ν_{\max} 1603 and 1735, respectively, and one of them was assigned to the ureic CO of the diazepinone. These carbonyl carbons were identified also by ^{13}C -NMR spectrum of the same compound: a singlet at $\delta = 164.3$ ppm was again assigned to the ureic CO whereas a quartet at $\delta = 156.17$ ppm with $J = 36.0$ Hz was afterwards assigned to CO of the trifluoroacetamidic group.

Since prior to the coupling reaction, 1,3,4,7-tetrahydro-2H-1,3-diazepin-2-one (**5**) was activated using bis(trimethylsilyl)trifluoroacetamide (BSTFA) as silylating agent to give **5a** (SCHEME 1),¹⁹ it was postulated that the C_1 carbon underwent nucleophilic attack both from the silylated base **5a** and from a trifluoroacetamido group arising from BSTFA (FIGURE 2).

^{19}F -NMR spectrum of compound **13** confirmed the presence of a fluorine signal at $\delta = 279.63$ ppm.

Moreover a trifluoroacetamido group on C_1 in compound **13** could explain the presence of:

- i) an exchangeable doublet at $\delta = 9.57$ ppm corresponding to the amidic proton in the 1H -NMR spectrum;

TABLE 2. CDA inhibitory activity of tetrahydrodiazepinone derivatives.

Compound	Ki (μM), CDA (human placenta)
1	1.66
2	0.075
9	0.025
10	0.075
13	inactive

Cytidine deaminase activity was measured by the decrease in absorbance at 282 nm with cytidine as substrate (See experimental section).

- ii) a signal at $\delta = 156.17$ (quartet, $J = 36.0$ Hz, long range coupling with the three fluorine atoms) corresponding to the carbon of the amidic carbonyl group in the ^{13}C -NMR spectrum;
- iii) a signal at $\delta = 116.19$ ppm (quartet, $J = 288.7$ Hz) corresponding to the carbon of the trifluoromethyl group in the ^{13}C -NMR spectrum.

BIOLOGICAL EVALUATION

The synthesized nucleoside analogs were evaluated as inhibitors of cytidine deaminase purified from human placenta, following the experimental procedure described earlier,¹⁵ and the results are reported in TABLE 2.

As already reported by Marquez *et al.* in studies on mouse kidney and human liver CDA,¹⁹ compound **2** is a potent inhibitor of the human placenta enzyme ($K_i = 7.5 \times 10^{-8}$ M).

Substitution of ribose by a 2-deoxyribose moiety led to compound **9** that appeared slightly more potent than **2**. This finding confirms our rationale based on the results obtained in the 1-deazaadenosine series⁹ and on the higher activity of 2'-deoxycoformycin (**4**) in comparison with that of coformycin (**3**).^{20,21}

Surprisingly, the corresponding α -2'-deoxynucleoside **10** acted as a very potent inhibitor of human placenta CDA, with a K_i similar to that of the β -ribofuranosyl derivative **2**. This finding is not in agreement with previous hypotheses that the α -anomers should not present any relevant inhibitory activity on cytidine deaminase.¹⁸ Moreover, since the reaction between **5a** and tetraacetyl ribose in our hand gave only the β -isomer **2**, as already described by

Marquez *et al.*,¹⁹ we could not test the α -ribofuranosyl derivative of **2**, in order to confirm the result obtained with the α -2'-deoxynucleoside derivative **10**.

The opening of the sugar in the condensation of **5a** with the 2,3-dideoxyribose **11** led to the N-[4,5-dihydroxy-1-(2-oxo-2,3,4,7-tetrahydro [1,3]diazepin-1-yl)-pentyl]-2,2,2-trifluoroacetamide (**13**), that was inactive as CDA inhibitor, in agreement with the very low activity showed by some previously reported N-alkyl and N-aryl derivatives of cytosine and uracil.^{15,25,26}

On the basis that coformycin (**3**) and deoxycoformycin (**4**), which are very potent inhibitors of adenosine deaminase (ADA), present a seven-membered ring in their chemical structure, the tetrahydrodiazepinone derivatives **2**, **9**, **10**, and **13** were also evaluated as inhibitors of ADA from calf intestine and found inactive.

EXPERIMENTAL SECTION

Chemistry.

Melting points were determined with a Buchi apparatus and are uncorrected. ¹H NMR spectra were obtained with a Varian VXR 300 MHz spectrometer. IR spectra were recorded on a Perkin-Elmer Model 297 spectrophotometer. TLC were carried out on pre-coated TLC plates with silica gel 60 F-254 (Merck). For column chromatography, silica gel 60 (Merck) was used. Elemental analyses were determined on a Carlo Erba model 1106 analyser.

1,3-Di(trimethylsilyl)-1,3,4,7-tetrahydro-2H-1,3-diazepin-2-one (**5a**).

To 0.3 g (2.7 mmol) of **5** in 30 ml of dry acetonitrile were added 2.7 ml of bis(trimethylsilyl)trifluoroacetamide (BSTFA) and the mixture was stirred at room temperature for 2 hours. After removing the solvent under *vacuum*, compound **5a** was obtained as a light yellow oil which was dissolved in 5 ml of dry methylene chloride and used for the coupling reactions without further purification.

1-(2-Deoxy-3,5-di-O-*p*-toluoyl- β -D-*erythro*-pentofuranosyl)-1,3,4,7-tetrahydro-2H-1,3-diazepin-2-one (**7**)

1-(2-Deoxy-3,5-di-O-*p*-toluoyl- α -D-*erythro*-pentofuranosyl)-1,3,4,7-tetrahydro-2H-1,3-diazepin-2-one (**8**).

To a ice-cooled solution of **5a**¹⁹ in 5 ml of dry methylene chloride were added 1.14 g (2.93 mmol) of freshly prepared 1-chloro-2-deoxy-3,5-di-O-*p*-toluoyl-D-

erythro-pentofuranose (6)²² and 0.5 ml of SnCl₄. The suspension was stirred at room temperature for 36 hours, then the solvent was removed, the residue neutralized with a saturated sodium bicarbonate solution and extracted with chloroform. The organic layer was washed with water, dried over Na₂SO₄, and evaporated under reduced pressure. The residue was chromatographed on silica gel column eluting with CHCl₃-MeOH (99-1) to give 2.9 g (68%) of a mixture of β and α anomers 7 and 8. Separation of anomers was achieved by flash chromatography: elution with cC₆H₁₂-CHCl₃ (80:20) gave 0.34 g (27%) of 7 and 0.24 g (19%) of 8 as vretuous chromatographically pure solids.

7: ¹H NMR (Me₂SO-*d*₆) δ 2.26 (m, 2H, H-2'a and H-2'b), 2.41 (s, 6H, CH₃), 3.62 (m, 4H, CH₂-4 and CH₂-7), 4.28 (m, 1H, H-4'), 4.52 (m, 2H, CH₂-5',), 5.48 (m, 1H, H-3'), 5.69 (m, 2H, CH=CH), 5.96 (m, 1H, H-1'), 6.27 (m, 1H, NH), 7.37 (d, 4H, J = 8.3 Hz, H-arom), 7.92 (d, 4H, J = 8.3 Hz, H-arom). Anal. Calcd. for C₂₆H₂₈N₂O₆: C, 67.23; H, 6.08; N, 6.03. Found: C, 67.52; H, 6.21; N, 5.89.

8: ¹H NMR (Me₂SO-*d*₆) δ 2.04 (m, 1H, H-2'b), 2.40 (s, 6H, CH₃), 2.79 (m, 1H, H-2'a), 3.58 (m, 2H, CH₂-4), 3.82 (m, 2H, CH₂-7), 4.41 (m, 2H, CH₂-5',), 4.60 (m, 1H, H-4'), 5.47 (m, 1H, H-3'), 5.78 (m, 2H, CH=CH), 5.97 (m, 1H, H-1'), 6.25 (m, 1H, NH), 7.37 (m, 4H, H-arom), 7.90 (d, 4H, J = 7.3 Hz, H-arom). Anal. C₂₆H₂₈N₂O₆: C, 67.23; H, 6.08; N, 6.03. Found: C, 67.49; H, 6.22; N, 5.87.

1-(2-Deoxy-β-D-*erythro*-pentofuranosyl)-1,3,4,7-tetrahydro-2H-1,3-diazepin-2-one (9).

To 0.3 g (0.65 mmol) of 7 was added 80 mL of methanol saturated at 0 °C with ammonia and the mixture was set aside at room temperature for 24 h. The reaction mixture was evaporated and the residue was flash chromatographed over silica gel eluting with CHCl₃-MeOH (from 97:3 to 95:5) to give 0.074 g (50%) of 9 as a chromatographically pure solid.

¹H NMR (Me₂SO-*d*₆) δ 1.73 (m, 1H, H-2'b), 1.89 (m, 1H, H-2'a), 3.45 (m, 2H, CH₂-5',), 3.57 (m, 3H, CH₂-4 and H-4'), 3.64 (m, 2H, CH₂-7), 4.08 (m, 1H, H-3'), 5.79 (m, 3H, H-1' and CH=CH), 6.05 (m, 1H, NH). ¹³C NMR (Me₂SO-*d*₆) δ 37.16 (C-2'), 40.61 (C-4 or C-7), 42.26 (C-7 or C-4), 62.26 (C-5'), 70.89 (C-3'), 85.83 (C-1' and C-4'), 126.92 (C-5 or C-6), 127.84 (C-6 or C-5), 164.63 (C-2). Anal. C₁₀H₁₆N₂O₄: C, 52.62; H, 7.07; N, 12.27. Found: C, 52.88; H, 7.21; N, 12.02.

1-(2-Deoxy-α-D-*erythro*-pentofuranosyl)-1,3,4,7-tetrahydro-2H-1,3-diazepin-2-one (10).

To 0.2 g (0.43 mmol) of 8 was added 50 mL of methanol saturated at 0 °C with ammonia and the mixture was set aside at room temperature for 24 h. The

reaction mixture was evaporated and the residue was chromatographed over silica gel preparative TLC eluting with CHCl_3 -MeOH (90:10) to give 0.050 g (40%) of **10** as a chromatographically pure solid.

^1H NMR ($\text{Me}_2\text{SO}-d_6$) δ 1.63 (m, 1H, H-2'b), 2.32 (m, 1H, H-2'a), 3.34 (m, 2H, CH_2 -5'), 3.57 (m, 2H, CH_2 -4), 3.79 (m, 3H, CH_2 -7 and H-4'), 4.08 (m, 1H, H-3'), 5.76 (m, 3H, H-1' and $\text{CH}=\text{CH}$), 6.02 (m, 1H, NH). Anal. $\text{C}_{10}\text{H}_{16}\text{N}_2\text{O}_4$: C, 52.62; H, 7.07; N, 12.27. Found: C, 52.82; H, 7.25; N, 12.01.

N-[4,5-dihydroxy-5-O-*p*-toluoyl-1-(2-oxo-2,3,4,7-tetrahydro[1,3]diazepin-1-yl)-pentyl]-2,2,2-trifluoroacetamide (12).

To a solution of **5a** in 5 ml of dry methylene chloride was added 0.74 g (2.96 mmol) of 1-methoxy-2,3-dideoxy-5-O-*p*-toluoyl-D-*glycero*-pentofuranose (**11**)²³ and 0.5 ml of TMS-triflate and the mixture was allowed to stand at room temperature for 24 hours. The solvent was evaporated to dryness and the residue neutralized with a saturated sodium bicarbonate solution and extracted with chloroform. The organic layer was washed with water, dried over Na_2SO_4 , and evaporated under reduced pressure. The residue was flash chromatographed on silica gel column eluting with CHCl_3 -MeOH (from 99.75:0.25 to 98:2) to give 0.54 g (45%) of **12** as a chromatographically pure solid.

12: ^1H NMR ($\text{Me}_2\text{SO}-d_6$) δ 1.47 (m, 2H, CH_2 -3), 1.86 (m, 2H, CH_2 -2), 2.40 (s, 3H, CH_3), 3.52 (m, 2H, CH_2 -4'), 3.65 (m, 1H, H-4), 3.79 (m, 2H, CH_2 -7'), 4.14 (m, 2H, CH_2 -5), 5.08 (t, 1H, OH-4), 5.69 (m, 3H, $\text{CH}=\text{CH}$ and H-1), 5.94 (m, 1H, NH-3'), 7.35 (d, 2H, $J = 8.1$ Hz, H-arom), 7.91 (d, 2H, $J = 8.1$ Hz, H-arom), 9.62 (d, 1H, $J = 8.1$ Hz, CHNH). Anal. $\text{C}_{20}\text{H}_{24}\text{F}_3\text{N}_3\text{O}_5$: C, 54.17; H, 5.46; N, 9.48. Found: C, 54.41; H, 5.67; N, 9.12.

N-[4,5-dihydroxy-1-(2-oxo-2,3,4,7-tetrahydro[1,3]diazepin-1-yl)-pentyl]-2,2,2-trifluoroacetamide (13).

To a solution of 0.5 g (1.13 mmol) of **12** in 10 ml of ethanol was added 0.6 g (4.35 mmol) of a suspension of K_2CO_3 in boiling ethanol and the reaction mixture was stirred at room temperature for 1 hour. The insoluble K_2CO_3 was filtered and the filtrate was evaporated to dryness. The residue was flash chromatographed on silica gel column eluting with CHCl_3 -MeOH (from 98:2 to 95:5) to yield 0.13 g (35%) of **13** as a chromatographically pure solid.

IR ν_{max} 1603 ($\text{C}=\text{O}$), 1735 ($\text{C}=\text{O}$); ^1H NMR ($\text{Me}_2\text{SO}-d_6$) δ 1.34 (m, 2H, CH_2 -3), 1.79 (m, 2H, CH_2 -2), 3.14-3.90 (m, 7H, CH_2 -5, CH_2 -4', CH_2 -7', H-4), 4.95 (m, 2H, OH-4 and OH-5), 5.65 (m, 1H, H-1), 5.73 (m, 2H, $\text{CH}=\text{CH}$), 5.93 (m, 1H, NH-3'),

9.57 (d, 1H, $J = 8.1$ Hz, CHNH); ^{13}C NMR ($\text{Me}_2\text{SO}-d_6$) δ 27.69 and 27.74 (C-3), 29.80 (C-2), 41.01 and 41.12 (C-7' or C-4'), 42.51 and 42.53 (C-4' or C-7'), 63.06 and 63.21 (C-5 or C-4), 66.18 and 66.29 (C-4 or C-5), 71.02 and 71.19 (C-1), 116.19 (q, $J = 288.7$ Hz, CF_3), 126.38 (C-5' or C-6'), 128.32 (C-6' or C-5'), 156.17 (q, $J = 36.0$ Hz, C=O), 164.27 and 164.3 (C-2'); ^{19}F NMR ($\text{Me}_2\text{SO}-d_6$) δ 279.63. Anal. $\text{C}_{12}\text{H}_{18}\text{F}_3\text{N}_3\text{O}_4$: C, 44.31; H, 5.58; N, 12.92. Found: C, 44.65; H, 5.71; N, 12.69.

N-[4,5-dihydroxy-4,5-O-di-*p*-nitrobenzoyl-1-(2-oxo-2,3,4,7-tetrahydro[1,3]diazepin-1-yl)-pentyl]-2,2,2-trifluoroacetamide (14)

N-[4,5-dihydroxy-4,5-O-di-*p*-nitrobenzoyl-1-(3-*p*-nitrobenzoyl-2-oxo-2,3,4,7-tetrahydro[1,3]diazepin-1-yl)-pentyl]-2,2,2-trifluoroacetamide (15).

To an ice-cooled solution of 0.25 g (0.77 mmol) of **13** in 5 ml of dry pyridine was added 0.43 g (2.3 mmol) di *p*-nitrobenzoyl chloride, and the mixture was allowed to stand at room temperature for 2 hours. The solvent was removed by distillation under vacuum, and the residue was purified on a silica gel column. Elution with cC_6H_{12} -AcOEt- C_6H_6 (60:30:10) gave 0.15 g (36%) of **14** and 0.05 g (10%) of **15** as chromatographically pure solids.

14: ^1H NMR ($\text{Me}_2\text{SO}-d_6$) δ 1.87 (m, 4H, CH_2 -3 and CH_2 -2), 3.50 (m, 2H, CH_2 -4'), 3.70 (m, 2H, CH_2 -7'), 4.63 (m, 2H, CH_2 -5), 5.50 (m, 1H, H-4), 5.70 (m, 3H, $\text{CH}=\text{CH}$ and H-1), 6.0 (m, 1H, NH-3'), 8.17 (m, 4H, H-arom), 8.37 (m, 4H, H-arom), 9.63 (dd, 1H, CHNH). Anal. $\text{C}_{26}\text{H}_{24}\text{F}_3\text{N}_5\text{O}_{10}$: C, 50.09; H, 3.88; N, 11.23. Found: C, 50.22; H, 4.06; N, 11.02.

15: ^1H NMR ($\text{Me}_2\text{SO}-d_6$) δ 1.74 (m, 2H, CH_2 -3), 1.90 (m, 2H, CH_2 -2), 3.81 (m, 2H, CH_2 -7'), 4.39 (m, 2H, CH_2 -4'), 4.71 (m, 2H, CH_2 -5), 5.47 (m, 1H, H-4), 5.68 (m, 1H, H-1), 5.80 (m, 2H, $\text{CH}=\text{CH}$), 7.80 (d, 2H, H-arom), 8.10 (d, 2H, H-arom), 8.16 (m, 4H, H-arom), 8.33 (m, 4H, H-arom), 9.96 (m, 1H, CHNH). Anal. $\text{C}_{32}\text{H}_{27}\text{F}_3\text{N}_6\text{O}_{13}$: C, 50.53; H, 3.58; N, 11.5. Found: C, 50.74; H, 3.69; N, 10.87.

Enzyme assays

Cytidine deaminase activity was determined by a direct spectrophotometric assay based on the decrease of absorbance at 282 nm, when the substrates (cytidine and deoxycytidine) are deaminated. The enzyme was obtained from human placenta following a purification methods already reported by Vita *et al.*¹⁵ In the inhibition experiments, cytidine was used as substrate and the reaction was followed at 282 nm using a UVIKON 940 (Kontron Instruments) spectrophotometer. The standard reaction mixture consisted of 0.167 mM

cytidine and 100 mM Tris-HCl, pH 7.5, in a final volume of 1.0 ml. The reaction was initiated by the addition of 0.02-0.04 enzyme units. One enzyme unit is defined as the amount of enzyme which catalyzes the deamination of 1 μ mol of cytidine pre minute at 37 °C.

The method used for the determination of activity against adenosine deaminase has been described by Lupidi *et al.* in a preceding paper.⁷

Acknowledgements

This work was supported by a grant (no 94.00599.CT11) from the Italian Research Council (CNR). We thank M. Brandi, F. Lupidi and G. Sagratella for technical assistance.

REFERENCES

- 1) Slingerland, R.J.; Van Gennip, A.H.; Bodlaender, J.M.; Voute P.A.; Van Kuilenburg, A.B.P. *Eur. J. Biochem.*, **1995**, *233*, 538-543.
- 2) Slingerland, R.J.; Bodlaender, J.M.; Van Lenthe, H.; Van Kuilenburg, A.B.P.; Van Gennip, A.H. *Clin. Chem. Enzym. Commun.* **1993**, *5*, 315-319.
- 3) Marijnen, Y.M.T.; De Korte, D.; Haverkort, W.A.; Den Breejen, E.J.S.; Van Gennip, A.H. *Biochim. Biophys. Acta* **1989**, *1012*, 148-155.
- 4) Van den Berg, A.A.; Van Lenthe, H.; Busch, S.; De Korte, D.; Roos, D.; Van Kuilenburg, A.B.P.; Van Gennip, A.H. *Eur. J. Biochem.* **1993**, *216*, 161-167.
- 5) Weber, G. *Eur. J. Biochem.* **1983**, *43*, 3466-3492.
- 6) Lupidi, G.; Riva, F.; Cristalli, G.; Grifantini, M. *It. J. Biochem.* **1982**, *31*, 398-403.
- 7) Lupidi, G.; Cristalli, G.; Marmocchi, F.; Riva F. *J. Enzyme Inhib.* **1985**, *1*, 67-75.
- 8) Lupidi, G.; Falasca, M.; Marmocchi, F.; Venardi, G.; Cristalli, G.; Riva, F. *Biochemistry Int.* **1992**, *26*, 1053-1063.
- 9) Cristalli, G.; Eleuteri, A.; Vittori, S.; Volpini, R.; Camaioni E.; Lupidi, G. *Drug Dev. Res.* **1993**, *28* (3), 253-258.
- 10) Cristalli, G.; Eleuteri, A.; Volpini, R.; Vittori, S.; Camaioni E.; Lupidi, G. *J. Med. Chem.* **1994**, *37*, 201-205.
- 11) Cristalli, G.; Vittori, S.; Eleuteri, A.; Volpini, R.; Camaioni E.; Lupidi, G.; Mahmood, N.; Bevilacqua, F.; Palu, G. *J. Med. Chem.* **1995**, *38*, 4019-4025.
- 12) Vita, A.; Amici, A.; Cacciamani, T.; Lanciotti, M.; Magni, G. *Biochemistry*, **1985**, *24*, 6020-6024.

- 13) Vita, A.; Amici, A.; Cacciamani, T.; Lanciotti, M.; Magni, G. *The International Journal of Biochemistry* **1986**, *18*, 431-436.
- 14) Vita, A.; Cacciamani, T.; Natalini, P.; Ruggieri, S.; Raffaelli, N.; Magni, G. *Comp. Biochem. Physiol.* **1989**, *93b*, 591-594.
- 15) Cacciamani, T.; Vita, A.; Cristalli, G.; Vincenzetti, S.; Natalini, P.; Ruggieri, S.; Amici, A.; Magni, G. *Arch. Biochem. Biophys.* **1991**, *290*, 285-292.
- 16) Marquez, V. E. in *Developments in Cancer Chemotherapy*; Glazer, R. I., Ed.; CRC Press: Florida, **1984**; pp 91-114, and references cited within.
- 17) Boothman, D. A.; Briggler, T. V.; Greer, S. *Pharmac. Ther.* **1989**, *42*, 65-88.
- 18) Marquez, V. E.; Liu, S. P.; Kelley, A. J.; Driscoll, J. S.; Mc Cormack, J. J. *J. Med. Chem.* **1980**, *23*, 715-717.
- 19) Liu, P. S.; Marquez, V. E.; Driscoll, J. S.; Fuller, R. W.; Mc Cormack, J. J. *J. Med. Chem.* **1981**, *24*, 662-666.
- 20) Cha, S.; Agarwal, R. P.; Parks, R. E. *Biochem. Pharmacol.* **1975**, *24*, 2187-2197.
- 21) Wentworth, D. F.; Wolfenden, R. *Biochemistry*, **1975**, *14*, 5099-5105.
- 22) Hoffer, M. *Chem. Ber.* **1960**, *93*, 2777-2781.
- 23) Abdel-Megied, A. E.-S.; Pedersen, E. B.; Nielsen, C. M. *Monat. Chem.* **1991**, *122*, 59-70.
- 24) Kreis, K. A.; Watanabe, K. A.; Fox, J. J. *Helv. Chim. Acta* **1978**, *61*, 1011-1016.
- 25) Baker, B. R.; Kelley, J. L. *J. Med. Chem.* **1968**, *11*, 682-686.
- 26) Baker, B. R.; Kelley, J. L. *J. Med. Chem.* **1968**, *11*, 686-690.

Received January 20, 1996

Accepted June 29, 1996