

This article was downloaded by:

On: 26 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713597286>

A Convenient Synthesis of Deoxynebularine Phosphoramidite

Douglas Picken^a, Vicki Gault^a

^a Link Technologies Limited, Glasgow, U.K.

To cite this Article Picken, Douglas and Gault, Vicki(1997) 'A Convenient Synthesis of Deoxynebularine Phosphoramidite', *Nucleosides, Nucleotides and Nucleic Acids*, 16: 7, 937 – 939

To link to this Article: DOI: 10.1080/07328319708006110

URL: <http://dx.doi.org/10.1080/07328319708006110>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

A CONVENIENT SYNTHESIS OF DEOXYNEBULARINE PHOSPHORAMIDITE

Douglas Picken* and Vicki Gault


Link Technologies Limited, 2 Napier Court, Cumbernauld, Glasgow G68 0LG, U.K.

ABSTRACT

Enzyme catalysis by a cell free preparation from *Lactobacillus leichmannii* is used to accomplish the transformation of thymidine to deoxynеbularine in the presence of purine. The resulting mixture of nucleosides is then converted to their 5'-dimethoxytrityl derivatives which are easily separated using conventional chromatography. Phosphitylation of the 5'-dimethoxytrityl-deoxynеbularine then gives the phosphoramidite derivative suitable for incorporation into oligonucleotides.

INTRODUCTION

Deoxynеbularine (9- β -D-2'-deoxyribofuranosylpurine) is the deoxy analogue of the naturally occurring antibiotic nebularine¹. It is of interest in the systematic study of the structural elements involved in hydrogen bonding² and for its ability to form the N..CG motif in triple helical structures³.




Deoxynеbularine

Previous reports of the synthesis of this compound involved either a multi-step 2' deoxygenation sequence⁴ starting from very expensive natural nebularine or the deamination of protected 2'-deoxyadenosine⁵ in a non-aqueous diazotization reaction involving prolonged treatment at elevated temperatures. This we found to give variable results and led to highly coloured material. Because neither of these alternatives seemed totally satisfactory for the synthesis of multi-gram amounts of

deoxynebularine, we developed a method based on the enzyme system⁶ derived from *Lactobacillus leichmannii*. This enzyme system has previously been shown to be of use in the preparation of a variety of nucleoside analogues such as the 2'-deoxy and 2',3'-dideoxynucleosides of 2-aminopurine⁷. A previously reported enzymic preparation of deoxynebularine suffers from the disadvantage of requiring expensive 2-deoxyribose-1-phosphate as substrate⁸.

RESULTS AND DISCUSSION

Nucleoside N-deoxyribosyltransferase (E.C. 2.4.2.6) catalyses the transfer between purine and pyrimidine bases and deoxynucleosides⁹.



The insensitivity of the enzyme from *Lactobacillus* spp. towards organic solvents allows the use of crude preparations from these sources. The addition of solvents inhibits the competing activities of degradative enzymes while maintaining the activity of interest. We have investigated a variety of water-miscible and immiscible solvents as well as different nucleosides in this reaction (B1=thymidyl, cytidyl, uridyl; B2 = purine). For our preparative purposes, thymidine in the presence of purine dissolved in buffer containing 10% ethanol is transformed into the desired product -deoxynebularine. Using a five-fold excess of nucleoside, the reaction reaches 100% conversion of purine in a period of several days, without other side reactions being detectable by hplc. The resulting crude mixture of nucleosides is then most conveniently converted to their 5'-dimethoxytrityl derivatives by standard methods prior to separation by column chromatography. We have found that this is possible even in the presence of citrate buffer residues. Pure 5'-dimethoxytrityl-deoxynebularine is subsequently phosphitylated to yield the phosphoramidite derivative suitable for incorporation into oligonucleotides by automated synthesisers.

This synthesis has the advantages of being environmentally friendly by minimising the use of organic solvents, using cheap starting material and achieving high conversion of heterocyclic base to product nucleoside.

EXPERIMENTAL

Enzyme preparation. *Lactobacillus leichmannii* (ATCC4797) was grown in 5L MRS broth at 37°C for 24 hours (1% starter inoculum). The cells were harvested by centrifugation at 10000 x g for 15 minutes at 4°C and washed twice by re-suspension and centrifugation in 20mM PIPES buffer (pH 6.5, 0.02% sodium azide). The cell pellet (wet weight approximately 15g) was then frozen and thawed before being passed twice through a French press and made up to a volume of 10ml in the same buffer. After

centrifugation at 25000 x g for 45 minutes at 4°C the supernatant was dialysed overnight against two changes of 4L PIPES buffer and then stored frozen until required for use.

Enzyme incubation and derivatisation. Thymidine (14.5g, 60mmol) and purine (2.4g, 20mmol) were dissolved in citrate buffer (0.1M, pH 6.0, 0.05% sodium azide) containing 10% ethanol (total volume 495ml) and enzyme extract was added (5ml). After incubation at 40°C for 6 days, the reaction mixture was evaporated to dryness, co-evaporated with pyridine (3 x 100ml) and the resulting solid was suspended in pyridine (100ml). Dimethoxytrityl chloride (20.5g, 60mmol) was added in several portions and the reaction allowed to proceed overnight. Following evaporation and aqueous workup, the mixture of nucleosides was separated by chromatography over silica gel, eluting with ethyl acetate. This yielded pure 5'-dimethoxytrityl-deoxynebularine (3.6g, 7 mmol) whose identity was confirmed by ¹H and ¹³C nmr spectroscopy.

Standard methods were then used to convert this into its cyanoethyl phosphoramidite derivative.

REFERENCES

1. Biescle, J.J. *Cancer* **1955**, 8, 87
2. For a recent example of the use of deoxynebularine in the investigation of hydrogen bonding in aptamers see Huizenga, D.E.; Szostak, J.W. *Biochemistry* **1995**, 34, 656-665
3. Stilz, H.U.; Dervan, P.B. *Biochemistry* **1993**, 32, 2177-2185
4. Jiricny, J.; Wood, S.G.; Martin, D.; Ubasawa, A. *Nucleic Acids Res.* **1986**, 14, 6579-90
5. Nair, V.; Chamberlain, S.D. *Synthesis* **1984**, 401-3
6. Huard, C.N.; Hutchinson, D.W. *Biocatalysis* **1992**, 6, 201-210
7. Hicks, N.; Hutchinson, D.W.; Mahmood, N.; Hay, A.J. *Antiviral Chem. Chemotherapy* **1992**, 3, 153-156
8. Caffaz, S.; Allegrini, S.; Cercignani, G. *Ital. J. Biochem.* **1990**, 39, 115-20
9. For a review of the enzymatic synthesis of nucleosides see Hanrahan, J.R.; Hutchinson, D.W. *J. Biotechnology* **1992**, 23, 193-210