


This article was downloaded by:

On: 26 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713597286>

Duplex Stability of Oligonucleotides Containing 7-Substituted 7-Deaza- and 8-Aza-7-Deazapurine Nucleosides

F. Seela^a; N. Ramzaeva^a; M. Zulauf^a

^a Laboratorium für Organische und Bioorganische Chemie, Institut für Chemie, Universität Osnabrück, Osnabrück, Germany

To cite this Article Seela, F. , Ramzaeva, N. and Zulauf, M.(1997) 'Duplex Stability of Oligonucleotides Containing 7-Substituted 7-Deaza- and 8-Aza-7-Deazapurine Nucleosides', *Nucleosides, Nucleotides and Nucleic Acids*, 16: 7, 963 – 966

To link to this Article: DOI: 10.1080/07328319708006116

URL: <http://dx.doi.org/10.1080/07328319708006116>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.


DUPLEX STABILITY OF OLIGONUCLEOTIDES CONTAINING 7-SUBSTITUTED 7-DEAZA- AND 8-AZA-7-DEAZAPURINE NUCLEOSIDES

F. Seela*, N. Ramzaeva, and M. Zulauf


Laboratorium für Organische und Bioorganische Chemie, Institut für Chemie, Universität Osnabrück, Barbarastr. 7, D-49076 Osnabrück, Germany.

ABSTRACT.- The synthesis of 7-substituted 7-deaza- and 8-aza-7-deazapurine 2'-deoxyribonucleosides, their incorporation into oligonucleotides, and the stability of corresponding duplexes is described.

The presence of methyl, as well as halogeno substituents in position 7 of a series of 7-deazapurine-containing oligonucleotides leads to duplex stabilization with retention of the B-DNA structure.^{1,2} In continuation of our investigations on the properties of base-modified oligodeoxynucleotides we now report on the synthesis of the phosphoramidites 1-3 of 7-substituted 8-aza-7-deaza-2'-deoxyadenosine, as well as 7-deazapurine 2'-deoxyribonucleosides. They are employed in solid-phase synthesis of oligonucleotides which are then investigated with regard to duplex stability.



Compounds 4a,b³, 6⁴, and 9⁵ served as precursors for the synthesis of the protected nucleosides 5a,b, 8a-c and 11a,b (Scheme 1), which were then converted into



Scheme 1

phosphoramidites. The ω -(trifluoroacetyl)aminoalkynyl- and hexynyl nucleosides **7a-c**, and **11a,b** were prepared using the palladium-catalyzed cross-coupling reaction⁶. All compounds were characterized by ^1H -, ^{13}C -, and ^{31}P -NMR spectra as well as by elemental analyses.

The oligonucleotides containing 7-bromo or 7-iodo-7-deazapurine 2'-deoxyribonucleosides exhibit significantly higher T_m -values than those containing 2'-deoxyadenosine or -guanosine (Table 1).

TABLE 1. T_m-Values of 7-deazapurine-containing oligonucleotides.

Oligonucleotide	T _m [°C] ^a	Oligonucleotide	T _m [°C] ^b
d(A - T) ₆	33	d(G - C) ₄	60
d(c ⁷ A - T) ₆	36	d(c ⁷ G - C) ₄	53
d(c ⁷ I ⁷ A - T) ₆	60	d(c ⁷ I ⁷ G - C) ₄	70
d(c ⁷ Br ⁷ A - T) ₆	55	d(c ⁷ Br ⁷ G - C) ₄	67
d(c ⁷ Me ⁷ A - T) ₆	41	d(c ⁷ Me ⁷ G - C) ₄	58
d(c ⁷ Hex ⁷ A - T) ₆	50	d(c ⁷ Hex ⁷ G - C) ₄	58
d(c ⁷ X ⁷ A - T) ₆ ^c	50		
d(c ⁷ Z ⁸ Br ⁷ A - T) ₆	52		
d(c ⁷ Z ⁸ I ⁷ A - T) ₆	56		
d(c ⁷ Z ⁸ A - T) ₆	36		

^a 8 μM Oligomer conc., 60 mM Na-cacodylate, pH 7.1, 100 mM MgCl₂, 1 M NaCl; ^b 10 μM oligomer conc., 10 mM Na-cacodylate, pH 7, 10 mM MgCl₂, 100 mM NaCl; ^c X = C≡CCH₂NH₃⁺.

Although, 7-deaza-2'-deoxyguanosine destabilizes the duplex structure [d(c⁷G-C)₄] compared to the parent oligonucleotide containing 2'-deoxyguanosine [d(G-C)₄], the replacement of the 7-hydrogen by a methyl [d(c⁷Me⁷G-C)₄] or a hexynyl group [d(c⁷Hex⁷G-C)₄] increases the stability of the duplex strongly.

On the other hand, the duplex d(c⁷A-T)₆ is slightly more stable than d(A-T)₆. The replacement of the 7-hydrogen by a methyl [d(c⁷Me⁷A-T)₆], a hexynyl [d(c⁷Hex⁷A-T)₆], as well as a 7-(3-aminopropynyl) group [d(c⁷X⁷A-T)₆] leads to further stabilization of the duplexes (Table 1). In the case of 8-aza-7-deaza-2'-deoxyadenosines the introduction of bromo or iodo substituents in position 7 (1a,1b) shows the same tendency of duplex stabilization as 7-deaza-2'-deoxyadenosines.

Table 1 indicates that substituents in the 7-position of 7- deazapurines have considerable steric freedom within the major groove of DNA and may carry bulky substituents without significantly interfering overall binding. Therefore, the 7-position of 7-deazapurines is an ideal position for the functionalization of DNA with reporter groups.

References and Notes

1. F. Seela, H. Thomas, *Helv. Chim. Acta* **1995**, *78*, 94.
2. F. Seela, N. Ramzaeva, Y. Chen, *Bioorg. Med. Chem. Lett.* **1995**, *5*, 3049.
3. F. Seela, M. Zulauf, unpublished results.
4. F. Seela, M. Zulauf, *Synthesis* **1996**, 726.
5. N. Ramzaeva, F. Seela, *Helv. Chim. Acta* **1995**, *78*, 1083.
6. F. W. Hobbs, *J. Org. Chem.* **1989**, *54*, 3420.
7. **5a**: $^1\text{H-NMR}$ ($\text{d}_6\text{-DMSO}$): δ , 2.30 (1H, m, H-2'), 2.84 (1H, m, H-2'), 3.06 (2H, m, 2H-5'), 3.76, 3.78 (6H, 2s, NMe_2), 3.71, 3.72 (6H, 2s, 2 OMe), 3.95 (1H, m, H-4'), 4.51 (1H, m, H-3'), 5.30 (1H, d, 3'-OH), 6.58 (1H, t, H-1'), 6.76-7.37 (13H, m, arom. H), 8.47 (1H, s, H-6), 8.98 (1H, s, CH).

- 8b**: $^1\text{H-NMR}$ ($\text{d}_6\text{-DMSO}$): δ , 1.74 (2H, m, CH_2), 2.23 (1H, m, H-2'), 2.46 - 2.48 (3H, m, CH_2 , H-2'), 3.11, 3.15 (6H, 2s, NMe_2), 3.13 (2H, m, 2H-5'), 3.33 (2H, m, CH_2), 3.74 (6H, s, OCH_3), 3.91 (1H, m, H-4'), 4.34 (1H, m, H-3'), 5.29 (1H, d, 3'-OH), 6.54 (1H, t, H-1'), 6.82-7.36 (13H, m, arom. H), 7.56 (1H, s, H-6), 8.29 (1H, s, H-2), 8.73 (1H, s, CH), 9.43 (1H, s, NH).

- 11b**: $^1\text{H-NMR}$ ($\text{d}_6\text{-DMSO}$): δ , 1.12 (6H, m, 2 Me), 1.74 (2H, m, CH_2), 2.23 (1H, m, H-2'), 2.41 (2H, m, CH_2), 2.76 (1H, m, CH), 3.11 (2H, m, 2H-5'), 3.30 (2H, m, CH_2), 3.72 (6H, s, 2 OCH_3), 3.90 (1H, m, H-4'), 4.32 (1H, m, H-3'), 5.27 (1H, d, 3'-OH), 6.39 (1H, t, H-1'), 6.85-7.37 (14H, m, arom. H, H-6), 9.47 (1H, t, NHCOCF_3), 11.53 (1H, s, NH), 11.81 (1H, s, NH).