

This article was downloaded by:

On: 26 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713597286>

Synthesis and Properties of Oligoribonucleotide Analogues Having Amide (3''-CH₂-CO-NH-5') Internucleoside Linkages

Eriks Rozners^{ab}; Roger Strömberg^{ab}

^a Department of Organic Chemistry, Arrhenius Laboratory, Stockholm University, Stockholm ^b

Laboratory of Organic and Bioorganic Chemistry, Department of Medical Biochemistry and Biophysics, Karolinska Institutet, Stockholm, Sweden

To cite this Article Rozners, Eriks and Strömberg, Roger(1997) 'Synthesis and Properties of Oligoribonucleotide Analogues Having Amide (3''-CH₂-CO-NH-5') Internucleoside Linkages', *Nucleosides, Nucleotides and Nucleic Acids*, 16: 7, 967 — 970

To link to this Article: DOI: 10.1080/07328319708006117

URL: <http://dx.doi.org/10.1080/07328319708006117>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

SYNTHESIS AND PROPERTIES OF OLIGORIBONUCLEOTIDE ANALOGUES
HAVING AMIDE (3'-CH₂-CO-NH-5') INTERNUCLEOSIDE LINKAGES


Eriks Rozners* and Roger Strömberg

Department of Organic Chemistry, Arrhenius Laboratory, Stockholm University,
S-10691 Stockholm and Laboratory of Organic and Bioorganic Chemistry, Department of
Medical Biochemistry and Biophysics, Karolinska Institutet, S-171 77 Stockholm,
Sweden

Abstract: Amide linked ribonucleoside dimer **5** was prepared and converted into selectively protected H-phosphonate building block **9**. Oligoribonucleotide duplexes having amide linkages at selected sites were synthesized and their stability was studied by UV melting experiments.

Oligonucleotides having the phosphodiester replaced with dephosphono linkages have been proposed as potential second generation antisense compounds¹. Most of these studies have been done on oligodeoxynucleotides and their analogues. Because the RNA-RNA duplexes are generally more stable than the DNA-RNA hybrids, oligoribonucleotide analogues having chemically and enzymatically stable dephosphono linkages could be promising as second generation antisense compounds. Recently we found that substitution of phosphodiester with formacetal linkage slightly improved the stability of oligoribonucleotide duplexes². Herein we report preliminary results on synthesis and stability of RNA-RNA duplexes having amide (3'-CH₂-CO-NH-5') linkages at selected positions.

Acid **3** was prepared (Scheme 1) from 2',5'-O-di-TBDMS-3'-O-phenoxythiocarbonyl uridine³ **1** according to procedures successfully used in the deoxy series: radical allylation⁴ gave **2**, cleavage of the double bond⁵ with OsO₄ and NaIO₄ and oxidation⁶ of the aldehyde with NaClO₂ gave **3**. Correct configuration at C3' was confirmed by NOESY spectra of the intermediate aldehyde. Informative NOEs were observed between H1' and 3'-CH₂ and between H3' and H6. Amine **4** was prepared from 5'-azido-5'-deoxyuridine⁷ *via* acylation with acetyl chloride in presence of pyridine and radical reduction with tributylstannane⁸ (61%, two steps). Coupling of **3** and **4** (Scheme 2) in presence of DCC (1 eq.) and 1-hydroxybenzotriazole (1 eq.) gave dimer **5**. Selective



(i) allyltributyltin, AIBN (75%); (ii) OsO₄, NaIO₄ (82%); NaClO₂ (82%).


Scheme 1.

removal of the 5'-O-TBDMS group (limited acetic acid hydrolysis), cleavage of the terminal 2' and 3' acetyl groups and protection of the 5'-OH as monomethoxytrityl ether gave dimer **8**. Synthesis of the dimeric H-phosphonate building block **9** was achieved *via* one pot selective 2'-O-acylation (1.1 eq. of *o*-chlorobenzoyl chloride) followed by 3'-O-phosphonylation².

Correct structures of all compounds were confirmed using ¹H and ¹³C NMR spectroscopy. The signal of H3' in the 3' uridine residue of **9** showed the coupling pattern expected (pentet, 1:2:2:2:1) from spin-spin interactions with H2' and H4' (*J* = ca 5 Hz) and with P (*J* = ca 10 Hz) thus confirming the correct structure of **9** as the 3'-O-hydrogenphosphonate.

Three model oligoribonucleotides (Table) having two, three and six internucleoside amide linkages (**x**) were synthesized using the automated solid phase H-phosphonate method for RNA synthesis⁹ with 2'-O-*o*-chlorobenzoyl protection¹⁰. Dimer **9** was used under standard coupling conditions. Unmodified sequences and the complementary oligoribonucleotides were also synthesized and the stability of the corresponding duplexes were characterized by UV melting experiments (Table). Correct nucleoside residue composition of all oligonucleotides were confirmed by enzymatic degradation (SVPD and AP) followed by RP HPLC analysis.

Substitution of selected phosphodiester linkages with amides (3'-CH₂-CO-NH-5') did not significantly affect the stability of RNA-RNA duplexes. However, the data reveal some sequence dependence which makes it difficult to rationalize the effect of this substitution.



(i) DCC, HOBT (68%); (ii) 80% AcOH, 50°C, 3h; (iii) NH₃/EtOH, 1:1 6h; (iv) 4-Monomethoxytrityl chloride (43%, 3 steps); (v) *o*-Chlorobenzoyl chloride, 1.1 eq, -78°C; (vi) PCl₃, Imidazole, NEt₃, -78°C (69%).

Scheme 2.

Table. Oligonucleotide sequences and affinity towards complementary RNA (t_m)^a.

Entry	Oligonucleotides ^b	unmodified ribo	amide modified ^c
1	AAGCGAUxUUxUGACACU	56.7	56.7 (0)
2	ACAUXUCGUxUGUXUCAGA	53.4	52.5 (-0.3)
3	(UXU) ₆ U	13.8	17.6 (+0.6)

^a Absorbance vs temperature profiles were measured at 260 nm in buffer containing 100 mM NaCl, 10 mM sodium phosphate (pH 7.2), 0.1 mM EDTA and 2 μM oligonucleotides ^b x denotes position of the amide linkage. ^c Δ t_m per modification are given in parenthesis.

RNA-RNA duplexes are generally more stable than the corresponding DNA-RNA complexes. Although there are many different sides to developing antisense compounds, favourable duplex formation is likely to be of major importance. Oligoribonucleotide analogues can from this sense be considered as having even greater potential than their deoxy counterparts. The duplex with the complementary RNA for the full deoxy version of the 16mer (Table, Entry 1) had a $t_m = 46.9^\circ\text{C}$, i.e. 9.8°C lower than for the RNA or amide modified oligomers. Other advantages are ready access to monomeric synthons (e.g. can often be readily prepared from modified sugars and heterocyclic bases) and less costly starting materials. Uniformly modified amide-linked oligoribonucleotides might be interesting candidates for use as second generation antisense compounds.

REFERENCES

1. (a) De Mesmaeker, A.; Altman, K.-H.; Waldner, A.; Wendeborn, S. *Current Opinion Structural Biology* **1995**, 5, 343-355. (b) Sanghvi, Y.S.; Cook, P.D. In *Carbohydrate Modifications in Antisense Research*, Sanghvi, Y.S. and Cook, P.D., Eds., ACS, Washington, DC: **1994**, p. 1-22.
2. Rozners, E.; Strömberg, R. manuscript in preparation.
3. Meier, C.; Huynh-Dinh, T. *Synlett* **1991**, 227-228.
4. De Mesmaeker, A.; Lebreton, J.; Waldner, A.; Fritsch, V.; Wolf, R.M.; Freier, S.M. *Synlett* **1993**, 733-736.
5. Sanghvi, Y.S.; Bharadwaj, R.; Debart, F.; De Mesmaeker, A. *Synthesis* **1994**, 1163-1166.
6. Dalcanale, E.; Montanari, F. *J.Org.Chem.* **1986**, 51, 567-569.
7. (a) Hata, T.; Yamamoto, I.; Sekine, M. *Chem.Lett.* **1975**, 977-980. (b) Yamamoto, I.; Sekine, M.; Hata, T. *J.Chem.Soc.Perkin Trans.I* **1978**, 306-310.
8. Samano, M.C.; Robins, M.J. *Tetrahedron Lett.* **1991**, 32, 6293-6296.
9. (a) Garegg, P.J.; Lindh, I.; Regberg, T.; Stawinski, J.; Strömberg, R.; Henrichson, C. *Tetrahedron Lett.* **1986**, 27, 4055-4059. (b) Westman, E.; Sigurdsson, S.; Stawinski, J.; Strömberg, R. *Nucleic Acids Res. symp. ser. #31* **1994**, 25-26.
10. (a) Rozners, E.; Strömberg, R.; Bizdena, E. *Nucleosides Nucleotides* **1995**, 14, 855-857. (b) Rozners, E.; Renhofa, R.; Petrova, M.; Popelis, J.; Kumpins, V.; Bizdena, E. *Nucleosides Nucleotides* **1992**, 11, 1579-1593.