

This article was downloaded by:

On: 26 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713597286>

Synthesis of Some Fully 3' and 4'-C-Branched-Chain Sugar Nucleoside Analogues

T. Ton-That^a

^a CNRS-UM II 5625, Laboratoire de Chimie Bioorganique, Université Montpellier II, Montpellier Cedex 5, France

To cite this Article Ton-That, T.(1999) 'Synthesis of Some Fully 3' and 4'-C-Branched-Chain Sugar Nucleoside Analogues', *Nucleosides, Nucleotides and Nucleic Acids*, 18: 4, 731 — 732

To link to this Article: DOI: 10.1080/15257779908041556

URL: <http://dx.doi.org/10.1080/15257779908041556>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

SYNTHESIS OF SOME FULLY 3' and 4'-C-BRANCHED-CHAIN SUGAR NUCLEOSIDE ANALOGUES.

T. Ton-That.

CNRS-UM II 5625, Laboratoire de Chimie Bioorganique.
Université Montpellier II, Place Eugène Bataillon.
34095 Montpellier Cedex 5-France.


ABSTRACT: The synthesis of some new 3'-azido-3'-C-substituted pyrimidine nucleoside analogues is described. The key step is the geminal disubstitution of an appropriated ketone carbohydrate, via the regioselective ring opening of the corresponding tosyl-epoxide derivative.

As a part of our studies on the stereospecific synthesis of highly functionalized branched-chain sugars¹, two sets of fully 3' and 4'-C-branched-chain sugar nucleoside analogues - 2',3',4'-trideoxypyranosyl and 2',3'-dideoxyfuranosyl - were prepared and evaluated as antiviral agents. These compounds were characterized by the presence of two functional groups (azido and hydroxymethyl or fluoromethyl) on the same carbon of the sugar moiety.

The synthesis of this novel type of nucleoside analogues was based on the formation, from ulose **1**², of α,β -epoxy-sulphones **2** followed by regioselective opening of their three-membered ring in the presence of azide ions. The resulting α -azido-aldehyde intermediates were transformed into the α -azido hydroxymethyl derivatives **3(a,b)** and subsequently into the α -azido fluoromethyl derivatives **3(c,d)** by exposure their corresponding triflates to fluoride ions . Acetolysis of **3** and condensation³ of the resulting branched-chain acetylated sugars **4** with silylated thymine afforded (35-60%) the first set of 2',3',4'-trideoxypyranosyl nucleoside analogues **4T**.

The access to the furanosyl series was based on the excision of the anomeric carbon by a two step procedure from the common intermediates **4**. The formation of a double bond **5** (90-95%) and its oxidative cleavage by the catalytic RuO₂-NaIO₄ method

resulted in the selective formation of the corresponding lactols (70-75%) which were acetylated to give the desired 2,3-dideoxyfuranosyl derivatives **6**. Condensation of the latter with silylated thymine (90%), deprotection and separation of anomers furnished the dideoxyfuranosylnucleoside analogues **6T** (Scheme 1).



Reagents. (a) $ClCH_2SO_2Ti$, $t\text{-BuOH}$, THF, 85%; (b) (i) NaN_3 , $MeOH:H_2O$, then $NaBH_4$, 85%; (ii) Tf_2O , Py, CH_2Cl_2 , then TBAF, THF, 80%; (c) Ac_2O , $BF_3 \cdot OEt_2$, 90%; (d) Silylated thymine, IK, 18-crown-6, CH_3CN then $NH_3/MeOH$; (e) $p\text{-TsCl}$, Py, DMAP, 90-95%; (f) (i) $NaIO_4$, RuO_2 , K_2CO_3 , CCl_4-H_2O , 70%; (ii) Ac_2O , Py.

Scheme 1

All the compounds synthesized, **4T** and **6T**, were tested for their *in vitro* inhibitory effects on HIV-1 and HBV replication and were found to be inactive.

Acknowledgements

I thank Drs A-M AUBERTIN (Univ. L. Pasteur, Strasbourg) and J.P. SOMADOSSI (Univ. Birmingham, Alabama) for the biological assays.

REFERENCES

1. a) Ton-That, T.; Laborde, M. de los A.; Olesker, A.; Lukacs, G. J. *Chem. Soc. Chem. Commun.* **1988**, 1581. b) Ton-That, T. *J. Carbohydr. Chem.* **1995**, *14*, 995.
2. Pecka, J.; Stanek, J.; Cerny, M. *Collect. Czech. Chem. Commun.* **1974**, *39*, 1192.
3. Azymah, M.; Chavis, C.; Lucas, M.; Imbach, J.L. *Tetrahedron Lett.* **1989**, *30*, 6165.