

This article was downloaded by:

On: 26 January 2011

Access details: Access Details: Free Access

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713597286>

Binding of Substrates by Purine Nucleoside Phosphorylase (PNP) from *Cellulomonas Sp.* - Kinetic and Spectrofluorimetric Studies

B. Wielgus-kutrowska^a; J. Tebbe^b; J. Wierzchowski^c; D. Shugar^a; W. Saenger^b; G. Koellner^b; A. Bzowska^a

^a Department of Biophysics, Institute of Experimental Physics, University of Warsaw, Warsaw, Poland

^b Institute für Kristallographie, Freie Universität Berlin, Berlin, Germany ^c Department of Organic Chemistry, Pedagogical University, Siedlce, Poland

To cite this Article Wielgus-kutrowska, B. , Tebbe, J. , Wierzchowski, J. , Shugar, D. , Saenger, W. , Koellner, G. and Bzowska, A.(1999) 'Binding of Substrates by Purine Nucleoside Phosphorylase (PNP) from *Cellulomonas Sp.* - Kinetic and Spectrofluorimetric Studies', *Nucleosides, Nucleotides and Nucleic Acids*, 18: 4, 871 — 872

To link to this Article: DOI: 10.1080/15257779908041586

URL: <http://dx.doi.org/10.1080/15257779908041586>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

BINDING OF SUBSTRATES BY PURINE NUCLEOSIDE PHOSPHORYLASE (PNP) FROM *CELLULOMONAS SP.* - KINETIC AND SPECTROFLUORIMETRIC STUDIES.

B. Wielgus-Kutrowska¹, J. Tebbe², J. Wierzchowski³, D. Shugar¹, W. Saenger², G. Koellner² and A. Bzowska^{1*}

¹Department of Biophysics, Institute of Experimental Physics, University of Warsaw, Żwirki i Wigury 93, 02 089 Warsaw, Poland; ²Institute für Kristallographie, Freie Universität Berlin, Takustraße 6, D-14195 Berlin, Germany; ³Department of Organic Chemistry, Pedagogical University, 3 Maja 54, 08-110 Siedlce, Poland

ABSTRACT: Dissociation constants and stoichiometry of binding for interaction of *Cellulomonas sp.* purine nucleoside phosphorylase with its substrates: inosine/guanosine, orthophosphate, guanine/hypoxanthine and D-ribose-1-phosphate were studied by kinetic and spectrofluorimetric methods.

Purine nucleoside phosphorylase (PNP, E.C. 2.4.2.1.) from *Cellulomonas sp.* catalyzes the reversible phosphorolysis of inosine (Ino) and guanosine (Guo), and some analogues, as follows: β -purine nucleoside + orthophosphate \rightleftharpoons purine base + α -D-pentose-1-phosphate. Adenosine is not a substrate, but is a competitive inhibitor of Ino phosphorolysis¹. Hence *Cellulomonas* PNP, although trimeric, cannot be assigned to one of the two main classes of PNPs: "high-molecular weight" hexameric bacterial enzymes, with broad specificity towards nucleosides^{2,3}, and "low-molecular weight" trimeric, mainly mammalian, PNPs specific for both binding and catalysis of 6-oxopurine nucleosides^{4,5}.

Commercial *Cellulomonas* PNP (Toyobo, Japan) was purified as previously described¹. Michaelis constants (K_m) and maximal velocities (V_{max}) were determined for both substrates, i.e. for Ino/Guo and orthophosphate (P_i). K_m is 43 and 11 μ M for Ino and Guo, respectively, and the Hill coefficient is 1. Kinetic plots for P_i were not linear, suggesting negative cooperativity or two classes of binding sites (as observed for both




Figure 1: **A** – Fluorescence spectra ($\lambda_{exc} = 290$ nm) of PNP from *Cellulomonas* sp. in 50 mM Hepes buffer pH=7(—), and (---) with 8 μ M Gua (saturating concentration); **B** – Fluorimetric titration ($\lambda_{obs} = 340$ nm) of *Cellulomonas* sp. (trimer E_0 , 1.05 μ M) with Gua (2.3 – 6.2 μ M) expressed as: $[L_0]/fr = K_d/(1-f_r) + N[E_0]$, where L_0 is total concentration of Gua; $f_r = (F_0 - F)/(F_0 - F_\infty)$; F , F_0 and F_∞ are the fluorescence intensities in the presence, absence and saturating concentration of Gua; K_d , dissociation constant; N , number of binding sites per PNP trimer.

"low-" and "high-" molecular weight PNPs). Formation of binary complexes of *Cellulomonas* PNP with guanine and hypoxanthine (Gua, Hx), as well as P_i and D-ribose-1-phosphate (rib-1-P), is accompanied by a marked increase of the intrinsic fluorescence of the protein, most pronounced for Gua (Fig. 1A). Dissociation constants and stoichiometry of binding were determined by fluorimetric titration of PNP in the presence of Hx, Gua, P_i and 1-rib-P. Gua and Hx bind with $K_d = 0.40 \pm 0.05$ μ M and 2.0 ± 0.5 μ M, and $N = 1.07 \pm 0.05$ and 0.9 ± 0.23 , respectively. Hence three molecules are bound per enzyme trimer (Fig. 1B). Assuming there is one binding site per monomer, $F(L_0) = F_0 - L_0(F_0 - F_\infty)/(K_d + L_0)$, titration with rib-1-P gives $K_d = 60 \pm 4$ μ M. For P_i , in line with kinetic results suggesting negative cooperativity, a model with two binding constants, or two classes of binding sites, $F(L_0) = F_0 - (F_0 - F_\infty)[N_1 L_0/(K_{d1} + L_0) + N_2 L_0/(K_{d2} + L_0)]$, gave the best fit. Low- and high- affinity binding sites have K_d values in the mM and μ M range, respectively. (Supported by Polish Committee for Scientific Research (KBN grant 6 P04A 062 09) and, in part, by the Howard Hughes Medical Institute (HHMI 75195-543401).

REFERENCES

1. Wielgus-Kutrowska, B.; Tebbe, J.; Schröder, W.; Luič, M.; Shugar, D.; Saenger, W.; Koellner, G.; Bzowska, A., *Adv. Exp. Med. Biol.* **1998**, *431*, 259-264.
2. Jensen, K.F.; Nygaard, P. *Eur. J. Biochem.* **1975**, *51*, 253-265.
3. Doskočil, J.; Holý, A. *Coll. Czechoslov. Chem. Commun.* **1977**, *42*, 370-383.
4. Stoeckler, J.D. *Developments in Cancer Chemotherapy*, Glazer, R.J. ed., **1984**, 35-60, CRC Press Inc., Boca Raton FL.
5. Montgomery, J.A. *Med. Res. Rev.* **1993**, *13*, 209-228, John Wiley and Sons.