

This article was downloaded by:

On: 26 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713597286>

Stability of Phosphorothioate Oligonucleotides in Aqueous Ammonia in Presence of Stainless Steel

Brett J. Turney^a; Zacharia S. Cheruvallath^a; Mark Andrade^a; Douglas L. Cole^a; Vasulinga T. Ravikumar^a

^a Isis Pharmaceuticals, Carlsbad, CA

To cite this Article Turney, Brett J. , Cheruvallath, Zacharia S. , Andrade, Mark , Cole, Douglas L. and Ravikumar, Vasulinga T.(1999) 'Stability of Phosphorothioate Oligonucleotides in Aqueous Ammonia in Presence of Stainless Steel', *Nucleosides, Nucleotides and Nucleic Acids*, 18: 6, 1187 – 1189

To link to this Article: DOI: 10.1080/07328319908044658

URL: <http://dx.doi.org/10.1080/07328319908044658>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

**STABILITY OF PHOSPHOROTHIOATE OLIGONUCLEOTIDES
IN AQUEOUS AMMONIA IN PRESENCE OF STAINLESS STEEL**

Brett J. Turney, Zacharia S. Cheruvallath, Mark Andrade,
Douglas L. Cole & Vasulinga T. Ravikumar*

Isis Pharmaceuticals, 2290 Faraday Avenue, Carlsbad, CA 92008


Antisense oligonucleotides as modulators of gene expression represent an exciting new drug technology. Oligodeoxyribonucleotide phosphorothioates are now among the most intensively investigated nuclease-resistant antisense analogs, as evidenced by a number of ongoing clinical trials against several disease targets. Structurally, these differ from natural oligonucleotides by the replacement of one of two nonbridging oxygen atoms by a sulfur atom at each internucleotide linkage. Among factors in the successful development of these complex molecules to support broad clinical trials have been advances made in automation, analysis and purification. The large scale synthesis of oligonucleotide phosphorothioates is presently carried out by initial formation of the internucleotide phosphite linkage using solid-phase phosphoramidite chemistry followed by sulfurization. Efficient synthesis of 20-mer oligophosphorothioates has been achieved on 0.15 mole scale with only 1.75-fold excess of amidite synthons. However, as the scale of synthesis increases to meet future market demands, issues related to fast and efficient synthesis, automation, scalability and product purification are also being investigated. One issue has been the protocol for final product deprotection. Since deprotection involves large quantities of saturated aqueous ammonium hydroxide, one might consider use of stainless steel reactors to withstand resulting vapor pressure at 55°C. A recent report,¹ however, discusses the instability of dimer phosphorothioates in aqueous ammonia in the

Table 1: Stability of oligophosphorothioates in ammonia in presence of stainless steel.

Contact Time (h)	% P=S based on SAX	% Full length based on CGE
0	87.85	94.17
2	79.04	93.64
4	76.28	93.42
6	73.75	89.79
10	68.68	88.50
34	65.10	86.60

presence of metal ions. As this is potentially an important issue for phosphorothioate oligonucleotide synthesis, we describe herein our findings regarding deprotection of a 20-mer oligodeoxyribonucleotide phosphorothioate with aqueous ammonia during process development studies.

In a typical experiment, 10 mg of a purified 20-mer oligophosphorothioate bulk drug substance (ISIS 5132/CGP 69846A) [ds(TCCCGCCTGTGACATGCATT)] were dissolved in aqueous ammonia (33%, 10 ml) and divided into six equal batches. Samples were taken up in bottles containing stainless steel shavings (type 316L), sealed, and held in an oven at 55°C. Samples were removed at time intervals of 2, 4, 6, 10, and 34 h, cooled, concentrated and analyzed by anion exchange chromatography and capillary gel electrophoresis. The amount of partially desulfurized material present was found to increase with contact time (Table 1).



In another recent literature report concerning deprotection of phosphorodithioates, low levels of base-catalyzed desulfurization were again reported² under standard

deprotecting conditions in a glass reactor, but desulfurization extent appeared to vary depending on the ammonia lot used. In view of this observation, we repeated the above experiment using ammonia from five different lots. We observed consistent levels ($\pm 5\%$) of desulfurization and chain cleavage across lots (data not shown).

Finally, in order to find whether stainless steel containers could be used for base-catalyzed deprotection, the following experiment was performed: The 20-mer oligonucleotide phosphorothioate was synthesized on a very large scale (80 mmole; loading 85 $\mu\text{mole/g}$) using polystyrene primer support on Pharmacia OligoProcess synthesizer. At the end of the synthesis, two 10 grams lots of the support were taken separately and incubated for 12 h with 200 ml of 30% aqueous ammonium hydroxide in a 2-litre glass container and an electropolished stainless steel (316L) container respectively. The solutions were filtered, solid support washed with ethanol and the combined mixture concentrated and analyzed by ^{31}P NMR and anion exchange HPLC. No significant difference was observed between the two experiments.

In summary, from the above results, it appears that surface integrity of the stainless steel reactor may be more important to get consistently good results than lot-to-lot ammonia variations.

REFERENCES:

1. Kodra, J. T.; Kehler, J.; Dahl, O. *Nucleic Acids Res.* **1995**, *23*, 3349.
2. Wiesler, W. T.; Caruthers, M. H. *J. Org. Chem.*, **1996**, *61*, 4272.