

This article was downloaded by:

On: 26 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713597286>

## Use of Phenylacetyl Disulfide (PADS) in the Synthesis of Oligodeoxyribonucleotide Phosphorothioates

Zacharia S. Cheruvallath<sup>a</sup>; Patrick D. Wheeler<sup>a</sup>; Douglas L. Cole<sup>a</sup>; Vasulinga T. Ravikumar<sup>a</sup>

<sup>a</sup> Isis Pharmaceuticals, Carlsbad, CA

**To cite this Article** Cheruvallath, Zacharia S. , Wheeler, Patrick D. , Cole, Douglas L. and Ravikumar, Vasulinga T.(1999) 'Use of Phenylacetyl Disulfide (PADS) in the Synthesis of Oligodeoxyribonucleotide Phosphorothioates', *Nucleosides, Nucleotides and Nucleic Acids*, 18: 6, 1195 – 1197

**To link to this Article:** DOI: 10.1080/07328319908044660

**URL:** <http://dx.doi.org/10.1080/07328319908044660>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

## **USE OF PHENYLACETYL DISULFIDE (PADS) IN THE SYNTHESIS OF OLIGODEOXYRIBONUCLEOTIDE PHOSPHOROTHIOATES**

**Zacharia S. Cheruvallath, Patrick D. Wheeler, Douglas L. Cole & Vasulinga T. Ravikumar\***

*Isis Pharmaceuticals, 2292 Faraday Avenue, Carlsbad, CA 92008*

Oligodeoxyribonucleotide phosphorothioates, where one of the nonbridging oxygen of the internucleotide phosphate is formally replaced by a sulfur atom, are the first class to undergo human clinical trials. Ongoing phosphorothioate clinical trials against several disease targets has necessitated manufacture of very large quantities of oligonucleotide active pharmaceutical ingredient (API). Clinical trial and future market demands have stimulated effort towards developing cost efficient large scale synthesis of these complex bio-molecules. This effort has culminated in the routine synthesis of 20-mer oligodeoxyribonucleotide phosphorothioates at 150 mmole scale using only 1.75-fold molar excess of amidites in less than 8 h total synthesis time.

The current chemistry of choice, the phosphoramidite route, requires sulfurization after each coupling. Hence, it is crucial that the sulfur transfer step be highly efficient. During the last few years, a variety of sulfurizing reagents have been investigated. Among these, 3H-1,2-benzodithiol-3-one 1,1-dioxide (Beaucage reagent) is most widely used because of its rapid sulfurization and commercial availability. However, the sulfoxide by-product formed during sulfurization is a potent oxidizing reagent and may be responsible for inconsistent reagent performance lot to lot. In addition, the synthesis of this reagent involves multiple steps and handling of difficult materials that makes it not amenable to large scale productions. Thus, there is an urgent need to replace this expensive reagent.

Phenylacetyl disulfide (PADS) has been reported in the literature as a sulfur transfer reagent in the synthesis of deoxyribonucleotide phosphorothioates,<sup>1,2</sup> but under the reported conditions it was found to be inefficient. While investigating a range of potential sulfur transfer reagents, we noticed that rate and efficiency of sulfurization are very dependent on the solvent system. Thus, bis(diisopropoxy phosphinothioyl) disulfide was efficient in pyridine whereas Beaucage reagent performed well in acetonitrile. This was also true for some of the other sulfur transfer reagents we investigated. Thus, we were prompted to investigate the efficiency of PADS under a variety of conditions.

Encouraged by our initial experiments on the synthesis of homo-thymidine phosphorothioate 20-mer using PADS under new conditions, we tested the sulfurization efficiency in the solid phase synthesis of a mixed deoxyoligonucleotide sequence (5'-TCC-CGC-CTG-TGA-CAT-GCA-TT, ISIS 5132). Several experiments (ca. 75 syntheses) were performed at 180-190  $\mu$ mole scale using a Pharmacia OligoPilot II DNA/RNA synthesizer on polystyrene support (90  $\mu$ mole/gram loading) by changing solvents (dichloromethane, dichloroethane, acetonitrile), bases (pyridine, *sym*-collidine, lutidine, 2-picoline, 4-picoline, 3-picoline), sulfurization time (600, 480, 360, 100, 60 sec), solvent to base ratio (100:0, 75:25, 50:50, 25:75, 0:100), concentration (2.0, 1.0, 0.8, 0.6, 0.5, 0.4, 0.2, 0.1M), and column volume (1.5, 1.0, 0.5). Based on the data obtained from these experiments, it was found that a 0.2M solution of PADS (6 molar equivalents) in 1:1 (v/v) CH<sub>3</sub>CN:3-picoline for 60 sec gave a sulfurization efficiency of >99.6%. The oligonucleotide phosphorothioate synthesized under these optimized conditions was analysed by <sup>31</sup>P NMR, capillary gel electrophoresis (CGE), SAX-HPLC and by measuring crude yield by UV. The following table compares sulfur transfer and synthesis efficiencies of Beaucage reagent and that of PADS.

Reagent	Crude Yield OD/ $\mu$ mole	Crude Full Length (%)	Full Length (n-1) (%)	P=S : P=O ( <sup>31</sup> P NMR)	P=S : P=O (SAX HPLC)
Beaucage	115-120	72	91-92 (2.3)	99.4 : 0.6	99.5 : 0.5
PADS	118-125	73	91-92 (2.2)	99.6 : 0.4	99.6 : 0.4

The above results demonstrate that PADS is an efficient sulfur transfer reagent in the large scale synthesis of oligodeoxyribonucleotide phosphorothioates, fully equivalent in performance to Beaucage reagent. To verify the efficiency of this reagent

for different sequences, this optimized condition was used to synthesize four other oligodeoxyribonucleotide phosphorothioates (5'-GCC-CAA-GCT-GGC-ATC-CGT-CA, ISIS 2302; 5'-GTT-CTC-GCT-GGT-GAG-TTT-CA, ISIS 3521; 5'-TCC-GTC-ATC-GCT-CCT-CAG-GG, ISIS 2503; 5'-GCG-TTT-GCT-CTT-CTT-CTT-GCG, ISIS 2922) (20 to 21-mer in length) and again found to be an efficient sulfur transfer reagent. In addition, during our investigation we noted that PADS gave significantly more reproducible results than Beaucage reagent. In addition, a solution of PADS in acetonitrile/3-picoline (1:1, v/v) stored at room temperature for two weeks gave results similar to those from a freshly prepared solution.<sup>3</sup> Another major advantage of PADS over Beaucage reagent is its cost. The current cost of Beaucage reagent is ca \$6000/Kg as compared to PADS which is ca \$500/Kg.<sup>4</sup> In conclusion, we have demonstrated that phenylacetyl disulfide, if used under proper conditions, is a very effective replacement for Beaucage reagent in the synthesis of phosphorothioate oligonucleotides.

#### REFERENCES:

1. Kamer, P. C. J.; Roelen, H. C. P. F.; van den Elst, H.; van der Marel, G. A.; van Boom, J. H. *Tetrahedron Lett.* **1989**, *30*, 6757.
2. Roelen, H. C. P. F.; Kamer, P. C. J.; van den Elst, H.; van der Marel, G. A.; van Boom, J. H. *Recl. Trav. Chim. Pays-Bas.*, **1991**, *110*, 325.
3. The solution turns dark on storage at room temperature for extended period of time.
4. PADS is now commercially available from different vendors.