

This article was downloaded by:

On: 26 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713597286>

An Improved Method for the Preparation of 4-Cyano-2-butenyl-deoxyribonucleosidephosphoramidites

G. Duckworth^a; J. Ackroyd^a

^a Cruachem Ltd Todd Campus, West of Scotland Science Park, Glasgow, Scotland

To cite this Article Duckworth, G. and Ackroyd, J.(1999) 'An Improved Method for the Preparation of 4-Cyano-2-butenyl-deoxyribonucleosidephosphoramidites', *Nucleosides, Nucleotides and Nucleic Acids*, 18: 6, 1227 — 1229

To link to this Article: DOI: 10.1080/07328319908044671

URL: <http://dx.doi.org/10.1080/07328319908044671>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

AN IMPROVED METHOD FOR THE PREPARATION OF 4-CYANO-2-BUTENYL-DEOXYRIBONUCLEOSIDEPHOSPHORAMIDITES.

G. Duckworth and J. Ackroyd*

Cruachem Ltd

*Todd Campus, West of Scotland Science Park, Acre Road, Glasgow,
Scotland. G20 0UA*

ABSTRACT An improved alternative synthesis of 4-cyano-2-butenyldeoxy nucleoside-phosphoramidites in >100g quantities is described via reaction of the phosphordiamidites with 4-cyano-2-buten-1-ol.

Currently, many pharmaceutical companies are investing significant effort into the synthesis of potential antisense therapeutic oligonucleotides¹. In order to achieve a cost-effective synthetic route and to improve the stability of the resultant oligonucleotide, investigations continue into a range of protecting groups for the base², sugar³ and internucleotidic phosphate linkage⁴.

As part of our studies in this area, we recently had cause to synthesise deoxyribonucleoside-4-cyano-2-butenyl protected phosphoramidites as reported⁴. Our attempts to repeat this approach, although partially successful on a 5g scale, met with significant difficulties upon scale-up to >100g quantities and consequently, an alternative approach was sought. In our hands, the problem was the preparation and isolation of the cyanobutenylphosphitylating agent in sufficient quality and quantity.

Our exploration of alternative methods led us to consider the versatility of the phosphordiamidites⁵ (**5**), a class of compounds that have been used in the preparation of deoxynucleoside phosphoramidites⁶ and deoxydinucleoside phosphoramidites⁷, as a precursor to the cyanobutenylphosphoramidites (**6**). This approach has the added advantage of generating the deoxynucleoside phosphordiamidite (**5**) suitable for reaction, *in situ*, with




FIG. 1: I) SOCl_2 , Pyr, Et_2O , 0°C ; II) KCN, NaI, MeCN, 20°C ; III) PCl_3 , DCM, -60°C ; IV) DIPA, (3), DCM, 20°C ; V) Tetrazole, (2), DCM, 20°C

4-cyano-2-buten-1-ol (2). The chemical synthesis of the 4-cyano-2-butenylphosphoramidites (6) from the phosphordiamidite synthons is described.

4-Cyano-2-buten-1-ol (2) was prepared by methods previously reported i.e. monohalogenation⁸ of 2-butene-1,4-diol followed by cyanation⁴ of the resulting 4-chloro-2-buten-1-ol (1). Bisdiisopropylaminochlorophosphine (3) was prepared by adding a solution of phosphorus trichloride in dichloromethane dropwise to a stirred solution of diisopropylamine (6.0 equiv.) in dichloromethane. The reaction was carried out at -60°C . Once the addition of phosphorus trichloride was complete, the cooling bath was removed and the reactants stirred at room temperature over night. Diisopropylamine hydrochloride was removed by filtration and the product used without further purification.

5'-O-Dimethoxytritylthymidine⁹ (4a) in dichloromethane was treated with diisopropylamine (1.5 equiv.) and bisdiisopropylaminochlorophosphine (3) (1.1 equiv.).

After stirring at room temperature for 1.5 hours TLC (5% methanol/dichloromethane) indicated complete conversion of (**4a**). 4-Cyano-2-butene-1-ol (**2**) (1.1equiv.) and tetrazole (0.25 equiv.) were added. TLC (diethyl ether) after 2.5 hours showed incomplete reaction. Further tetrazole (0.25 equiv.) was added and the reactants stirred overnight. After the usual aqueous work up the product was purified by chromatography on silica gel using a gradient of ethyl acetate in hexane (N.B. all solvents contained 1.0% triethylamine). Yield = 58%. The remaining three phosphoramidites (**6b-d**) were prepared in a similar manner in 42%, 40% and 41% yields respectively, the spectroscopic data was as reported⁴.

In summary, this method provides an improved alternative route for the synthesis of 4-cyano-2-butenyl protected phosphoramidites and allows the preparation of the title compounds in >100g quantities.

REFERENCES

- 1 S.M. Freier, K.H. Altmann, *Nucleic Acid Research*, **1997**, *25*, 4429.
- 2 M.P. Reddy, N.B. Hanna, F. Farooqui, *Nucleosides and Nucleotides*, **1997**, *16*, 1589.
- 3 L. Beigelman, A. Karpeisky, J. Matulic-Adamic, P. Haeberli, D. Sweedler, N. Usman, *Nucleic Acid Research*, **1995**, *23*, 4434.
- 4 V.T. Ravikumar, Z.S. Cheruvallath, D.L. Cole, *Tetrahedron Letters*, **1996**, *37*, 6643.
- 5 G. Beaton, W.K.-D. Brill, A. Grandas, Y.-X. Ma, J. Nielsen, E. Yau, M. H. Caruthers, *Tetrahedron*, **1991**, *47*, 2377.
- 6 S.L. Beaucage, *Tet. Lett.*, **1984**, *25*, 375; A.D. Barone, J.-Y. Tang, M.H. Caruthers, *Nucleic Acid Research*, **1984**, *12*, 4051.
- 7 J.E. Marugg, A. Burik, M. Tromp, G.A. van der Marel, J.H. van Boom, *Tetrahedron. Letters*, **1986**, *27*, 2271; B. Uznanski, A. Wilk, W.S. Stec, *Tetrahedron Letters*, **1987**, *28*, 3401.
- 8 J. Cologne, G. Poilane, *Bull. Soc. Chim. Fr.*, **1955**, 953.
- 9 F. Morvan, B. Rayner, J.-L. Imbach. *In Methods in Molecular Biology, Protocols for Oligonucleotides and Analogs*. Edited by: S. Agarawal; Humana Press Inc., Totowa, New Jersey, **1993**; Vol. 20, 261-283.