

This article was downloaded by:

On: 26 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713597286>

New Phosphitylating Reagents Containing P-F Bond and Their Application in the Synthesis of P-Fluoro Nucleosidyl Phosphates, Thiophosphates and Selenophosphates

W. Dąbkowski^a; I. Tworowska^a; E. Poniatowska^a; J. Michalski^a; F. Cramer^b

^a Centre of Molecular and Macromolecular Studies, Polish Academy of Sciences, Sienkiewicza, Poland

^b Max-Planck-Institut für experimentelle Medizin, Hermann-Rein Strasse, Germany

To cite this Article Dąbkowski, W. , Tworowska, I. , Poniatowska, E. , Michalski, J. and Cramer, F.(1999) 'New Phosphitylating Reagents Containing P-F Bond and Their Application in the Synthesis of P-Fluoro Nucleosidyl Phosphates, Thiophosphates and Selenophosphates', *Nucleosides, Nucleotides and Nucleic Acids*, 18: 6, 1247 – 1248

To link to this Article: DOI: 10.1080/07328319908044679

URL: <http://dx.doi.org/10.1080/07328319908044679>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

NEW PHOSPHITYLATING REAGENTS CONTAINING P-F BOND AND THEIR APPLICATION IN THE SYNTHESIS OF P-FLUORO NUCLEOSIDYL PHOSPHATES, THIOPHOSPHATES AND SELENOPHOSPHATES

W. Dąbkowski,^a I. Tworowska,^a E. Poniatowska,^a J. Michalski^{a*} and F. Cramer^b


^aCentre of Molecular and Macromolecular Studies, Polish Academy of Sciences,
90-363 Łódź, Sienkiewicza 112, Poland

^bMax-Planck-Institut für experimentelle Medizin, 3400 Göttingen,
Hermann-Rein Strasse 3, Germany

Abstract: An efficient and general synthesis of phosphorofluoridates RO-P(X)(OH)F (X=O) and their analogues (X=S, Se) based on two new phosphitylating reagents: 2-cyanoethyl-N,N-diisopropylfluorophosphoroamidite F-P(NPr₂)OCH₂CH₂CN and tert-butyl-N,N-diisopropylfluorophosphoroamidite F-P(NPr₂)OBu^t is described.


Compounds of the general formula R-O-P(X)(F)OH **1** where X=O (R=nucleosid-3'-yl or nucleosid-5'-yl) have been first obtained by Wittmann ¹. Wittmann method and procedures described afterward are not fully satisfactory. We searched for more general synthesis of **1** especially providing approach towards thio- and selenofluorophosphates ².

In this communication we present an expedient general method based on a phosphoro-amide approach. New phosphitylating reagents containing P-F bond **2** and **3** were synthesized and applied in the synthesis of compounds **1** (X = O, S, Se). The synthetic strategy for the preparation of the phosphitylating reagents **2** and **3** is delineated below.



Preparation of amidites **2** and **3** can be performed in one-flask procedure in over 90% yield. Both compounds are stable colourless liquids which can be purified by distillation *in vacuo* or by silica gel column chromatography and stored at room temperature.

Phosphitylating reagents **2** and **3** react with alcohols in the presence of activators such as tetrazole, benzoyl chloride or trimethylsilyl chloride (TMSCl) at room temperature to give phosphorofluoridites **4** and **5** in very high yield. TMSCl proved to be superior to others.



Oxidation of phosphorofluoridites **4** and **5** by hydrogen tert-butyl peroxide or addition of elemental sulfur in the presence of triethylamine gave the corresponding phosphorofluoridates **6** and **7** which undergo thermal elimination of 2-methyl-1-propene (80°C, 2h) or β-elimination of vinyl cyanide [pyridine-triethylamine (3:1 v/v), r.t., 10 min] to form the final compounds **1** (X=O, S) in excellent yield. All these reactions are best performed as one-flask procedures. It is noteworthy that in the coupling procedure (**3** → **5**) activated by TMSCl the tert-butyl group in unaffected. Preliminary experiments showed that this procedure is applicable for the synthesis of **1** (X=Se).

Examination of the phosphitylation procedure by ³¹P and ¹⁹F NMR spectroscopy using phosphorofluoridite **2** clearly shows intermediate formation of fluorophosphorotetrazolidite FP(OCH₂CH₂CN)CHN₄ **8**. Formation of fluorochlorophosphine FP(OCH₂CH₂CN)Cl **9** was observed when amidite **2** was allowed to react with benzoyl chloride or TMSCl. It is most likely that the compounds **8** and **9** act as phosphitylating species towards alcohols.

Acknowledgment. This work was supported by the State Committee for Scientific Research, Poland (No 3 T09A 155 10), Stiftung für Deutsch-Polnische Zusammenarbeit (2789/96/LN) and German-Polish project (POL-211-96).

REFERENCES

1. Wittmann, R. *Chem. Ber.*, **1963**, *96*, 771-779.
2. Dąbkowski, W.; Tworowska, I. *Chemistry Letters*, **1995**, 727-728 and references therein.