This article was downloaded by:


On: 26 January 2011

Access details: Access Details: Free Access

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-

41 Mortimer Street, London W1T 3JH, UK



# Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information: http://www.informaworld.com/smpp/title~content=t713597286

## **Erratum**

To cite this Article (2000) 'Erratum', Nucleosides, Nucleotides and Nucleic Acids, 19: 3, 703

To link to this Article: DOI: 10.1080/15257770008035017 URL: http://dx.doi.org/10.1080/15257770008035017

# PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: http://www.informaworld.com/terms-and-conditions-of-access.pdf

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

#### **ERRATUM**

## SYNTHESIS OF OLIGODEOXYNUCLEOTIDES CONTAINING 2-SUBSTITUTED GUANINE DERIVATIVES USING 2-FLUORO-2'-DEOXYINOSINE AS COMMON NUCLEOSIDE PRECURSOR.

Antonio R. Díaz, Ramon Eritja\* and Ramon Güimil García.

Nucleosides & Nucleotides 16 (10&11), 2035-2051 (1997)

Page 2042. Melting temperatures of duplexes containing the control unmodified oligonucleotide were wrong. Table 2 should read as follows.

TABLE 2. Melting temperatures (°C) of duplexes containing N-2-substituted guanine derivatives in 0.15 M NaCl, 0.05 M tris.HCl pH 7.4.

5' CAC CXA CGG CGC 3' 3' GTG GYT GCC GCG 5'

		X		
Y	G	$G^{API}$	$G^{SPM}$	$G^{SPD}$
C	63	74	73	72
A	50	58	53	53
T	57	63	61	61
G	54	54	57	59