

This article was downloaded by:

On: 26 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713597286>

3'-DEOXYRIBOFURANOSE DERIVATIVES OF 1-DEAZA AND 3-DEAZA-ADENOSINE AND THEIR ACTIVITY AS ADENOSINE DEAMINASE INHIBITORS

S. Costanzi; C. Lambertucci; R. Volpini; S. Vittori; G. Lupidi^a; G. Cristalli

^a Dipartimento di Biologia M.C.A., University of Camerino, Camerino, Italy

Online publication date: 31 March 2001

To cite this Article Costanzi, S. , Lambertucci, C. , Volpini, R. , Vittori, S. , Lupidi, G. and Cristalli, G.(2001) '3'-DEOXYRIBOFURANOSE DERIVATIVES OF 1-DEAZA AND 3-DEAZA-ADENOSINE AND THEIR ACTIVITY AS ADENOSINE DEAMINASE INHIBITORS', *Nucleosides, Nucleotides and Nucleic Acids*, 20: 4, 1037 — 1041

To link to this Article: DOI: 10.1081/NCN-100002486

URL: <http://dx.doi.org/10.1081/NCN-100002486>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

3'-DEOXYRIBOFURANOSE DERIVATIVES OF 1-DEAZA AND 3-DEAZA-ADENOSINE AND THEIR ACTIVITY AS ADENOSINE DEAMINASE INHIBITORS

**S. Costanzi,¹ C. Lambertucci,¹ R. Volpini,¹ S. Vittori,¹
G. Lupidi,² and G. Cristalli^{1,*}**

¹Dipartimento di Scienze Chimiche, via S. Agostino 1, Italy

²Dipartimento di Biologia M.C.A., University of Camerino,
I-62032, Camerino, Italy

ABSTRACT

2,6-Dichloro-1-deazapurine and 2,6-dichloro-3-deazapurine were coupled with 1,2-*O*-diacetyl-5-*O*-benzoyl-3-deoxy- β -D-ribofuranose. Deprotection of the obtained compounds and reaction with liquid ammonia gave the desired 2-chloroadenine nucleosides, which were dechlorinated to afford the corresponding 1-deaza and 3-deazaadenosine derivatives. Biological studies performed on ADA from calf intestine showed that these new nucleosides are inhibitors of the enzyme.

INTRODUCTION

Adenosine deaminase (ADA) is an important catabolic enzyme, which converts adenosine and deoxyadenosine to inosine and deoxyinosine, respectively. Moreover ADA is involved in the pathogenesis of the Severe Combined Immuno-deficiency Disease (SCID) and is responsible for deamination of potentially useful drugs with a nucleosidic structure. Elevated serum ADA activity has been described in various autoimmune and inflammatory diseases, including Systemic Lupus Erythematosus (SLE) (1).

*Corresponding author.

Modifications of the purine moiety or/and substitution of the sugar moiety of adenosine with aliphatic chains led to derivatives with good ADA inhibitory activity (2). We reported in previous papers that 1-deazaadenosine analogues exhibit good inhibitory activity on ADA from several sources. In this research, we compared coupling of 2,6-dichloro-1-deazapurine or 3-deazapurine with 1,2-*O*-diacetyl-5-*O*-benzoyl-3-deoxy- β -D-ribofuranose and evaluated the obtained compounds as ADA inhibitors.

CHEMISTRY


Coupling of 2,6-dichloro-1-deazapurine (**1**) with 1,2-*O*-diacetyl-5-*O*-benzoyl-3-deoxy- β -D-ribofuranose (**2**), using tin tetrachloride as the acidic-catalyst, yielded only the β -anomer **3** (Scheme 1).

Deprotection with methanolic ammonia gave the dichloro-1-deazanucleoside **5**. On the other hand, coupling of compounds **1** and **2** using the fusion method yielded a mixture of β - and α -anomers as shown in Scheme 1.

Room temperature standing of the protected compounds with methanolic ammonia gave the corresponding dichloro-1-deazanucleosides **5** and **6** (3,4).


In the case of 2,6-dichloro-3-deazapurine (**7**), coupling with sugar **2** in the presence of tin tetrachloride gave the β -anomer **8** in low yield. Deprotection with methanolic ammonia afforded the dichloro-1-deazanucleoside **12**. 2,6-Dichloro-3-deazapurine (**7**) was also coupled with **2** under fusion reaction conditions as shown in Scheme 2 (5).

The glycosylation site and the anomeric configuration were assigned on the basis of UV data and ^1H NMR spectra, including 1D ^1H NOE difference spectra, of the deprotected nucleosides **12**–**15**, obtained by treating compounds **8**–**11** with methanolic ammonia at room temperature. In a previous work we isolated,



Scheme 1.






Scheme 2.

besides the N(9)- β - and the N(9)- α -3'-deoxyribonucleosides **12** and **13**, also the N(9)- α -3'-deoxyarabinonucleoside **15** (5). Now with further studies we also identified the N(9)- β -3'-deoxyarabino derivative **14**.

The UV spectra of the four nucleosides were practically identical. Saturation of H-C(1') resulted in a NOE at the H-C(8) and H-C(3) signals in all four cases. The latter excludes without any doubt N(7) as glycosylation site. NOE at the H-C(4') upon saturation of H-C(1') of **12** established the β configuration. In the case of **14**, saturation of H-C(1') gave a strong NOE at the H-C(2') signal but not at the H-C(4') one. This is probably due to the distortion induced in the sugar moiety by the different position of the OH-C(2'). NOE at the Ha-C(3') upon saturation of H-C(1') of **13** and **15** established the α configuration. The epimerization at C(2') of nucleosides **14** and **15** has been demonstrated by the NOE observed at the H_b-C(3') signal upon saturation of either H-C(2') or H-C(4'). This result confirmed that the three protons are located on the same side of the sugar ring.


Reaction of compounds **5** and **12** with liquid ammonia at 130°C gave the nucleosides **16** and **18**, which were dechlorinated by catalytic hydrogenolysis to obtain the corresponding 3'-deoxy-1-deazaadenosine (**17**) and 3'-deoxy-3-deazaadenosine (**19**) as shown in Scheme 3. The same procedure could be utilized in order to introduce different amines in 6 position of these molecules.





Scheme 3.

Table 1. ADA Inhibitory Activity of 1-deaza and 3-deaza Adenosines

Cpd.	X	Y	Z	R	R ₁	K _i (μM)	
Ia	CH	N	Cl	OH	OH	30	
Ib	CH	N	H	OH	OH	0.38	
Ic	CH	N	Cl	H	OH	23	
Id	CH	N	H	H	OH	0.19	
16	CH	N	Cl	OH	H	>100	
17	CH	N	H	OH	H	2.5	
IIa	N	CH	Cl	OH	OH	37	
IIb	N	CH	H	OH	OH	400	
IIc	N	CH	Cl	H	OH	61	
18	N	CH	Cl	OH	H	152	
19	N	CH	H	OH	H	411	

RESULTS

The 3'-deoxy-1-deazaadenosines (**16** and **17**) and 3'-deoxy-3-deazaadenosines (**18** and **19**) were tested for their ability to inhibit calf intestine ADA (6). In Table 1 the activity of these compounds is compared to that of the corresponding ribose and 2'-deoxyribose derivatives.

2'-Deoxy-1-deazaadenosine (**Id**, K_i = 0.19 μM) proved to be the most potent compound of the series, and it resulted more active than 1-deazaadenosine itself (**Ib**, K_i = 0.38 μM). The 3-deaza derivatives are still inhibitors of the enzyme although always less potent than the corresponding 1-deaza analogues.

It is interesting to note that in the case of the 1-deaza nucleosides the presence of a chlorine atom on C(2) produces a decrease in the ADA inhibitory activity (**Ib**, K_i = 0.38 μM vs **Ia**, K_i = 30 μM), while in the case of the 3-deaza analogues the same substitution leads to more potent compounds (**IIb**, K_i = 400 μM vs **IIa**, K_i = 37 μM).



ACKNOWLEDGMENTS

Work supported by the Italian Ministry of Research (Grant n. 9905155535, 1999) and University of Camerino (Fondo di Ricerca d'Ateneo).

REFERENCES

1. Cristalli, G.; Costanzi, S.; Lambertucci, C.; Lupidi, G.; Vittori, S.; Volpini, R.; Camaioni, E. *Med. Res. Rev.*, in press.
2. Cristalli, G.; Eleuteri, A.; Vittori, S.; Volpini, R.; Camaioni, E.; Lupidi, G. *Drug. Dev. Res.*, **1993**, 28, 253–258.
3. Vittori, S.; Camaioni, E.; Costanzi, S.; Volpini, R.; Cristalli, G. *Nucleosides & Nucleotides*, **1999**, 18, 587–590.
4. Volpini, R.; Costanzi, S.; Vittori, S.; Cristalli, G.; Lupidi, G. *Helv. Chim. Acta*, **1999**, 82, 2112–2118.
5. Volpini, R.; Camaioni, E.; Costanzi, S.; Vittori, S.; Cristalli, G. *Helv. Chim. Acta*, **1998**, 81, 2326–2331.
6. Lupidi, G.; Cristalli, G.; Marmocchi, F.; Riva, F.; Grifantini, M. *J. Enzyme Inhib.*, **1985**, 31, 1179–1183.



Request Permission or Order Reprints Instantly!

Interested in copying and sharing this article? In most cases, U.S. Copyright Law requires that you get permission from the article's rightsholder before using copyrighted content.

All information and materials found in this article, including but not limited to text, trademarks, patents, logos, graphics and images (the "Materials"), are the copyrighted works and other forms of intellectual property of Marcel Dekker, Inc., or its licensors. All rights not expressly granted are reserved.

Get permission to lawfully reproduce and distribute the Materials or order reprints quickly and painlessly. Simply click on the "Request Permission/Reprints Here" link below and follow the instructions. Visit the [U.S. Copyright Office](#) for information on Fair Use limitations of U.S. copyright law. Please refer to The Association of American Publishers' (AAP) website for guidelines on [Fair Use in the Classroom](#).

The Materials are for your personal use only and cannot be reformatted, reposted, resold or distributed by electronic means or otherwise without permission from Marcel Dekker, Inc. Marcel Dekker, Inc. grants you the limited right to display the Materials only on your personal computer or personal wireless device, and to copy and download single copies of such Materials provided that any copyright, trademark or other notice appearing on such Materials is also retained by, displayed, copied or downloaded as part of the Materials and is not removed or obscured, and provided you do not edit, modify, alter or enhance the Materials. Please refer to our [Website User Agreement](#) for more details.

[Order now!](#)

Reprints of this article can also be ordered at

<http://www.dekker.com/servlet/product/DOI/101081NCN100002486>