

This article was downloaded by:

On: 26 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713597286>

### SYNTHESIS OF 3'-THIOAMIDO-MODIFIED 3'-DEOXYTHYMIDINE-5'-TRIPHOSPHATES AND THEIR USE AS CHAIN TERMINATORS IN SANGER-DNA SEQUENCING

K. Schwarzer<sup>a</sup>; C. Wojczewski<sup>a</sup>; J. W. Engels<sup>a</sup>

<sup>a</sup> Johann Wolfgang Goethe-Universität, Frankfurt am Main, Germany

Online publication date: 31 March 2001

**To cite this Article** Schwarzer, K. , Wojczewski, C. and Engels, J. W.(2001) 'SYNTHESIS OF 3'-THIOAMIDO-MODIFIED 3'-DEOXYTHYMIDINE-5'-TRIPHOSPHATES AND THEIR USE AS CHAIN TERMINATORS IN SANGER-DNA SEQUENCING', *Nucleosides, Nucleotides and Nucleic Acids*, 20: 4, 879 — 882

**To link to this Article:** DOI: 10.1081/NCN-100002450

**URL:** <http://dx.doi.org/10.1081/NCN-100002450>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

## SYNTHESIS OF 3'-THIOAMIDO-MODIFIED 3'-DEOXYTHYMIDINE-5'-TRIPHOSPHATES AND THEIR USE AS CHAIN TERMINATORS IN SANGER-DNA SEQUENCING

K. Schwarzer, C. Wojczewski, and J. W. Engels\*

Johann Wolfgang Goethe-Universität, Frankfurt am Main, Germany

### ABSTRACT

The thioamide derivatives 3'-deoxy-5'-*O*-(4,4'-dimethoxytrityl)-3'-[(2-methyl-1-thioxo-propyl)amino]thymidine **1** and 3'-deoxy-5'-*O*-(4,4'-dimethoxytrityl)-3'-{6-[[*(9H*-(fluoren-9-ylmethoxy)carbonyl]-amino)-1-thioxohexyl]amino}thymidine **2** were synthesized by regioselective thionation of their corresponding amides **7** and **8** with 2,4-bis(4-methoxyphenyl)-1,3,2,4-dithiadiphosphetane-2,4-disulfide (*Lawesson's* reagent). The thioamides were converted into the corresponding 5'-triphosphates **3** and **4**. Compound **3** was chosen for DNA sequencing experiments and **4** was further labelled with fluorescein.

### INTRODUCTION

Dye-terminator sequencing is characterized by an enzymatic incorporation of chain-terminating, fluorescent nucleotides. These terminators are attached to fluorescent dyes at their base moiety. Apart from this labeling strategy, there are other potential positions within nucleosides that may be used for labeling. We have shown that 3'-amino- and amido-modified nucleoside-5'-triphosphates are excellent terminators (1,2). However, the ability of DNA polymerases to hydrolyze ester- and amide-bonds at the 3'-position limits the use of these 3'-dye-terminators for DNA sequencing since the label is lost during the incorporation process (3). Therefore, we

---

\*Corresponding author.

decided to alter the link between dye and sugar moiety to find a bond that is stable against enzymatic degradation. In this context, we describe an efficient synthesis of 3'-thioamido-modified thymidines. We synthesized the thioamide derivative **1** as a model compound for the investigation of the thionation reaction with *Lawesson's* reagent. In a second synthetic route we optimized the thionation reaction. The addition of exact amounts of pyridine to the reaction mixture proved to be essential for an efficient transformation. The corresponding 5'-triphosphate **4** was further labeled with fluorescein and **3** was chosen for DNA sequencing experiments.

## RESULTS AND DISCUSSION

A broad spectrum of reactions exists for the synthesis of thioamides, only the conversion of amides into thioamides seemed to be suitable for nucleoside chemistry. Therefore, we synthesised the readily accessible amide **7** as a model compound for the investigation of the thionation reaction (Fig. 1). Direct action of hydrogen sulfide or phosphorus pentasulfide on **7/8** would result in an undesired exchange of the two carbonyl oxygen atoms at C(2) and C(4) of the base moiety. A more sophisticated thionation reagent is *Lawesson's* reagent, which converts carbonyl groups into thiocarbonyl groups in high yield (4–6). Therefore, **7** was reacted with *Lawesson's* reagent for 30 minutes at RT.

No by-products were observed under these reaction conditions, but the yield of **1** was rather poor (22%). 33% of the reactant was recovered unchanged. The synthesis of **2** under the same conditions resulted in even lower yields caused mainly by 5'-hydroxy deprotection. Therefore we tested basic solvents as scavengers for the phosphoric acids.

Thionation of **8** in pure anh. pyridine instead of anh. THF did not at all indicate any formation of **2** even when the amount of *Lawesson's* reagent was drastically increased. Thus, the reaction of **8** with *Lawesson's* reagent was accomplished in mixtures of anh. THF/pyridine. We observed that lowering the amount of




Figure 1. Synthesis of the thioamide derivatives **1** and **2**.

pyridine gradually resulted in an increased yield. Finally, the addition of **1** equiv. of pyridine and 4 equiv. of *Lawesson's* reagent in anh. THF as solvent yielded **2** in 50%; almost 50% of the intact educt was recovered. **1** and **2** were deprotected in 80% aq. AcOH solution at room temperature. The syntheses of the triphosphates **3** and **4** were achieved according to the procedure of Ludwig and Eckstein (7). The aminofunction of **4** was deprotected with 20% piperidine in pyridine and DMF at room temperature and succeedingly coupled with fluoresceinisothiocyanate (FITC) in aq. 0.1M NaHCO<sub>3</sub> (pH = 9.3) and DMF at room temperature for 24 hours. Both thioamides **1/2** were clearly identified by mass spectrometry and NMR spectroscopy. The electrospray-ionization MS (ESI-MS) showed formation of only one product with an increased weight of 16 Da compared with the educt, representing the exchange of an O-atom against a S-atom. The <sup>13</sup>C-NMR spectrum revealed the identity of the carbonyl group involved. The <sup>13</sup>C-chemical shifts of the carbon atoms C(2) and C(4) of the base moiety remained unchanged, whereas the carbonyl signal at the 3'-terminal moved downfield as expected.

The thioamides **1** and **2** were converted into the corresponding 5'-triphosphates **3** and **4**. Compound **4** was further labelled with fluorescein and **3** was investigated in DNA sequencing experiments (8). No band pattern was detected with Sequenase or with Thermosequenase. With Taq DNA polymerase, we obtained a band pattern which did not correlate with the standard ddTTP sequence. It was, however, identical with the band pattern obtained with a 3'-thioether 3'-deoxy-thymidine-5'-triphosphate. Although **3** can obviously not be defined as specific terminator for the enzymatic DNA synthesis under the chosen conditions, the result can not be called nonselective, since two structurally different nucleotides were incorporated at exactly the same positions within the DNA. This excludes a random process. Variation of the terminator concentration only resulted in a different fragment length distribution. The reason for this could be a conformationally driven selection process.

## CONCLUSION

We demonstrated that 3'-thioamido modified nucleosides can efficiently be synthesised from 3'-amido tethered nucleosides by a regioselective thionation strategy applying *Lawesson's* reagent. Addition of 1 equiv. of pyridine to the reaction mixture was necessary to prevent degradation of the nucleoside and ensure good yields at the same time. Thioamide **3** was accepted as a substrate by the Taq DNA polymerase. However, specific incorporation did not occur. Rather the band pattern showed a distinct correlation with a thioether tethered nucleoside triphosphate.

## ACKNOWLEDGMENTS

We would like to thank the Bundesministerium für Bildung und Forschung (BEO0311088), the Verband der Chemischen Industrie and the Deutsche Forschungsgemeinschaft (Graduiertenkolleg) for financial support as well as Roche, 270 Madison Avenue, New York, New York 10016.



Diagnostics for cooperation and financial support. DNA sequencing experiments were done by Dr. S. Klingel and Dr. G. Sagner at Roche Diagnostics, Penzberg, Germany.

### REFERENCES

1. Herrlein, M. K.; Konrad, R. E.; Engels, J. W.; Holletz, T.; Cech, D., *Helv. Chim. Acta* **1994**, *77*, 586.
2. Wojczewski, C.; Faulstich, K.; Engels, J. W., *Nucleosides Nucleotides* **1997**, *16*, 751.
3. Canard, B.; Cardona, B.; Sarfati, R. S., *Proc. Natl. Acad. Sci. USA* **1995**, *92*, 10859.
4. Yde, B.; Yousif, N. M.; Pedersen, U.; Thomsen, I.; Lawesson, S.-O., *Tetrahedron* **1984**, *40*, 2047.
5. Cava, M. P.; Levinson, M. I., *Tetrahedron* **1985**, *41*, 5061.
6. Cherkasov, R. A.; Kutyrev, G. A.; Pudovik, A. N., *Tetrahedron* **1985**, *41*, 2567.
7. Ludwig, J.; Eckstein, F., *J. Org. Chem.* **1989**, *54*, 631.
8. Wojczewski, C.; Schwarzer, K.; Engels, J. W., *Helv. Chim. Acta* **2000**, *83*, 1268.



## **Request Permission or Order Reprints Instantly!**

Interested in copying and sharing this article? In most cases, U.S. Copyright Law requires that you get permission from the article's rightsholder before using copyrighted content.

All information and materials found in this article, including but not limited to text, trademarks, patents, logos, graphics and images (the "Materials"), are the copyrighted works and other forms of intellectual property of Marcel Dekker, Inc., or its licensors. All rights not expressly granted are reserved.

Get permission to lawfully reproduce and distribute the Materials or order reprints quickly and painlessly. Simply click on the "Request Permission/Reprints Here" link below and follow the instructions. Visit the [U.S. Copyright Office](#) for information on Fair Use limitations of U.S. copyright law. Please refer to The Association of American Publishers' (AAP) website for guidelines on [Fair Use in the Classroom](#).

The Materials are for your personal use only and cannot be reformatted, reposted, resold or distributed by electronic means or otherwise without permission from Marcel Dekker, Inc. Marcel Dekker, Inc. grants you the limited right to display the Materials only on your personal computer or personal wireless device, and to copy and download single copies of such Materials provided that any copyright, trademark or other notice appearing on such Materials is also retained by, displayed, copied or downloaded as part of the Materials and is not removed or obscured, and provided you do not edit, modify, alter or enhance the Materials. Please refer to our [Website User Agreement](#) for more details.

**[Order now!](#)**

Reprints of this article can also be ordered at

<http://www.dekker.com/servlet/product/DOI/101081NCN100002450>