

This article was downloaded by:

On: 26 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713597286>

Synthesis of a Base-Protected α -L-LNA Guanine Nucleoside

Gilles Gaubert^a; Jesper Wengel^a

^a Nucleic Acid Center, Department of Chemistry, University of Southern Denmark, Odense M, Denmark

Online publication date: 09 August 2003

To cite this Article Gaubert, Gilles and Wengel, Jesper(2003) 'Synthesis of a Base-Protected α -L-LNA Guanine Nucleoside', *Nucleosides, Nucleotides and Nucleic Acids*, 22: 5, 1155 — 1157

To link to this Article: DOI: 10.1081/NCN-120022824

URL: <http://dx.doi.org/10.1081/NCN-120022824>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Synthesis of a Base-Protected α -L-LNA Guanine Nucleoside

Gilles Gaubert and Jesper Wengel*

Nucleic Acid Center, Department of Chemistry, University of Southern
Denmark, Odense M, Denmark

ABSTRACT

Synthesis of (1*R*,3*R*,4*S*,7*R*)-7-hydroxy-1-hydroxymethyl-3-(2-*N*-isobutyroylguanin-9-yl)-2,5-dioxabicyclo[2.2.1]heptane (**12**), a protected α -L-LNA guanine nucleoside, has been accomplished using a convergent synthetic strategy starting from 1,2-di-*O*-acetylfuranose **4**.


Key Words: LNA (locked nucleic acids); α -L-LNA; Guanine nucleoside.

The unprecedented thermal stability of duplexes involving LNA^[1] (Locked Nucleic Acid) and α -L-LNA^[2] (α -L-*ribo* configured LNA; adenin-9-yl, 5-methylcytosin-1-yl and thymine-1-yl monomers) has stimulated us to synthesize the guanine α -L-LNA derivative **12** (Sch. 1).

Coupling between di-*O*-acetyl-di-*O*-mesyl furanose **4**^[2] and per(trimethylsilyl)-2-*N*-acetyl-6-*O*-(diphenylcarbamoyl) guanine^[3] afforded stereoselectively the desired *N*-9 regioisomer **5** in 59% yield. Treatment of nucleoside **5** with half-saturated methanolic ammonia gave chemoselectively the *O*-deacylated derivative **6** in 95% yield. To accomplish epimerisation at C2', nucleoside **6** was reacted with trifluoromethanesulfonic anhydride in a mixture of pyridine and dichloromethane and

*Correspondence: Jesper Wengel, Nucleic Acid Center, Department of Chemistry, University of Southern Denmark, Campusvej 55, DK-5230 Odense M, Denmark; Fax: +45 66 15 87 80; E-mail: jwe@chem.sdu.dk.





Scheme 1. Reagents and conditions: a) $G^{Ac,DPC}(SiMe_3)_2$, TMSOTf, toluene, 59%; b) half sat. NH_3 , MeOH, 95%; c) 1) Tf_2O , pyridine, CH_2Cl_2 , 2) AcOK, 18-crown-6, toluene, 63%; d) half sat. NH_3 , MeOH, 96%; e) 1) NaH, THF; 2) AcOK, 18-crown-6, dioxane; f) NH_3 , MeOH, 35%; g) 1) TMSCl, pyridine; 2) $t-Bu_2O$; 3) NH_4OH , 72%; h) HCO_2NH_4 , Pd/C, MeOH, 46%.

subsequently with potassium acetate and 18-crown-6 in toluene to afford the desired derivative **7** in 46% yield. Selective O-deacylation of **7** using half-saturated methanolic ammonia gave in 96% yield nucleoside **8**, the C2'-epimer of nucleoside **6**. Nucleoside **8** was treated with sodium hydride in THF to induce cyclization, which was followed by substitution of the remaining mesyloxy group with an acetate group by reaction with potassium acetate and 18-crown-6 in dioxane at reflux, affording the bicyclic nucleoside derivative **9**. Complete deacetylation of **9** in saturated methanolic ammonia gave nucleoside **10** in 35% yield (from compound **8**). 2-N-Isobutyroyl protection of **10** was accomplished in 72% yield using transient protection^[4] of the free hydroxy groups followed by acylation of the amine function giving nucleoside **11**. Debenzylation of this intermediate (ammonium formate, Pd/C, 46% yield) eventually afforded the desired 2-N-isobutyroyl α -L-LNA guanine nucleoside **12** (Sch. 1).

In spite of many synthetic steps leading to a relatively low overall yield, a viable strategy for the preparation of a α -L-LNA nucleoside of guanine has been developed.

ACKNOWLEDGMENTS

Exiqon A/S is thanked for financial support.

REFERENCES

1. Koshkin, A.A.; Singh, S.K.; Nielsen, P.; Rajwanshi, V.K.; Kumar, R.; Meldgaard, M.; Olsen, C.E.; Wengel, J. LNA (Locked Nucleic Acids): Synthesis of the adenine, cytosine, guanine, 5-methylcytosine, thymine and uracil bicyclic nucleoside monomers, oligomerisation, and unprecedented nucleic acid recognition. *Tetrahedron* **1998**, *54*, 3607 pp.

2. Sørensen, M.D.; Kværnø, L.; Bryld, T.; Håkansson, A.E.; Verbeure, B.; Gaubert, G.; Herdewijn, P.; Wengel, J. α -L-ribo-Configured Locked Nucleic Acid (α -L-LNA): Synthesis and properties. *J. Am. Chem. Soc.* **2002**, *124*, 2164 pp.
3. Robins, M.J.; Zou, R.; Guo, Z.; Wnuk, S.F. Nucleic acid related compounds. 93. A solution for the historic problem of regioselective sugar-base coupling to produce 9-glycosylguanines or 7-glycosylguanines. *J. Org. Chem.* **1996**, *61*, 9207 pp.
4. Ti, G.S.; Gaffney, B.L.; Jones, R.A. Transient protection: efficient one-flask syntheses of protected deoxynucleosides. *J. Am. Chem. Soc.* **1982**, *104*, 1316 pp.



