

This article was downloaded by:

On: 26 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713597286>

### Oxidative Transformations of Nucleoside Fluorenemethyl H-Phosphonoselenoate Diesters

Martin Kullberg<sup>a</sup>; Jacek Stawinski<sup>ab</sup>

<sup>a</sup> Department of Organic Chemistry, Arrhenius Laboratory, Stockholm University, Stockholm, Sweden

<sup>b</sup> Institute of Bioorganic Chemistry, Polish Academy of Sciences, Poznan, Poland

**To cite this Article** Kullberg, Martin and Stawinski, Jacek(2005) 'Oxidative Transformations of Nucleoside Fluorenemethyl H-Phosphonoselenoate Diesters', *Nucleosides, Nucleotides and Nucleic Acids*, 24: 5, 659 — 661

**To link to this Article:** DOI: 10.1081/NCN-200060173

**URL:** <http://dx.doi.org/10.1081/NCN-200060173>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

## OXIDATIVE TRANSFORMATIONS OF NUCLEOSIDE FLUORENEMETHYL H-PHOSPHONSELENOATE DIESTERS

**Martin Kullberg** □ *Department of Organic Chemistry, Arrhenius Laboratory, Stockholm University, Stockholm, Sweden*

**Jacek Stawinski** □ *Department of Organic Chemistry, Arrhenius Laboratory, Stockholm University, Stockholm, Sweden and Institute of Bioorganic Chemistry, Polish Academy of Sciences, Poznan, Poland*

□ *9-Fluorenmethyl H-phosphonoselenoate monoester has been used to produce thymidine 3'-O-phosphoroselenoate monoester from which various P(V) derivatives containing multiple modifications at phosphorus were obtained; e.g., thymidine 3'-O-phosphoroselenofluoridate, 3'-O-phosphoroselenothioate, or 3'-O-phosphorodiselenoate monoesters.*

### INTRODUCTION

We have previously reported on the use of 9-fluorenmethyl H-phosphonoselenoate monoester as a useful reagent for transferring of an H-phosphonoselenoate moiety<sup>[1]</sup> to nucleosides. In this article, we expand the uses of 9-fluorenmethyl H-phosphonoselenoate monoester by performing several oxidative transformations on the intermediate nucleoside fluorenmethyl H-phosphonoselenoate diester **1** prior to the removal of the fluorenmethyl protecting group. This gives access to several novel phosphoroselenoate monoesters.

### RESULTS AND DISCUSSION

The nucleoside fluorenmethyl H-phosphonoselenoate diester **1** is easily available through previously published procedures.<sup>[1]</sup> The most basic oxidative transformation is oxidation of **1** into the corresponding phosphoroselenoate diester **3**. By performing this reaction using standard protocol developed for H-phosphonate diesters<sup>[2]</sup> a significant deselenization occurred, even with equimolar

Financial support from the Swedish Research Council is gratefully acknowledged.


Address correspondence to Jacek Stawinski, Institute of Bioorganic Chemistry, Polish Academy of Sciences, Noskowskiego 12/14, Poznan 61-704, Poland.

amounts of iodine used. However, when the reaction was run in MeCN with iodine (1 eq), pyridine (20 eq), and water (60 eq), the deselenization of the product could be almost completely suppressed. Changing from iodine to carbon tetrachloride as an oxidizing agent simplified further the synthetic protocol. The reaction with CCl<sub>4</sub> (10 eq), pyridine (20 eq), and water (60 eq) in MeCN resulted in clean and fast (15 min) formation of the desired phosphoroselenochloridate **2**, which upon hydrolysis afforded product **3** in isolated yield of 80%.

Deprotection of diester **3** to produce monoester **4** using the protocol developed for the sulfur analogue<sup>[3]</sup> that involved treatment with aqueous ammonia was accompanied by severe side products formation. It was found that the cleanest deprotection of **3** could be achieved by using TMS-Cl (10 eq) and TEA (20 eq) in pyridine (10 min). Under such conditions, **4** was the only observable product and it was isolated in 96% yield by precipitation from diethyl ether. The obtained phosphoroselenoate monoester **4** was, however, unstable in aqueous solution and loses selenium within hours, in contrast to the corresponding diesters of type **3**, which were much more stable (Scheme 1).

The intermediate phosphoroselenochloridate **2** can also be treated with 1 eq triethylammonium trifluoroborate (TAF) to produce fast (5 min) and quantitatively (<sup>31</sup>P NMR) the corresponding phosphoroselenofluoridate **5**. Crude **5** could easily be deprotected using TEA (10 eq) in pyridine and the monoester **6** was isolated in 93% yield.

The diastereoisomers of **1** could easily be separated by flash column chromatography and their sulfurization gave rise to novel phosphate monoester analogues, phosphoroselenothioates, with a chiral phosphorus center. Sulfurization of **1** with elemental sulfur (3 eq) and pyridine (20 eq) in MeCN was uneventful and



**SCHEME 1** Oxidative transformations of nucleoside fluorenylmethyl H-phosphonoselenoate diesters.

yielded within 20 min the diesters **7a** and **7b** essentially quantitatively ( $^{31}\text{P}$  NMR; isolated yields ca 90%).

The deprotection of diesters **7a** and **7b** using aqueous conditions turned out to be a difficult task. Changing to anhydrous conditions with thiophenol and triethyl amine gave the desired monoesters **8a** and **8b**, but the reaction was slow (2 h) and side products were formed over time. However, using piperidine (20 eq) in pyridine furnished clean formation of the desired products. The produced ammonium salts of **8a** and **8b** were somewhat unstable, but titration of the salts with DBU stabilized the products sufficiently to permit their isolation and characterization. These monoesters were, however, even more prone to degradation than phosphoroselenoate **4** and in the presence of air or moisture, compound **8** decomposed within hours.

Selenization of **1** turned out to be less straightforward than sulfurization because using elemental selenium and pyridine in MeCN gave slow conversion and some by-products formation. Changing to KSeCN as a selenizing agent gave a somewhat faster reaction, but formation of side products was not completely suppressed. Attempts to use triphenyl phosphine selenide or triphenyl phosphoroselenoates gave no formation of the desired product. The best results were obtained with selenization of **1** with 1.1 eq of 3*H*-1,2-benzothiaselenol-3-one<sup>[4]</sup> (BTSe) and 20 eq of pyridine in MeCN. This gave clean and fast (5 min) formation of diselenoate **9**, which upon treatment with piperidine (20 eq) afforded monoester **10** ( $^{31}\text{P}$  NMR). Unfortunately, phosphorodiselenoate **10** was too unstable to allow its isolation and more detailed characterization.

## CONCLUSION

Nucleoside 9-fluorenemethyl H-phosphonoselenoate monoester **1** was found to be a convenient starting material for the preparation of various P(V) derivatives with multiple modifications at the phosphorus center. By oxidative transformations of **1**, several novel selenium-containing nucleotide analogues have been synthesized.

## REFERENCES

1. Kullberg, M.; Stawinski, J. 9-Fluorenemethyl H-phosphonoselenoate—a versatile reagent for transferring an H-phosphonoselenoate group. *Nucleosides Nucleotides Nucleic Acids* **2003**, *22*, 1463–1465.
2. Garegg, P.J.; Regberg, T.; Stawinski, J.; Strömberg, R. Studies on the oxidation of nucleoside Hydrogenphosphonates. *Nucleosides Nucleotides* **1987**, *6*, 429–432.
3. Jankowska, J.; Cieslak, J.; Kraszewski, A.; Stawinski, J. 9-Fluorenemethyl H-phosphonothioate, a versatile reagent for the preparation of H-phosphonothioate, phosphorothioate, and phosphorodithioate monoesters. *Tetrahedron Lett.* **1997**, *38*, 2007–2010.
4. Stawinski, J.; Thelin, M. Nucleoside H-phosphonates. 14. Synthesis of nucleoside phosphoroselenoates and phosphorothioselenoates via stereospecific selenization of the corresponding H-phosphonate and H-phosphonothioate diesters with the aid of new selenium-transfer reagent 3*H*-1,2-benzothiaselenol-3-one. *J. Org. Chem.* **1994**, *59*, 130–136.