

This article was downloaded by:

On: 26 January 2011

Access details: Access Details: Free Access

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713597286>

An Intramolecular 1,3-Dipolar Cycloaddition Reaction Towards the Synthesis of Chiral Azetidine Nucleoside Analogues: the d-glucose Case

P. Pádár^a; P. Forgó^b; Z. Kele^a; N. M. Howarth^c; L. Kovács^a

^a Department of Medicinal Chemistry, Heriot-Watt University, Riccarton, Edinburgh, UK ^b Department of Organic Chemistry, University of Szeged, Szeged, Hungary ^c Department of Chemistry, School of Engineering and Physical Sciences, Heriot-Watt University, Riccarton, Edinburgh, UK

To cite this Article Pádár, P. , Forgó, P. , Kele, Z. , Howarth, N. M. and Kovács, L.(2005) 'An Intramolecular 1,3-Dipolar Cycloaddition Reaction Towards the Synthesis of Chiral Azetidine Nucleoside Analogues: the d-glucose Case', Nucleosides, Nucleotides and Nucleic Acids, 24: 5, 743 — 745

To link to this Article: DOI: 10.1081/NCN-200060056

URL: <http://dx.doi.org/10.1081/NCN-200060056>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

AN INTRAMOLECULAR 1,3-DIPOLAR CYCLOADDITION REACTION TOWARDS THE SYNTHESIS OF CHIRAL AZETIDINE NUCLEOSIDE ANALOGUES: THE D-*GLUCO* CASE

P. Pádár □ *Department of Medicinal Chemistry, Heriot-Watt University, Riccarton, Edinburgh, UK*

P. Forgó □ *Department of Organic Chemistry, University of Szeged, Szeged, Hungary*

Z. Kele □ *Department of Medicinal Chemistry, Heriot-Watt University, Riccarton, Edinburgh, UK*

N. M. Howarth □ *Department of Chemistry, School of Engineering and Physical Sciences, Heriot-Watt University, Riccarton, Edinburgh, UK*

L. Kovács □ *Department of Medicinal Chemistry, Heriot-Watt University, Riccarton, Edinburgh, UK*

□ *The diastereoselective intramolecular 1,3-dipolar cycloaddition reaction of unsaturated nitrones, derived from methyl α -D-glucopyranoside with 2-furaldehyde and 2-(benzyloxy)acetaldehyde has been studied. In our previous studies with 2-furaldehyde, the cycloaddition resulted 3 diastereoisomers in a 3:1:1 ratio. In this article, how the number of the possible isomers generated by 1,3-cycloaddition could be reduced from 4 to 1 when 2-(benzyloxy)acetaldehyde was employed as an aldehyde is shown.*

RESULTS AND DISCUSSION

In an attempt to overcome some of the undesirable properties of peptide nucleic acids (PNAs), we have designed conformationally restricted, chiral azetidine nucleic acids (ANAs). The key step in the synthesis of the ANA monomers, which are needed for construction of the oligomers, is a diastereoselective intramolecular 1,3-dipolar cycloaddition reaction involving unsaturated nitrones derived from carbohydrate precursors (Figure 1). The intramolecular 1,3-

The financial support of grants Wellcome Trust 063879/Z/01/Z and EU FP6 TRIoH LSHB-CT-2003-503480 is gratefully acknowledged.

Address correspondence to L. Kovács, Department of Medicinal Chemistry, Heriot-Watt University, Riccarton, Edinburgh, UK; E-mail: kovacs@ovrisc.mdche.u-szeged.hu




FIGURE 1 a) Zn, sonication^[3] or Zn and Co(II)-phthalocyanine,^[4] b) NH₂OH · HCl, NaHCO₃,^[5] c) NaBH₃CN, HCl/dioxane, d) 2-furaldehyde or 2-(benzyloxy)acetaldehyde, toluene, 4 Å MS, 50°C, 18 h, e) toluene, 120°C, 4 Å MS, Lewis acid catalysts, **7**, **16**: 9-oxa-1-azabicyclo[4.2.1]nonane skeleton.

dipolar cycloaddition has the potential to reduce the number of possible resulting isomers by virtue of steric constraint. Thus, condensation of the hydroxylamines (**5** and **12**) with different aldehydes 2-furaldehyde (**14**), 2-(benzyloxy)acetaldehyde (**15**), took place successfully at 60–90°C (toluene, 4 Å MS) to afford the corresponding unsaturated nitrones. Subsequent ring closure yielded heterocycles (**7a–d** and **16**) as diastereomeric mixtures.

The effect of side chain protection on the regio- and diastereoselectivity of the intramolecular 1,3-cycloaddition has been studied in detail for the *D-gluco* series. The reaction involving the benzyl protected nitron (**13**) proved to be sluggish (toluene, reflux, 1 week) and low-yielding; therefore, it was abandoned. All the cycloaddition products isolated exhibited the same regiochemistry. After 12–48 h, the benzoyl protected nitron was found to afford only 9-oxa-1-azabicyclo[4.2.1]-nonane diastereoisomers **7b–7d** (Figure 2). We were unable to isolate the fourth diastereoisomer (**7a**) as it was produced in a negligible amount. We assume that the alternative 8-oxa-1-azabicyclo[4.2.1]nonane derivatives did not form because of steric hindrance between the furyl chain and oxazepane ring.




FIGURE 2 The structure of diastereoisomers **7a–7d** and **16**.

The effect of different Lewis acid catalysts (ZnCl_2 , MgCl_2 , BF_3OEt_2), solvents (toluene, benzene, dioxane) and time (12–48 h) on yields and diastereomeric ratios have been examined in detail for reactions involving 2-furaldehyde (**14**). It was found that **7b** was the main isomer in all cases (except when dioxane and ZnCl_2 was used). We have ascertained that the best solvent for this reaction is toluene in presence of MgCl_2 as a Lewis acid catalyst (90% yield). The use of a hard Lewis acid catalyst caused elimination of the benzoyl protecting group and, sometimes, decomposition and conversion of the furyl group, too (yields: 17–32%). The stereoselectivity of the reaction did not alter much under the conditions explored. The cycloaddition was found to afford three 9-oxa-1-azabicyclo[4.2.1]nonane diastereomers in a 3:1:1 ratio.

Condensation of 2-(benzyloxy)acetaldehyde (**15**) with benzoyl protected hydroxylamine (**5**) resulted in formation of the single cycloaddition product (**16**), bearing the 9-oxa-1-azabicyclo[4.2.1]nonane skeleton (Figure 2), and the expected intermediate unsaturated nitron was not observed. The reaction took place at 90°C (toluene, 4 Å MS) without the addition of a Lewis acid catalyst. This surprising result implies that we could potentially reduce the number of the possible diastereoisomers generated by the intramolecular 1,3-cycloaddition from 4 to 1. The NMR measurements ^1H , ^{13}C , HSQC, and HMBC suggest the same *trans* arrangement for protons H-6 and H-8 in **16**. We have determined both the configuration of the newly formed chiral centers (*C*-6, *C*-8), and the conformation of the 1,2-oxazepane ring for the all diastereoisomers **7b–7d** and **16**. Finally, the results show that the single isomer of **16** has the same configuration and conformation as the main isomer, **7b**. As a conclusion, various isoxazolidine precursors of chiral azetidine moieties have been successfully synthesised by employing an intramolecular 1,3-dipolar cycloaddition. Out of the three aldehydes examined to date, two of them, aldehydes **14** and **15** have been shown to be suitable candidates for use in this reaction. Our preliminary studies have also found that, when **15** is employed, one diastereoisomer of 9-oxa-1-azabicyclo[4.2.1]nonane is formed exclusively, out of a possible four. The configuration and conformation of the diastereoisomers of the resulting cycloadducts have been examined in detail using NMR studies. The effect of Lewis acid catalysts and solvents on yield and diastereomeric ratios have been investigated.

REFERENCES

1. Bernet, B.; Vasella, A. Carbocyclic compounds from monosaccharides 1. Transformations in the glucose series. *Helv. Chim. Acta* **1979**, *62*, 1990–2016.
2. Garegg, P.J.; Samuelsson, B. Novel reagent system for converting a hydroxy-group into an iodo-group in carbohydrates with inversion of configuration 2. *J. Chem. Soc., Perkin Trans. 1* **1980**, 2866–2869.
3. Skaanderup, P.R.; Hyldtoft, L.; Madsen, R. Zinc-mediated fragmentation of methyl 6-deoxy-6-iodo-hexopyranosides. *Monatsh. Chem.* **2002**, *133*(4), 467–472.
4. Kleban, M.; Kautz, U.; Greul, J.; Hilgers, P.; Kugler, R.; Dong, H.Q.; Jäger, V. Vitamin B-12 catalysis of zinc-mediated 6-deoxy-6-iodopyranoside fragmentation: a mild and convenient preparation of ω -unsaturated hexose derivatives (5-hexenoses). *Synthesis* **2000** (7), 1027–1033.
5. Vasella, A. Synthesis and transformation of isoxazolidine nucleosides. *Helv. Chim. Acta* **1977**, *60*, 426–446.