

This article was downloaded by:

On: 25 January 2011

Access details: Access Details: Free Access

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713597286>

### Degradation of Pyrimidines in *Saccharomyces Kluyveri*: Transamination of $\beta$ -Alanine

K. D. Schnackerz<sup>a</sup>; G. Andersen<sup>a</sup>; D. Dobritzsch<sup>b</sup>; J. Piškur<sup>a</sup>

<sup>a</sup> Cell- and Organism Biology, Lund University, Lund, Sweden <sup>b</sup> Department. of Medical Biochemistry and Biophysics, Karolinska Institutet, Stockholm, Sweden

**To cite this Article** Schnackerz, K. D. , Andersen, G. , Dobritzsch, D. and Piškur, J.(2008) 'Degradation of Pyrimidines in *Saccharomyces Kluyveri*: Transamination of  $\beta$ -Alanine', *Nucleosides, Nucleotides and Nucleic Acids*, 27: 6, 794 — 799

**To link to this Article:** DOI: 10.1080/15257770802145983

**URL:** <http://dx.doi.org/10.1080/15257770802145983>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

## DEGRADATION OF PYRIMIDINES IN *SACCHAROMYCES KLUYVERI*: TRANSAMINATION OF $\beta$ -ALANINE

K. D. Schnackerz,<sup>1</sup> G. Andersen,<sup>1</sup> D. Dobritzsch,<sup>2</sup> and J. Piškur<sup>1</sup>

<sup>1</sup>Cell- and Organism Biology, Lund University, Lund, Sweden

<sup>2</sup>Department of Medical Biochemistry and Biophysics, Karolinska Institutet, Stockholm, Sweden

□ *Beta-alanine is an intermediate in the reductive degradation of uracil. Recently we have identified and characterized the *Saccharomyces kluyveri* PYD4 gene and the corresponding enzyme  $\beta$ -alanine aminotransferase (*SkPyd4p*), highly homologous to eukaryotic  $\gamma$ -aminobutyrate aminotransferase (GABA-AT). *S. kluyveri* has two aminotransferases, GABA aminotransferase (*SkUga1p*) with 80% and *SkPyd4p* with 55% identity to *S. cerevisiae* GABA-AT. *SkPyd4p* is a typical pyridoxal phosphate-dependent aminotransferase, specific for  $\alpha$ -ketoglutarate ( $\alpha$ KG),  $\beta$ -alanine (BAL) and  $\gamma$ -aminobutyrate (GABA), showing a ping-pong kinetic mechanism involving two half-reactions and substrate inhibition. *SkUga1p* accepts only  $\alpha$ KG and GABA but not BAL, thus only *SkPyd4p* belongs to the uracil degradative pathway.*

**Keywords** Aminotransferase;  $\beta$ -alanine;  $\gamma$ -aminobutyrate; gene duplication; *Saccharomyces kluyveri*

### INTRODUCTION

In biological systems,  $\beta$ -alanine (BAL) plays a major role as a precursor of pantothenic acid,<sup>[1]</sup> a constituent of dipeptides such as anserine and carnosine in the brain<sup>[2]</sup> and as an intermediate in the catabolic pathway of pyrimidines.  $\beta$ -Alanine is a structural analog of  $\gamma$ -aminobutyrate (GABA) and glycine, two major inhibitory neurotransmitters.<sup>[3]</sup> Moreover, BAL itself has been suggested to be involved in synaptic transmission. Recent studies revealed that BAL activates the same receptors on the membranes of rat sacral dorsal commissural neurons as glycine.<sup>[4]</sup> In mammalian systems, uracil is degraded to BAL in three consecutive enzymatic steps by dihydropyrimidine dehydrogenase, dihydropyrimidine amidohydrolase and  $\beta$ -alanine synthase. In contrast, microorganisms produce BAL mostly by direct

decarboxylation of L-aspartate.<sup>[5]</sup> Bacteria and plants can also degrade spermidine to BAL.<sup>[6]</sup>

Mammals degrade BAL to malonate semialdehyde (MSA) using either BAL: $\alpha$ -ketoglutarate aminotransferase (BAL-AT) or D-3-aminoisobutyrate: pyruvate aminotransferase (DAIB-AT). The produced MSA is subsequently metabolized by semialdehyde dehydrogenase to acetyl-CoA.<sup>[7]</sup> In humans, BAL-AT and GABA aminotransferase activity resides within a single enzyme.<sup>[8]</sup>

Even though *Saccharomyces cerevisiae* can synthesize BAL from polyamines, it cannot use BAL as sole nitrogen source.<sup>[9]</sup> GABA, in contrast, is readily used as sole nitrogen source by the action of GABA-AT and succinic semialdehyde dehydrogenase (SSADH), encoded by *UGA1* and *UGA2*, respectively.<sup>[10]</sup> Several yeast species, like *Saccharomyces kluyveri* and *Schizosaccharomyces pombe* can utilize BAL as sole nitrogen source.<sup>[11]</sup>

In this study, we present the identification and characterization of two *S. kluyveri* enzymes involved in transamination of BAL and GABA, encoded by the *PYD4* and *UGA1* genes.

## MATERIALS AND METHODS

All chemicals and enzymes were purchased from Sigma (St. Louis, MO, USA). Yeast nitrogen base without amino acids and ammonium sulfate were obtained from Difco (BD Biosciences, San Jose, CA, USA) and chelating Sepharose fast flow from Amersham Biosciences (Piscataway, NJ, USA). Oligos were purchased from DNA Technology (Aarhus, Denmark).

### Strains and Media

The yeast and bacterial strains used in this study were grown in YPD medium (1% yeast extract, 2% bactopectone, 2% glucose) or synthetic defined medium (1% succinic acid, 0.6% sodium hydroxide, 2% glucose, 0.67% yeast nitrogen base without amino acids and ammonium sulfate) supplemented with different nitrogen sources (0.5% ammonium sulfate, 0.1% for all other types). For solid medium (plates) 2% agar was added. For plasmid rescue, *E. coli* strain XL1-Blue was used, and for cloning TOP10 and for overexpression of fusion proteins, BL21 Star. All bacteria were grown at 37°C in LB medium supplemented with ampicillin ( $100 \text{ mg} \times \text{L}^{-1}$ ).

### Mutagenesis

*S. kluyveri* mutants were generated from strains Y156 and 159 with ethyl methanesulfonate.<sup>[11]</sup> Mutagenized cells were plated and grown for 2–3 days at 25°C. The plates were replicated onto new plates containing ammonium sulfate (control) or dihydrouracil (DHU) as sole nitrogen

source. After 5–7 days at 25°C, colonies were selected based on their inability to grow on DHU plates. Putative mutants were tested for growth on  $\beta$ -ureidopropionate (BUP), with BAL as sole nitrogen source.

### **Cloning, Expression, and Purification of SkPyd4p**

The cloning, expression and purification are described in detail elsewhere.<sup>[12]</sup>

Clones of succinic semialdehyde dehydrogenase (SSADH) and malonate semialdehyde decarboxylase (MSADC) were kindly provided by M. D. Toney (University of California at Davis) and C. P. Whitman (University of Texas, Austin) and the proteins obtained as previously described.<sup>[13,14]</sup>

### **Spectrophotometric Measurements**

All measurements were performed on a Varian Cary 3 UV/Visible spectrophotometer equipped with a temperature controlled automatic multi-cell changer.

### **Enzyme Assays**

In a continuous assay, the production of MSA by BAL-AT is coupled with MSADC to produce acetaldehyde, which in a NADH-dependent alcohol dehydrogenase reaction generates NAD<sup>+</sup> and ethanol. The MSADC-generated acetaldehyde was monitored by observing the NADH depletion at 340 nm.<sup>[12]</sup> In a continuous assay, succinic semialdehyde (SSA) produced by Ugalp and Pyd4p was converted by the NADP<sup>+</sup>-dependent SSADH to NADPH and succinate. The absorbance change at 340 nm was monitored.<sup>[12]</sup>

### **Data Analysis**

Kinetic data analysis was performed as described previously.<sup>[12]</sup>

## **RESULTS**

### **Identification of the *PYD4* Gene Encoding a $\beta$ -Alanine Aminotransferase**

While screening mutagenized *S. kluyveri* cells grown on DHU, a mutant, Y947, was isolated which could not use DHU, BUP ( $\beta$ -ureidopropionate) or BAL as sole nitrogen source. This mutant behaved like a double mutant, since both BUP and BAL can serve as a direct nitrogen source via the action of  $\beta$ -alanine synthase and BAL-AT. However, complementation with the genomic library yielded plasmid P733, which supported growth on all three

nitrogen sources (DHU, BUP, and BAL). An open reading frame, termed *PYD4* (DQ512721), encoding a protein with 55% identity at the protein level to *S. cerevisiae* *UGA1* gene product, GABA-AT, was identified. A BLAST search of *ScUGA1* in the *S. kluyveri* genome revealed another homologous gene, termed *SkUGA1* (DQ512722). Its gene product has 80% identity to *ScUga1p*, and 57% to *SkPyd4p*.

### Sequence Analysis of the *PYD4* Gene

Phylogenetic analysis of the homologous proteins from yeast shows that *PYD4* is only found in two other sequenced yeasts (*Debaryomyces hansenii* and *Candida albicans*). Both yeasts also contain a putative *UGA1* gene. A search for  $\beta$ -alanine synthase (encoded by *PYD3*) and dihydropyrimidine amidohydrolase (encoded by *PYD2*) revealed that *S. kluyveri* is the only yeast carrying all three *PYD* genes. *C. albicans* and *D. hansenii* lack *PYD2*, and *Kluyveromyces lactis* has *PYD2* and *PYD3*, but lacks *PYD4*. *S. cerevisiae* lacks *PYD2*, *PYD3*, and *PYD4*.

### Purification and Properties of *SkPyd4p* and *SkUga1p*

The purification of the His-tagged *SkPyd4p* and *SkUga1p* was accomplished after disruption of the respective cells in three steps using streptomycin treatment, affinity chromatography on nickel-chelating Sepharose and size exclusion gel filtration on Sepharose S-12. From 1 liter of cell culture about 20 mg of pure protein could be obtained. The molecular weight of the subunits of both proteins was found to be about 56-57 kDa based on SDS/PAGE gel. Purified *SkPyd4p* and *SkUga1p* had native molecular masses of 111 (dimers) and 135/240 kDa (dimers/tetramers), respectively. The spectra of purified *SkPyd4p* and *SkUga1p* show two maxima at 340 and 410 nm, indicating the equilibrium of the ketoenamine (410 nm) and enolimine (340 nm) tautomers which can be shifted by changing the pH of the protein solution.

### Initial Velocity Studies

In the initial reaction of *SkPyd4p*,  $\alpha$ KG and BAL (or GABA) are converted to glutamate and MSA (or SSA), respectively. In a second reaction, either the amount of MSA or SSA was determined. The substrate specificity of *SkPyd4p* was tested and it could be shown that *SkPyd4p* accepts both BAL and GABA as amino donors but with a preference for BAL. In contrast, *SkUga1p* uses only GABA as substrate not BAL.

Both *SkPyd4p* and *SkUga1p* were subjected to a full kinetic analysis. Since *SkUga1p* shows no activity towards BAL, only *SkPyd4p* was analyzed with both substrates. When the initial velocity was determined at various  $\alpha$ KG


concentrations and several fixed levels of BAL, a pattern composed of parallel lines is seen at low reactant concentrations in a double reciprocal plot. At high  $\alpha$ KG concentrations, the linear lines are deflected upwards, indicating substrate inhibition. Values of  $K_{M,BAL}$ ,  $K_{M,GABA}$  and  $K_{M,\alpha KG}$  with  $8.2 \pm 0.4$ ,  $1.8 \pm 0.2$ ,  $2.9 \pm 0.2$  mM, respectively, were determined for  $^{Sk}Pyd4p$ . Since the maximal velocity for  $^{Sk}Pyd4p$  (7.5 units/mg) is more than eight times higher for BAL,  $^{Sk}Pyd4p$  can be considered a BAL-AT rather than a GABA-AT.  $^{Sk}Uga1p$  exhibits  $K_{M,GABA}$  and  $K_{M,\alpha KG}$  values of  $3.2 \pm 0.2$  and  $0.22 \pm 0.01$  mM, respectively. The obtained kinetic behaviour is consistent with a ping-pong kinetic mechanism.

## DISCUSSION

Mammals have only one gene encoding a GABA-AT which exhibits both BAL-AT and GABA-AT activity. *S. cerevisiae* has a single gene encoding GABA-AT ( $^{Sc}UGA1$ ), while *S. kluyveri* has two  $^{Sc}UGA1$ -like genes. Similarly, the genomes of *C. albicans* and *D. hansenii* each contain an  $UGA1$ -like gene. This study identifies that one of the two genes in *S. kluyveri*,  $PYD4$ , encodes a BAL-AT involved in BAL degradation. Apparently, the original  $PYD4/UGA1$  gene was duplicated in one yeast lineage. Later on, one of the duplicated genes was lost in some descendant lineages, such as *S. cerevisiae*. The time point for the gene duplication is estimated to be  $\sim 200$  million years ago, when *Yarrowia lipolytica* split from the *Saccharomyces/Candida/Debaryomyces* lineages.<sup>[15]</sup>

The pattern of an array of parallel lines for different fixed concentrations of the second substrate is consistent with a ping-pong kinetic mechanism. All aminotransferases adhere to such a mechanism, where the keto substrates exert substrate inhibition. Both  $^{Sk}Pyd4p$  and GABA-AT from *E. coli* show comparable substrate inhibition for  $\alpha$ KG with inhibition constants of  $28.3 \pm 6.7$  and  $10.2 \pm 2.3$  mM,<sup>[14]</sup> respectively. Both  $^{Sk}Pyd4p$  and  $^{Sk}Uga1p$  show a reaction mechanism involving two half-reactions.

Our study shows that  $PYD4/^{Sk}Pyd4p$  is a part of the reductive pathway of pyrimidine nucleic acid precursors (Figure 1).



**FIGURE 1** Reductive degradation of pyrimidine nucleic acid precursors. Uracil is degraded via dihydrouracil, N-carbamyl- $\beta$ -alanine,  $\beta$ -alanine to malonate semialdehyde by dihydropyrimidine dehydrogenase (1, encoded by  $PYD1$ ), dihydropyrimidine amidohydrolase (2, encoded by  $PYD2$ ),  $\beta$ -alanine synthase (3, encoded by  $PYD3$ ), and  $\beta$ -alanine aminotransferase (4, encoded by  $PYD4$ ), respectively.<sup>[16]</sup>

## REFERENCES

1. Cronan, J.E.; Little, K.J.; Jackowski, S. Genetic and biochemical analyses of pantothenate biosynthesis in *Escherichia coli* and *Salmonella typhimurium*. *J. Bacteriol.* **1982**, 149, 916–922.
2. Bauer, K.; Salnikow, J.; deVitry, F.; Tixier-Vidal, S.; Kleinkauf, H. Characterization and biosynthesis of omega-aminoacyl amino acids from rat brain and the C-6 glioma cell line. *J. Biol. Chem.* **1979**, 254, 6402–6407.
3. Enna, S.J.; Gallagher, J.P. Biochemical and electrophysiological characteristics of mammalian GABA receptors. *Int. Rev. Neurobiol.* **1983**, 24, 181–212.
4. Wang, D.S.; Zhu, H.L.; Li, J.S. Beta-alanine acts on glycine receptors in the rat dorsal commissural neurons. *Int. J. Neurosci.* **2003**, 113, 293–305.
5. Williamson, J.M.; Brown, G.M. Purification and properties of L-aspartate-alpha-decarboxylase, an enzyme that catalyzes the formation of beta-alanine in *Escherichia coli*. *J. Biol. Chem.* **1979**, 254, 8074–8082.
6. Large, J.P. Enzymes and pathways of polyamine breakdown in microorganisms. *FEMS Microbiol. Rev.* **1992**, 88, 249–262.
7. Goodwin, G.W.; Rougraff, P.M.; Davis, E.J.; Harris, R.A. Purification and characterization of methylmalonate semialdehyde dehydrogenase from rat liver. Identity to malonate semialdehyde dehydrogenase. *J. Biol. Chem.* **1989**, 264, 14965–14971.
8. Schor, D.S.; Struys, E.A.; Hogema, B.M.; Gibson, K.M.; Jacobs, C. Development of a stable-isotope dilution assay for gamma-aminobutyric acid (GABA) transaminase in isolated leukocytes and evidence that GABA and beta-alanine transaminase are identical. *Clin. Chem.* **2001**, 47, 525–531.
9. Ramos, F.; el Guezzer, M.; Grenson, M.; Wiame, J.M. Mutations affecting the enzymes involved in the utilization of 4-aminobutyric acid as nitrogen source by yeast *Saccharomyces cerevisiae*. *Eur. J. Biochem.* **1985**, 149, 401–404.
10. LaRue, T.A.; Spencer, J.F. The utilization of purines and pyrimidines by yeasts. *Can. J. Microbiol.* **1968**, 14, 19–86.
11. Gojkovic, Z.; Paracchini, S.; Piškur, J. A new model organism for studying the catabolism of pyrimidines and purines. *Adv. Exper. Med. Biol.* **1998**, 431, 475–479.
12. Andersen, G.; Andersen, B.; Dobritzsch, D.; Schnackerz, K.D.; Piškur, J. A gene duplication led to specialized  $\gamma$ -aminobutyrate and  $\beta$ -alanine aminotransferases in yeast. *FEBS J.* **2007**, 274, 1804–1817.
13. Poelarends, G.J.; Johnson, W.H.; Murzin, A.G.; Whitman, C.P. Mechanistic characterization of a bacterial malonate semialdehyde decarboxylase: identification of a new activity on the tautomerase superfamily. *J. Biol. Chem.* **2003**, 278, 48674–48683.
14. Liu, W.; Peterson, P.E.; Carter, R.J.; Zhou, X.; Langston, J.A.; Fisher, A.J.; Toney, M.D. Crystal structures of unbound and aminooxyacetate *Escherichia coli*  $\gamma$ -aminobutyrate aminotransferase. *Biochemistry* **2004**, 43, 10896–10905.
15. Piškur, J.; Rozpedowska, E.; Polakova, S.; Merico, A.; Compagno, C. How did *Saccharomyces* evolve to become a good brewer? *Trends Genet.* **2006**, 22, 183–186.
16. Piškur, J.; Schnackerz, K.D.; Andersen, G.; Björnberg, O. Comparative genomics reveals novel biochemical pathways. *Trends Genet.* **2007**, 23, 369–372.