

Stereochemistry of the Cyclization-Rearrangement of (+)-Copalyl Diphosphate to (–)-Abietadiene Catalyzed by Recombinant Abietadiene Synthase from *Abies grandis*


Matthew M. Ravn,[†] Robert M. Coates,^{*,†} Janice E. Flory,[‡] Reuben J. Peters,[‡] and Rodney Croteau[‡]

Department of Chemistry, University of Illinois, 600 S. Mathews Avenue, Urbana, Illinois 61801, and Institute of Biological Chemistry, Washington State University, Pullman, Washington 99164

r-coates@uiuc.edu

Received November 6, 1999

ABSTRACT



Syntheses and enzymatic cyclizations of 8 α -hydroxy-17-nor copalyl diphosphate (8a), (15*R*)-[15-²H₁] 8b, and (15*R*,17*E*)-[15-³H₁,17-²H₁] copalyl diphosphate ([²H,³H] 2) catalyzed by recombinant abietadiene synthase (rAS) gave 17-nor manoyl oxide (9a), (16*E*)-[16-²H₁] 9b, and (15*S*,16*R*)-[16-²H₁,16-³H₁] abietadiene ([²H₁,³H₁] 4), respectively. These and other results indicate that conversion of CPP (2) to abietadiene (4) occurs by anti S_N' cyclization to a sandaracopimar-15-en-8-yl carbocation intermediate (13⁺, 13 β -methyl) followed by hydrogen transfer and methyl migration suprafacially on the *si* face of the vinyl group.

The abietanes comprise a large family of perhydrophenanthrene-type diterpenes characterized by an isopropyl group, or its functionalized equivalent, at C13.¹ Familiar examples are abietic acid (5) and the isomeric levopimaric, neoabietic, and palustric acids, which are major diterpene constituents of conifer oleoresin.² This defensive secretion is produced by pines, firs, spruces, and other conifers as a primary response to wounds caused by physical injuries, insects, large

herbivores, and microbial diseases.³ The derivation of the abietane carbon skeleton from the predicted head-to-tail connectivity of isoprene units led L. Ruzicka to propose that the aberrant isopropyl group might arise by proton-induced rearrangement of a regular isoprenoid precursor, such as pimaric acid.⁴ The acid-induced conversion of pimaric and isopimaric acids to abietic acid affords chemical precedent for this biogenetic relationship.⁵

The native enzymes responsible for the cyclization of (*E,E,E*)-geranylgeranyl diphosphate (1) to (–)-abietadiene (4) (Scheme 1) were isolated in partially purified form from the stems of both wound-induced grand fir (*Abies grandis*)

* Tel: (217) 333-4280. FAX: (217) 244-8068.

[†] University of Illinois.

[‡] Washington State University.

(1) *CRC Handbook of Terpenoids: Diterpenoids Vol I–IV*; Dev, S., Misra, R., Eds.; CRC Press: Boca Raton, FL, 1985–1986.


(2) Soltes, J.; Zinkel, D. F. In *Naval Stores: Production, Chemistry and Utilization*; Zinkel, D. F., Russel, J., Eds.; Pulp Chemicals Association: New York, 1989; Chapter 9.

(3) Phillips, M. A.; Croteau, R. *Trends Plant Sci.* **1999**, 4, 184.

(4) Ruzicka, L. *Experientia* **1953**, IX, 357.

(5) Wenkert, E.; Chamberlin, J. W. *J. Am. Chem. Soc.* **1959**, 81, 688.

Scheme 1



and lodgepole pine (*Pinus contorta*) saplings.⁶ Cytochrome P450 mixed function oxygenases and a dehydrogenase from the same sources, which carry out the sequential oxidation of the C18 methyl group of **4** to produce abietic acid, have also been characterized.⁷ The gene encoding abietadiene synthase (AS) from grand fir has been cloned, and the cDNA was functionally expressed as a single polypeptide bearing a putative *N*-terminal plastidial targeting sequence.⁸ Recombinant AS (rAS) was heterologously expressed in *E. coli* as a pseudomature enzyme without the *N*-terminal 84 amino acids⁹ and used in crude bacterial extracts after lysis and clarification (at 0.1 g wet cell weight/mL buffer). Although the resulting recombinant AS catalyzed the divalent metal ion-dependent cyclizations of both **1** and the predicted bicyclic intermediate (+)-copalyl diphosphate (**2**) to **4**,^{9,10} none of the four plausible pimaradiene intermediates was converted into the rearranged diterpene to an appreciable extent under the same assay conditions.¹⁰ Recent deuterium labeling experiments revealed that the cyclization-rearrangement of (+)-CPP to abietadiene occurs with stereospecific “intramolecular” transfer of a hydrogen atom from the 17-*pro-E* position of **2** to the terminal carbon of the side chain, which becomes the *pro-S* methyl of the isopropyl group.^{9,10} In this communication, we present evidence indicating a 13 β -methyl group in the presumed pimaradiene (**3b**) or pimaranyl carbocation intermediate and a suprafacial relationship between the hydrogen reincorporated at C16 and the subsequent methyl migration to C15.¹¹

(6) LaFever, R. E.; Vogel, B. S.; Croteau, R. *Arch. Biochem. Biophys.* **1994**, *313*, 139.

(7) Funk, C.; Croteau, R. *Arch. Biochem. Biophys.* **1994**, *308*, 258.

(8) Vogel, B. S.; Wildung, M. R.; Vogel, G.; Croteau, R. *J. Biol. Chem.* **1996**, *271*, 23262.


(9) Ravn, M. M.; Coates, R. M.; Jetter, R.; Croteau, R. *Chem. Commun.* **1998**, 21.

(10) Peters, R. J.; Flory, J.; Jetter, R.; Croteau, R., manuscript in preparation.

(11) The occurrence of an analogous proton transfer to form the *pro-S* methyl group of a presumptive abietadiene intermediate was inferred from the labeling pattern of ginkgolide A biosynthesized from [6,6,6-²H₃] mevalonate in *Ginkgo biloba* cell cultures: (a) Schwarz, M.; Arigoni, D. In *Isoprenoids Including Carotenoids and Steroids*; Cane, D. E., Ed.; Vol. 2 in *Comprehensive Natural Products Chemistry*; Barton, D., Nakanishi, K., Meth-Cohn, O., Eds.; Elsevier: Oxford, 1999; Chapter 14. (b) Schwarz, M. K. Dissertation 10951, ETH Zürich, Switzerland, 1994.

The coupled cyclization-rearrangement of **2** was intercepted by use of an oxa analogue (**8a**) bearing an 8 α -hydroxyl function in place of the exocyclic methylene group (Scheme 2). 8-Oxo-17-nor copalol (**6a**), obtained in three

Scheme 2



a. MnO₂, hexane. b. (*R*)-Alpine borane, THF, 0 °C; H₂O₂, OH⁻. c. Li/NH₃, Et₂O, EtOH, -33 °C. d. Cl(O)P(OEt)₂, py, CH₂Cl₂, 0 °C. e. (Bu₄N)₃HP₂O₇·(H₂O)₃, CH₃CN, 4d.

steps from methyl copalate,¹² underwent reduction with Li/NH₃ (EtOH, ether, -33 °C) to 8 α -hydroxy-17-nor copalol (**7a**, 83%).^{13,14} Conversion to the corresponding allylic diphosphate **8a** was accomplished in two steps by phosphorylation [(EtO)₂P(O)Cl, py, CH₂Cl₂, 0 °C, 3 h],¹⁵ displacement with pyrophosphate anion [(Bu₄N)₃HP₂O₇, CH₃CN, rt, 3.7 d], and purification according to the Poulter methods (66%).^{14,16} Incubation of **8a** with rAS^{8,10,17} effected clean cyclization to 17-nor manoyl oxide **9a** (yield, ca. 150 μ g, 46% by ¹H NMR estimate), the 13 β -methyl configuration of which was established by NOE measurements.¹⁴ Acid-catalyzed dehydration (*p*TsOH·H₂O, ether, rt, 1.5 h) of **7a** afforded a 1.5:1 mixture of **9a** and its 13 α -methyl isomer (76%). The formation of 17-nor manoyl oxide in the enzymatic cyclization implicates a β -methyl group at C13

(12) (a) catalytic OsO₄, NMO, acetone; 3:1 mixture of 8,17- and 13,14-diols (78%); (b) LiAlH₄ (or LiAlD₄ for **6b**), ether; (c) NaIO₄, aqueous acetone (60% for two steps).

(13) All new compounds were characterized at high purity by appropriate ¹H NMR, ¹³C NMR, and IR spectra. Purity \geq 95% was established by satisfactory combustion analysis and/or inspection of ¹H and ¹³C NMR spectra. Detailed experimental procedures and characterization data are provided in the Supporting Information. Key data are presented in the text or in footnote 14.

(14) Key spectral and physical data for selected compounds: **7a**, mp 85–86 °C, ¹H NMR δ 3.41 (td, *J* = 10.6, 4.9 Hz, 1H, H₈); **8a**, ³¹P NMR δ -7.02, -9.72 (2d, *J* = 22 Hz); **9a**, NOE irradiat at 3.52 obs 0.74 (C20 CH₃, 7.3%), 1.21 (C17 CH₃, 6.7%); **9b**, ¹H NMR δ 5.37 and 6.04 (2d, *J* = 17.4 Hz, 1H each, CH=CHD).

(15) Ravn, M. M.; Jin, A. Q.; Coates, R. M. *Eur. J. Org. Chem.*, in press.


(16) For leading references, see: (a) Davisson, V. J.; Woodside, A. B.; Neal, T. R.; Stremmer, K. E.; Muehlbacher, M.; Poulter, C. D. *J. Org. Chem.* **1986**, *51*, 4768. (b) Woodside, A. B.; Huang, Z.; Poulter, C. D. *Org. Synth., Coll. Vol. VIII*, **1993**, 616.

(17) Incubation conditions:^{8,10} 30 mM HEPES, pH 7.2, 5.0 mM dithiothreitol, 7.5 mM MgCl₂, 20 μ M MnCl₂, 5% (v/v) glycerol, 80 μ M **8a** (0.40 μ mol total), 3 \times 1 mL rAS cell lysate; 31 °C, 16 h. Purification by pipet chromatography over MgSO₄ and silica gel; yield, ca. 150 μ g (46%) by ¹H NMR estimate.

of the tricyclic intermediate corresponding to sandaracopimaradiene (**3b**) which in fact is a consistent, albeit very minor, byproduct (ca 2%) that accompanies abietadiene in the extractable products from incubations of **1** with rAS.^{8,10}

The stereochemistry of the enzyme-catalyzed S_N' cyclization was determined by deuterium labeling at C15 (Scheme 2). Allylic oxidation (MnO_2 , hexane, rt) of **6b** followed by *re*-selective hydride reduction with (*R*)-Alpine borane¹⁸ (THF , H_2O_2 , NaOH)¹⁹ and Li/NH_3 reduction gave **7b**. The 15*S* configuration and stereohomogeneity (98% de) of the intermediate keto alcohol **6c** were confirmed by ^1H NMR analysis of the corresponding camphanate ester.²⁰ Conversion to **8b** by diphosphate displacement as described above for unlabeled **7a** was assumed to proceed with complete inversion of configuration, as established for (*S*)-[1- $^2\text{H}_1$] geraniol.^{15,20} rAS-catalyzed cyclization of **8b** gave rise to **9b**, the 16*E* stereochemistry of which was established by the large coupling ($J = 17$ Hz) between the two vinyl protons.¹⁴ Hence, the stereochemistry of this enzyme-catalyzed S_N' cyclization, and by inference that of **2**, is anti, in agreement with similar S_N' cyclizations that occur in diterpene biosynthesis.²¹

The stereochemistry of the hydrogen reincorporation at C16 during the rearrangement that generates the isopropyl group was elucidated by double isotope labeling (Schemes 3 and 4). (*15R,17E*)-[15- $^3\text{H}_1$,17- $^3\text{H}_1$] **2** was synthesized from




methyl copalate as shown in Scheme 3. Bromination of the exocyclic double bond followed by diisobutylaluminum

(18) (a) Midland, M. M. *Chem. Rev.* **1989**, 89, 1553. (b) Midland, M. M.; Greer, S.; Tramantano, A.; Zderic, S. A. *J. Am. Chem. Soc.* **1979**, 101, 2352. (c) Prabhakaran, P. C.; Gould, S. J.; Orr, G. R.; Coward, J. K. *J. Am. Chem. Soc.* **1988**, 110, 5779.

(19) (a) Midland, M. M.; Graham, R. S. *Org. Synth., Coll. Vol. VIII*, **1990**, 402. (b) Krishnamurthy, S.; Brown, H. C. *J. Org. Chem.* **1977**, 42, 1197.

(20) (a) Gerlach, H.; Zagalak, B. *J. Chem. Soc., Chem. Commun.* **1973**, 274. (b) Gerlach, H.; Kappes, D.; Boeckman, R. K.; Maw, G. N. *Org. Synth., Coll. Vol. IX*, **1998**, 151.

Scheme 4



hydride reduction and dehydrobromination and afforded (17*E*)-17-bromocopalol (**11**). The corresponding tetrahydropyranyl ether was lithiated, deuterated, and hydrolyzed to (17*E*)-[17- $^2\text{H}_1$] copalol (**12**). A four-step reaction sequence consisting of MnO_2 oxidation, NaBH_3T reduction, MnO_2 oxidation, and (*R*)-Alpine borane reduction gave (15*S*,17*E*)-[15- $^2\text{H}_1$,17- $^3\text{H}_1$] copalol ([$^2\text{H},^3\text{H}$] **12**).^{18,19} Conversion to the diethyl phosphate followed by displacement with $(\text{Bu}_4\text{N})_3\text{P}_2\text{O}_7\text{H}$ in CH_3CN with inversion of configuration¹⁵ provided (15*R*,17*E*)-[17- $^3\text{H}_1$,15- $^2\text{H}_1$] **2** (0.13 mCi, 40% radiochemical yield, 21.8 mCi/mmol).

Incubation with rAS¹⁷ and addition of carrier afforded [16- $^2\text{H}_1$,16- $^3\text{H}_1$] abietadiene (**4**) (1.8 mg, 1.12 $\mu\text{Ci}/\mu\text{mol}$, 1.57×10^8 dpm), which was subjected to Kuhn–Roth oxidation (aq H_2CrO_4 , reflux, 16 h)²² (Scheme 4). The resulting [$^2\text{H},^3\text{H}$] acetic acid (3.1×10^7 dpm, 20% radiochemical yield, 90% radiochemical purity) was isolated by steam distillation and analyzed by enzymatic chirality assay with malate synthase and fumarase.²³ The *F* values observed, 74.1 ± 2.1 and 74.7 ± 1.6 , establish *R* stereochemistry for the chiral acetate, formation of (15*S*,16*R*)-[16- $^2\text{H}_1$,16- $^3\text{H}_1$] abietadiene (**4**) in the enzymatic reaction, and overall retention of configuration in the replacement of diphosphate of [$^2\text{H},^3\text{H}$] **2** with deuterium.

Because an anti S_N' cyclization of doubly labeled **2** would generate a vinyl group bearing tritium in the 16*E* position of the pimarenyl intermediate (e.g., **13**⁺), the deuterium atom from C17 and the methyl group from C13 both migrate to the *si,si* face of the vinyl group of **3b**, i.e., a formal suprafacial relationship.²⁴ This stereochemical finding is at variance with either a concerted pericyclic mechanism

(21) (a) MacMillan, J.; Beale, M. H. In *Isoprenoids Including Carotenoids and Steroids*; Cane, D. E., Ed.; Vol. 2 in *Comprehensive Natural Products Chemistry*; Barton, D., Nakanishi, K., Meth-Cohn, O., Eds.; Elsevier: Oxford, 1999; Chapter 8 (b) Cane, D. E. *Tetrahedron* **1980**, 36, 1109.

(22) Phillips, G. T.; Clifford, K. H. *Eur. J. Biochem.* **1976**, 61, 271.

(23) (a) Floss, H. G.; Tsai, M. D. *Adv. Enzymol.* **1982**, 50, 243. (b) Unpublished protocols and calibration standards of (*R*)- and (*S*)-[$^2\text{H},^3\text{H}$] acetate were kindly provided by H. G. Floss and S. Lee, University of Washington, Seattle, WA. The *F* values obtained for the chiral acetate standards at the UI were as follows: *R*, 78.0 ± 3.6 ; *S*, 28.1 ± 3.6 .

predicted to be antarafacial on the basis of orbital symmetry considerations^{11b} or a synchronous proton addition to C16 and antiperiplanar methyl migration to C15. Consequently a secondary sandaracopimaren-15-yl carbocation (**14**⁺)/diphosphate anion pair, or alternatively a covalently bonded pimarenyl–enzyme adduct,^{25,26} is implicated prior to the methyl group rearrangement and proton elimination steps leading to abietadiene.

The occurrence of bona fide secondary carbocation intermediates in terpene synthase-catalyzed cyclizations and rearrangements is rare, reflecting the higher energy of these species. Examples are found in mechanistic schemes leading to monoterpenes (bornyl diphosphate and fenchol synthases),^{27a} sesquiterpenes (pentalene and trichodiene synthases),^{27b} tetracyclic diterpenes (e.g., *ent*-kaurene synthase),^{21a} and polycyclic triterpenes (squalene and oxido-squalene synthases).^{27c,d} However, in all of these cases the 15 kcal/mol thermodynamic energy barrier between the secondary ion²⁸ and a preceding tertiary carbocation would be offset by the substantial enthalpic gains associated with cyclizations into C=C double bonds ($\Delta H_{BE} \approx -20$ kcal/mol) or with relief of ring strain (bicyclo[3.1.1]heptane/bicyclo[2.2.1]-heptane, $\Delta\Delta H_{SE} = -19$ kcal/mol),²⁹ together with the stabilization arising from charge delocalization (bridged ions) and concerted bond-forming reactions.

(24) (a) The relatively efficient enzymatic cyclization of **8a** and the unproductive binding of its 8 β -OH isomer ($K_i = 0.18 \mu\text{M}$),^{24b} as well as precedent,²¹ suggest that the S_N' cyclization probably takes place on the *re, re*(α) face of C17 to generate **13**⁺ with a 14 α deuterium substituent. (b) Ravn, M. M.; Coates, R. M.; Peters, R. J.; Croteau, R., unpublished results.

(25) Cornforth, J. W. *Angew. Chem., Int. Ed. Engl.* **1968**, 7, 903.

(26) Incubations of (13*S*,15*S*)-isopimar-8(14)-en-15-yl diphosphate and a mixture of (13*S*,15*S*) and (13*S*,15*R*) isomers with rAS did not produce detectable quantities of abietadiene. Thus, internal return of OPP[−] to form a freely exchangeable covalent diphosphate intermediate appears to be excluded.

(27) (a) Wise, M. L.; Croteau, R. In *Isoprenoids Including Carotenoids and Steroids*; Cane, D. E., Ed.; Vol. 2 in *Comprehensive Natural Products Chemistry*; Barton, D., Nakanishi, K., Meth-Cohn, O., Eds.; Elsevier: Oxford, 1999; Chapter 5. (b) Cane, D. E. In *Isoprenoids Including Carotenoids and Steroids*; Cane, D. E., Ed.; Vol. 2 in *Comprehensive Natural Products Chemistry*; Barton, D., Nakanishi, K., Meth-Cohn, O., Eds.; Elsevier: Oxford, 1999; Chapter 6. (c) Abe, I.; Prestwich, G. D. In *Isoprenoids Including Carotenoids and Steroids*; Cane, D. E., Ed.; Vol. 2 in *Comprehensive Natural Products Chemistry*; Barton, D., Nakanishi, K., Meth-Cohn, O., Eds.; Elsevier: Oxford, 1999; Chapter 10. [d] Poralla, K. In *Isoprenoids Including Carotenoids and Steroids*; Cane, D. E., Ed.; Vol. 2 in *Comprehensive Natural Products Chemistry*; Barton, D., Nakanishi, K., Meth-Cohn, O., Eds.; Elsevier: Oxford, 1999; Chapter 11.

(28) (a) Bittner, E. W.; Arnett, E. M.; Saunders, M. J. *Am. Chem. Soc.* **1976**, 98, 2724. (b) Arnett, E. M.; Petro, C. J. *Am. Chem. Soc.* **1978**, 100, 5408.

(29) Liebman, J. F.; Greenberg, A. *Chem. Rev.* **1976**, 76, 320.

The thermodynamically uphill isomerization of the tertiary pimarenyl ion **13**⁺ to the apparently localized secondary ion **14**⁺ in the absence of an exothermic counterbalance or the possibility of a lower energy concerted mechanism is unprecedented. Although the 15 kcal/mol thermodynamic barrier between tertiary and secondary carbenium ions²⁸ should be surmountable at room temperature, it nevertheless seems likely that AS participates overtly in catalyzing the process and lowering the thermodynamic deficit. Some important factors to consider in the enzyme-induced destabilization of **13**⁺/OPP[−] and/or stabilization of **14**⁺/OPP[−] carbocation-diphosphate anions are ion pair distances and forces,³⁰ homoallyl interaction in **14**⁺, π -complexation with the aromatic rings of amino acid side chains at the active site,³¹ interactions with proximal peptide carbonyl dipoles or nucleophilic heteratoms,³² and electrostatic field gradients. Interestingly, binding of a sandaracopimarenyl amine, mimicking the secondary carbocation intermediate, is greatly enhanced (~ 1000 -fold) by the addition of inorganic diphosphate,^{24b} suggesting that AS utilizes the diphosphate anion to stabilize the secondary carbocation intermediate. The X-ray crystallographic structure of rAS and its complexes with pimarenyl ion mimic inhibitors may reveal what specific interactions and mechanisms are involved in the remarkable pimarenyl⁺–abietadienyl⁺ rearrangement.

Acknowledgment. We thank the National Institutes of Health for research grants GM 13956 (R.M.C.) and GM 31354 (R.B.C.) and Drs. H.G. Floss and S. Lee for authentic standards of (*R*)- and (*S*)-[²H₁,³H₁] acetates and assistance with the enzymatic chirality assays. R.M.C. thanks D. Arigoni and F.-G. Klärner for helpful discussions. R.J.P. is a fellow of the Jane Coffin Childs Memorial Fund for Medical Research, and this investigation has been supported in part by the Jane Coffin Childs Memorial Fund for Medical Research.

Supporting Information Available: Experimental procedures and full characterization data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL991230P

(30) (a) Poulter, C. D. *Acc. Chem. Res.* **1990**, 23, 70. (b) Poulter, C. D.; Capson, T. L.; Thompson, M. D.; Bard, R. S. *J. Am. Chem. Soc.* **1989**, 111, 3734.

(31) Dougherty, D. A. *Science* **1996**, 271, 163.

(32) Jenson, C.; Jorgensen, W. L. *J. Am. Chem. Soc.* **1997**, 119, 10846.