

Addition of Organozinc Species to Cyclic 1,3-Diene Monoepoxide


Song Xue,* Yali Li, Kaizhen Han, Wen Yin, Meng Wang, and Qingxiang Guo

Department of Chemistry, University of Science and Technology of China,
Hefei, 230026, P. R. China

xuesong@ustc.edu.cn

Received December 20, 2001

ABSTRACT



The reaction of organozinc reagents (ZnEt_2 , ZnPh_2) with cyclic 1,3-diene monoepoxides in the presence of CF_3COOH gave the *cis*-addition products. Lewis acids such as $(\text{CF}_3\text{CO}_2)_2\text{Zn}$ and ZnCl_2 mediated the nucleophilic addition of ZnEt_2 to cyclooctadiene monoepoxide with high stereoselectivity.

Vinyl epoxides are valuable synthetic building blocks, and their reactions have been extensively investigated.¹ The addition of organometallic reagents to vinyl epoxides is a fundamental preparation of allylic and homoallylic alcohols.² Grignard reagents,³ allylstannanes,⁴ and alkyllithiums⁵ generally provide 1,2-addition products, and copper reagents⁶ and copper-catalyzed organozinc reagents⁷ yield 1,4-addition products. Mixtures of regio- and/or stereoisomers are often obtained in favor of a *trans* relationship between the hydroxy group and the substituent. In contrast, the *syn* addition of carbon nucleophiles toward epoxide has received less attention.⁸

Recently, Zaidlewicz and Krzeminski reported that 1,2-*syn* addition was favored in the reaction of allyldiethylborane

with six-, seven-, and eight-membered 1,3-diene monoepoxide.⁹ The ratio of *cis/trans*-1,2-addition was up to 72/28 when 3,4-epoxycyclohexene reacted in ether at room temperature. α -C-Glycosides were also synthesized by the *syn* addition of aluminum or boron reagents to glycal epoxides.¹⁰ We now wish to report a stereoselective addition of organozinc species to cyclic 1,3-diene monoepoxides.

Our investigation started with 1,3-cyclooctadiene monoepoxide **1c** as a substrate. The active organozinc species XZnR could be generated by reaction of ZnR_2 ($\text{R} = \text{Et}, \text{Ph}$) with acids (HX). When a solution of organozinc $\text{CF}_3\text{CO}_2\text{ZnEt}$ and substrate **1c** in CH_2Cl_2 was stirred at 0 °C for 2 h,

(9) They found a novel rearrangement reaction to occur in the reaction of allyldiethylborane with 5- membered rings. Zaidlewicz, M.; Krzeminski, M. P. *Org. Lett.* **2000**, 2, 3897.

(10) Rainier, J. D.; Cox, J. M. *Org. Lett.* **2000**, 2, 2707.

(11) **Addition of $\text{CF}_3\text{COOZnEt}$ to **1c**.** To a solution of ZnEt_2 (1 M in *n*-hexane, 1.2 mL, 1.2 mmol) in 2 mL of CH_2Cl_2 at 0 °C was added CF_3COOH (92 μL , 1.2 mmol) very slowly via syringe under N_2 . After 30 min of stirring, a solution of **1c** (124 mg, 1.0 mmol) in CH_2Cl_2 (1 mL) was added. The mixture was stirred for 2 h at 0 °C and then quenched with saturated aqueous NH_4Cl . The mixture was extracted with Et_2O (3×10 mL), washed with brine, dried (Na_2SO_4), and concentrated. Column chromatography afforded 102 mg (66%) of homoallylic alcohol *cis*-**2c**. GC–MS analysis of the crude reaction mixture revealed three peaks: **1c** (8%), 3-cycloocten-1-one of the rearrangement product from epoxide (19%), and *cis*-**2c** (73%).

(12) The stereochemistry of *cis*-**2c** was assigned from a similarity of its NMR spectrum to that of the known *cis*-2-allyl-3-cycloocten-1-ol in ref 9.

(13) Crystal data of *cis*-**2c**: $\text{C}_{10}\text{H}_{18}\text{O}$, crystal system, tetragonal; $a = 22.583(2)$ Å, $b = 22.583(2)$ Å, $c = 7.6273(9)$ Å, $V = 3889.9(7)$ Å³; space group, $I4(1)/a$; $Z = 16$; $F(000) = 1376$; $\mu = 0.065$ mm⁻¹; full matrix least-squares refinement on F^2 ; residuals, $R = 0.042$, $wR_2 = 0.0826$.

(1) (a) Trost, B. M.; McEachern, E. J.; Toste, F. D. *J. Am. Chem. Soc.* **1998**, 120, 12702. (b) Johannes, C. W.; Visser, M. S.; Weatherhead, G. S.; Hoveyda, A. H. *J. Am. Chem. Soc.* **1998**, 120, 8340. (c) Caldwell, C. G.; Derguini, F.; Bigge, C. F.; Chen, A. H.; Hu, S.; Wang, J.; Sastry, L.; Nakanishi, K. *J. Org. Chem.* **1993**, 58, 3533. (d) Tueting, D. R.; Echavarren, A. M. and Stille, J. K. *Tetrahedron* **1989**, 45, 979.

(2) Gorzynski-Smith, J. *Synthesis* **1984**, 629.

(3) (a) Söderberg, B. C.; Austin, L. R.; Davis, C. A.; Nyström, J. E.; Vagborg, J. O. *Tetrahedron* **1994**, 50, 61. (b) Bloodworth, A. J.; Curtis, R. J.; Spencer, M. D.; Tallant, N. A. *Tetrahedron* **1993**, 49, 2729.

(4) Naruta, Y.; Maruyama, K. *Chem. Lett.* **1987**, 963.

(5) Wieland, D. M.; Johnson, C. R. *J. Am. Chem. Soc.* **1971**, 93, 3047.


(6) (a) Marshall, J. A. *Chem. Rev.* **1989**, 89, 1503. (b) Marshall, J. A.; Crute, T. D., III; Hsi, J. D. *J. Org. Chem.* **1992**, 57, 115.

(7) (a) Lipshutz, B. H.; Woo, K.; Gross, T.; Buzard, D. J.; Tirado, R. *Synlett* **1997**, 477. (b) Badalassi, F.; Crotti, P.; Macchia, F.; Pineschi, M.; Arnold, A.; Feringa, B. L. *Tetrahedron Lett.* **1998**, 39, 7795.

(8) Trost, B. M.; Molander, G. A. *J. Am. Chem. Soc.* **1981**, 103, 5969.

only 1,2-addition product *cis*-**2c** was obtained with 66% yield (Table 1).¹¹ The relative stereochemistry between the hydroxy

Table 1. Addition of ZnEt₂ to Epoxides in the Presence of Acids



entry	epoxide	solvent	acid (HX)	yield ^a (%)
1	1c	CH ₂ Cl ₂	CF ₃ CO ₂ H	66
2	1c	toluene	CF ₃ CO ₂ H	52
3	1c	<i>n</i> -hexane	CF ₃ CO ₂ H	7
4	1c	Et ₂ O	CF ₃ CO ₂ H	0
5	1c	CH ₂ Cl ₂	CCl ₃ CO ₂ H	20
6	1c	CH ₂ Cl ₂		0
7	1a	CH ₂ Cl ₂	CF ₃ CO ₂ H	58
8	1b	CH ₂ Cl ₂	CF ₃ CO ₂ H	63

^a Isolated yields.

group and the ethyl group was assigned on the basis of the proton NMR spectrum.¹² The relative configuration was further confirmed as *cis* by X-ray analysis of the crystal structure (Figure 1).¹³ The *trans* isomer of the 1,2-addition




Figure 1. Crystal structure of *cis*-**2c**.


product was not observed by ¹H NMR. The choice of the solvent was crucial to the success of the reaction. CH₂Cl₂ and toluene were suitable solvents. The use of a more nonpolar solvent such as hexane gave a 7% yield of *cis*-**2c**. However, Et₂O failed to result in the reaction as a result of coordination to Lewis acidic zinc species thereby attenuating its reactivity. The reactivity of the organozinc species XZnEt was highly dependent on the X group. The species CCl₃-CO₂ZnEt, in place of CF₃CO₂ZnEt, only afforded a 20% yield using similar reaction conditions. In addition, when acid HX was anhydrous TsOH or CH₃COOH, no reaction occurred after stirring for 2 h at 0 °C.

Similarly, the reaction of CF₃CO₂ZnEt with cyclopentadiene and 1,3-cyclohexadiene monoepoxides gave the same

type of *cis*-1,2-addition with 56% and 63% yield, respectively. These results obtained show that ZnEt₂ is an effective nucleophile to cyclic 1,3-diene monoepoxide in the presence of CF₃COOH.¹⁴ The byproducts of these reactions were the rearrangement products of monoepoxides of cyclic dienes, suggesting that organozinc species XZnEt have Lewis acid character in these reactions.

In addition to ZnEt₂, ZnPh₂ underwent the same reaction to give exclusively **3c** in 74% yield when CF₃CO₂ZnPh reacted with **1c** in CH₂Cl₂ at 0 °C for 5 h. However, cyclopentadiene and 1,3-cyclohexadiene monoepoxides gave both 1,2- and 1,4-regioisomers when reacting with CF₃CO₂-ZnPh (Table 2). The 1,4-regioisomer was assigned from the

Table 2. Addition of ZnPh₂ to Epoxides in the Presence of CF₃CO₂H



entry	epoxide	3/4 ^a	yield ^b (%)
1	1a	39/61	81
2	1b	25/75	79
3	1c	100	74


^a Determined by GC–MS analysis. ^b Isolated yields.

chemical shift of the carbinol proton owing to its allylic character. On the other hand, the *cis* stereochemistry of **4a**, for example, is supported by the relative value of the chemical shifts of H_c (δ 2.87) and H_d (δ 1.58). The difference between H_c and H_d is larger than 1 ppm, demonstrating that the product is *cis*-1,4-disubstituted cyclopentene.¹⁵

Encouraged by this initial result, the reaction was investigated using Lewis acids in place of acid HX, since organozinc species XZnEt could be formed by the reaction of ZnEt₂ with ZnX₂. As shown in Table 3, 60% yield of **2c** could be obtained when ZnEt₂ reacts with (CF₃CO₂)₂Zn, which is formed in situ by the reaction of ZnEt₂ with CF₃-COOH. Analysis of the crude reaction mixture using GC–MS shows that a small amount of 1,4-addition product is always obtained under the reaction conditions used. It is worth noting that ZnCl₂ was found to be reactive under these reaction conditions, affording good yield (78%) of **2c** but with a mixture of stereoisomer. The same reaction carried out with ZnBr₂ gave the homoallylic alcohol **2c** in a low yield (39%) with a decrease in both stereoselectivity and regioselectivity.

(14) Shi reported that CF₃CO₂H accelerated the cyclopropanation of olefins. Yang, Z. Q.; Lorenz, J. C.; Shi, Y. *Tetrahedron Lett.* **1998**, 39, 8621.

(15) Marino, J. P.; de la Pradilla, R. F.; Laborde, E. *J. Org. Chem.* **1987**, 54, 4898.


Table 3. Addition of ZnEt_2 to **1c** Mediated by Lewis Acids


entry	ZnEt_2 (equiv)	Lewis acid (equiv)	yield 2c ^a (<i>cis/trans</i>) ^b	2c/5c ^c
1	0.6	$(\text{CF}_3\text{CO}_2)_2\text{Zn}$ (0.6)	60 (100)	99/1
2	0.6	$(\text{CF}_3\text{CO}_2)_2\text{Zn}$ (1.0)	53 (100)	98/2
3	1.2	$(\text{CF}_3\text{CO}_2)_2\text{Zn}$ (0.6)	36 (100)	98/2
4	1.2	$(\text{CF}_3\text{CO}_2)_2\text{Zn}$ (1.0)	52 (100)	98/2
5	0.6	ZnCl_2 (0.6)	26 (91/9)	97/3
6	1.2	ZnCl_2 (0.6)	52 (88/12)	94/6
7	1.2	ZnCl_2 (1.0)	78 (94/6)	94/6
8	1.2	ZnBr_2 (1.0)	39 (80/20)	89/11

^a Isolated yields. ^b Determined by ^1H NMR [*cis* (δ 3.81), *trans* (δ 3.52), CH-OH]. ^c Determined by GC-MS analysis.

The *syn* addition appears to be occurring via the coordination of zinc to oxygen of the epoxide and intramolecular transfer of R group. The mechanism is closely related to that proposed by Rainier and Cox in their work⁹ and outlined in Scheme 1. We considered the transfer of R group from XZnR occurred from the same face as the oxygen of epoxide and resulted in a *syn* addition, since organozinc reagents XZnR could act as a Lewis acid system as well as being nucleophilic in character.

In conclusion, we have demonstrated a new nucleophilic ring-opening reaction of cyclo-1,3-diene monoepoxide using

Scheme 1

organozinc species. The *cis*-addition products were obtained when reacting with $\text{CF}_3\text{CO}_2\text{ZnEt}$, which is formed by mixture of ZnEt_2 with $\text{CF}_3\text{CO}_2\text{H}$ or $(\text{CF}_3\text{CO}_2)_2\text{Zn}$. Additional studies of the scope are in progress.

Acknowledgment. We thank University of Science and Technology of China for financial support and the Natural Science Foundation of Anhui province.

Supporting Information Available: Spectroscopic data for all new products. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL017279O