

Asymmetric Synthesis of α -Amino 1,3-Dithioketals from Sulfinimines (*N*-Sulfinyl Imines). Synthesis of (2*S*,3*R*)-(–)-3-Hydroxy-3-methylproline


Franklin A. Davis,* Tokala Ramachandar, and Hu Liu

Department of Chemistry, Temple University, Philadelphia, Pennsylvania 19122

fdavis@temple.edu

Received July 20, 2004

ABSTRACT




N-Sulfinyl α -amino 1,3-dithioketals are prepared in high de and good yield by treating sulfinimines with lithio-1,3-dithianes. Selective removal of the *N*-sulfinyl or the thioketal groups affords stable α -amino 1,3-dithioketals and *N*-sulfinyl α -amino ketones, respectively. This new sulfinimine-derived chiral building block is employed in the asymmetric synthesis of polyoxypeptin amino acid (2*S*,3*R*)-(–)-3-hydroxy-3-methylproline.

The asymmetric syntheses of α -amino ketones **1** is currently receiving a great deal of attention because of the value of these building blocks in the asymmetric syntheses of heterocyclic compounds and natural products.¹ Generally enantiopure α -amino ketones are prepared by reaction of lithium and Grignard reagents with suitably *N*-protected α -amino acid derivatives; however, these syntheses are limited by the availability of the starting material.^{2,3} In addition, while α -amino ketones are more stable than α -amino aldehydes, racemization can occur on storage. The harsh conditions necessary to remove the *N*-protecting group can also be problematic.^{1,4}

There is however, another way to think about the syntheses and utilization of α -amino ketones, and that is to protect the keto group and leave the amine free for further elaboration.

The keto group is then unmasked later in the synthetic sequence. α -Amino 1,3-dithianes **2** are a class of compounds that can be used in this manner because the dithioacetal group is stable to both acidic and basic conditions (Scheme 1).⁵

Scheme 1



(1) (a) Coppola, G.; Schuster, H. *Asymmetric Synthesis: Construction of Chiral Molecules Using Amino Acids*; Wiley: New York, 1987. (b) Reetz, M. T. *Angew. Chem., Int. Ed. Engl.* **1991**, *30*, 1531. (c) Sardina, F. J.; Rapoport, H. *Chem. Rev.* **1996**, *96*, 1825.

(2) For leading references to the asymmetric synthesis of α -amino ketones, see: Paleo, M. R.; Calaza, M. I.; Sardina, F. J. *J. Org. Chem.* **1997**, *62*, 6862.

(3) For an alternative method, see: Myers, A. G.; Yoon, T. *Tetrahedron Lett.* **1995**, *36*, 9429.

(4) For reviews on α -amino aldehydes and ketones, see: (a) Jurczak, J.; Golebiowski, A. *Chem. Rev.* **1989**, *89*, 149. (b) Fisher, L. E.; Muchowski, J. M. *Org. Prep. Proced. Int.* **1990**, *22*, 399.


Although racemic α -amino 1,3-dithianes **2** have been described,⁶ the only optically active examples were those prepared from D-glucosamine.⁷ Furthermore, in only one instance, a racemic example, was the 1,3-dithiane success-

(5) For a review of the application of 1,3-dithianes in natural product synthesis, see: Yus, M.; Najera, C.; Foubelo, F. *Tetrahedron* **2003**, *59*, 6147.

(6) (a) Wan, Y.; Angleson, J. K.; Kutateladze, A. G. *J. Am. Chem. Soc.* **2002**, *124*, 5610. (b) Mitkin, O. D.; Kurchan, A. N.; Wan, Y.; Schiwal, B. F.; Kutateladze, A. G. *Org. Lett.* **2001**, *3*, 1841. (c) Page, P. C. B.; van Niel, M. B.; Westwood, D. J. *Chem. Soc., Chem. Commun.* **1987**, 775.

fully deprotected to give the ketone moiety.⁸ We describe here a general method for the asymmetric synthesis of α -amino 1,3-dithianes **2** and the utility of this new sulfinimine-derived chiral building block for the asymmetric synthesis of α -amino ketones **1** and the polyoxypeptin amino acid (2*S*,3*R*)-(-)-3-hydroxy-3-methylproline (**14**).

Synthesis of α -Amino-1,3-dithianes. Addition of 1.5 equiv of a preformed -78°C THF solution of 2-lithio-2-methyl-1,3-dithiane to sulfinimines (*S*)-(+)-*N*-benzylidene-*p*-toluenesulfinamide (**3a**) and (*S*)-(+)-*N*-isobutylidene-*p*-toluenesulfinamide (**3b**) readily gave the corresponding *N*-sulfinyl α -amino-1,3-dithianes (*S*_S,*S*)-(+)-**4a** and (*S*_S,*S*)-(+)-**4b** (Scheme 2).^{9,10} The diastereoselectivities, determined




by ^1H NMR on the crude reaction mixtures, were excellent (92–97% de), and the yields of the major diastereoisomers, isolated by flash chromatography, were very good (76–84%).

Selective removal of the *N*-sulfinyl group in (+)-**4** can be accomplished in several ways. Treatment of the *N*-sulfinyl α -amino 1,3-dithianes **4** with the Dess–Martin periodinane (DMP) reagent afforded the corresponding amino 1,3-dithianes (*S*)-(+)-**5a** and (*S*)-(+)-**5b** in 76 and 71% yield, respectively (Scheme 2).¹¹ The usual method for removing

N-sulfinyl groups in sulfinamides, TFA/ H_2O , also works (see below).⁸ A chiral shift reagent experiment indicated that the C–N stereocenter in (+)-**5a** was not compromised in the deprotection step.

Unless α -amino ketones are suitably *N*-protected, they are unstable and generally cannot be isolated.^{1,4} Consequently, a method for the selective hydrolysis of the dithioacetal moiety, in the presence of the *N*-sulfinyl group in (+)-**4**, is required if *N*-sulfinyl α -amino 1,3-dithianes are to be useful chiral building blocks (Scheme 1). It is unlikely that the *N*-sulfinyl group would survive most of the traditional methods for thioacetal hydrolysis, i.e., mercury(II) salts and oxidative conditions.¹² However, we found that bis(trifluoroacetoxy)iodobenzene [$\text{PhI}(\text{O}_2\text{CCF}_3)_2$], described earlier by Stork and Zhao for selective thioacetal hydrolysis, proved to be ideal for this purpose.¹³ Treatment of **4a,b** with 2 equiv of $\text{PhI}(\text{O}_2\text{CCF}_3)_2$ in acetonitrile–water afforded methyl ketones (*S*_S,*S*)-(-)-**6a** and (*S*_S,*S*)-(-)-**6b** in 72 and 60% isolated yields (Scheme 2). The fact that racemization was not detected in **6** further illustrates the unique nitrogen protecting group abilities of the sulfinyl group.⁹

Chelation-control arguments, based on six-membered chairlike transition states where the metal ion is chelated to the sulfinyl oxygen, have been used to rationalize the chiral recognition for the addition of enolates, Grignard reagents, DIBAL-H, and Et_2AlCN to sulfinimines.^{9d,14} On the other hand, steric arguments have been evoked to explain the stereochemical preference in additions of benzyl Grignard, α -metallo phosphonates, 1,3-dipoles, and glycine iminoester enolates to sulfinimines.^{9d,14} The absolute configuration of the newly created stereogenic center in (+)-**4** was determined to be (*S*)- by *m*-CPBA oxidation of **6a** to (*S*)-(+)-**7**, which has a known absolute configuration (Scheme 3).¹⁵ This



assignment means that 2-methyl-2-lithio-1,3-dithiane additions to sulfinimines fall into the chelation-control category and may go through a transition state such as **TS-1** (Scheme 3).

(11) Langille, N. F.; Dakin, L. A.; Panek, J. S. *Org. Lett.* **2003**, *4*, 575. These workers reported the used DMP as a selective method for removal of thioacetals and thioacetals.

(7) (a) Barton, D. H. R.; Gateau-Olesker, A.; Anaya-Mateos, J.; Cleophax, J.; Gero, S. D.; Chironi, A.; Riche, C. *J. Chem. Soc., Perkin Trans. 1* **1990**, 3211. (b) Anaya, J.; Barton, D. H. R.; Gero, S. D.; Grande, M.; Hernando, J. I. M.; Laso, N. M. *Tetrahedron: Asymmetry* **1995**, *6*, 609.


(8) Padwa, A.; Dharan, M.; Smolanoff, J.; Wetmore, S. I. *J. Am. Chem. Soc.* **1973**, *95*, 1954.

(9) For reviews on the chemistry of sulfinimines, see: (a) Zhou, P.; Chen, B.-C.; Davis, F. A. In *Advances in Sulfur Chemistry*; Rayner, C. M., Ed.; JAI Press: Stamford, CT, 2000; Vol. 2, pp 249–282. (b) Davis, F. A.; Zhou, P.; Chen, B.-C. *Chem. Soc. Rev.* **1998**, *27*, 13. (c) Ellman, J. A.; Owens, T. D.; Tang, T. P. *Acc. Chem. Res.* **2002**, *35*, 984. (d) Zhou, P.; Chen, B.-C.; Davis, F. A. *Tetrahedron* **2004**, *60*, in press.

(10) Davis, F. A.; Zhang, Y.; Andemichael, Y.; Fang, T.; Fanelli, D. L.; Zhang, H. *J. Org. Chem.* **1999**, *64*, 1403.

(-)-3-Hydroxy-3-methylproline. The aim of new synthetic methodology is to solve important problems in synthesis. To highlight the utility of our new chiral building block, we employed an α -amino 1,3-dithiane in the asymmetric synthesis of (2*S*,3*R*)-(-)-3-hydroxy-3-methylproline (**14**). This nonproteinogenic amino acid is a key component of the polyoxypeptins, a 19-membered cyclic hexadepsipeptide antibiotic reported to be a potent inducer of apoptosis in human pancreatic adenocarcinoma ASPC-1 cells.^{16,17}

Our synthetic strategy was to use the phenyl group as a masked carboxylic acid and to construct the pyrrolidine heterocycle directly from a substituted dithiane and benzaldehyde-derived sulfinimine (*R*)-(-)-**3a**. To this end, a preformed -78 °C solution of 1.5 equiv of dilithio species **8**, prepared from 2-(2-hydroxyethyl)-1,3-dithiane,¹⁸ was added to (*R*)-(-)-**3a** at -78 °C (Scheme 4). Addition of Ts-Cl and quenching gave pyrrolidine (-)-**9** in 91% de and 70% isolated yield of the major diastereoisomer for the single-flask reaction. Next, the *N*-sulfinyl group was removed (TFA/H₂O) and replaced with a tosyl group in one operation. The thioketal in (+)-**10** was hydrolyzed with PhI(O₂CCF₃)₂ affording the 3-oxo pyrrolidine (*R*)-(-)-**11** in 78% yield. The ketone, on treatment with excess methylmagnesium bromide, gave alcohol (-)-**12** as a single isomer in 85% yield. Oxidation of the phenyl group to the carboxylic acid (-)-**12** was easily affected by RuCl₃/NaIO₄. To facilitate isolation, the acid was immediately converted to the methyl ester (2*S*,3*R*)-(-)-**13** with TMSCHN₂. Attempts to remove the *N*-tosyl group in the crude acid resulted in low conversions to (-)-**14** (20–30%). Alternatively, the somewhat longer sequence involving hydrolysis of the methyl ester in (-)-**13** and then reductive removal of the *N*-tosyl group with Na/liquid NH₃ gave (2*S*,3*R*)-(-)-3-hydroxy-3-methylproline (**14**) in 72% yield (Scheme 4).¹⁹ Recognizing that a number of



the 10 steps are one-flask transformations, we consider our synthesis of this amino acid to be one of the most concise to date.

In summary, sulfinimines and lithio-1,3-dithianes afford *N*-sulfinyl- α -amino-1,3-thianes **4** in high de and good yields. This new sulfinimine-derived chiral building block can be selectively deprotected to give either α -amino-1,3-dithianes **5** or *N*-sulfinyl α -amino ketones **6** as stable entities. The synthetic utility of *N*-sulfinyl- α -amino-1,3-thianes is illustrated by the asymmetric synthesis of the polyoxypeptins amino acid (2*S*,3*R*)-(-)-3-hydroxy-3-methylproline (**14**).

Acknowledgment. We thank the National Institute of General Medical Sciences for generous support of this work (GM 57870 and GM51982).

Supporting Information Available: Experimental procedures and spectroscopic data for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL0485971

(12) Greene, T. W.; Wuts, P. G. M. *Protecting Groups in Organic Synthesis*, 3rd ed.; Wiley: New York, 1999; p 297.

(13) Stork, G.; Zhao, K. *Tetrahedron Lett.* **1989**, 30, 287.

(14) For leading references, see: (a) Davis, F. A. Lee, S.; Yan, H.; Titus, D. D. *Org. Lett.* **2001**, 3, 1757. (b) Reference 9d of this article.

(15) Adam, W.; Roschmann, K. J.; Saha-Moller, C. R. *Eur. J. Org. Chem.* **2000**, 557.

(16) Umezawa, K.; Nakasawa, K.; Uchihata, Y.; Otsuka, M. *Adv. Enzyme Regul.* **1999**, 39, 145.

(17) For earlier asymmetric syntheses of this amino acid, see: (a) Noguchi, Y.; Uchiro, H.; Yamada, T.; Kobayashi, S. *Tetrahedron Lett.* **2001**, 42, 5253. (b) Makino, K.; Kondoh, A.; Hamada, Y. *Tetrahedron Lett.* **2002**, 43, 4695. (c) Qin, D.-G.; Zha, H.-Y.; Yao, Z.-J. *J. Org. Chem.* **2002**, 67, 1038. (d) Shen, J.-W.; Qin, D.-G.; Zhang, H.-W.; Yao, Z.-J. *J. Org. Chem.* **2003**, 68, 7479. (e) Merio, P.; Revuelta, J.; Tejero, T.; Cicchi, S.; Goti, A. *Eur. J. Org. Chem.* **2004**, 776.

(18) Seebach, D.; Jones, N. R.; Corey, E. J. *J. Org. Chem.* **1968**, 33, 300.

(19) Properties of (-)-**14** were consistent with literature values. See ref 17d of this article.