

Suzuki–Miyaura Cross-Coupling of Benzylic Carbonates with Arylboronic Acids


Ryoichi Kuwano* and Masashi Yokogi

Department of Chemistry, Graduate School of Sciences, Kyushu University,
6-10-1 Hakozaki, Higashi-ku, Fukuoka 812-8581, Japan

rkuwascc@mbbox.nc.kyushu-u.ac.jp

Received January 14, 2005

ABSTRACT



The cross-coupling of benzylic carbonates with arylboronic acids gave the corresponding diarylmethanes in high yields by use of the palladium catalyst generated in situ from $[\text{Pd}(\eta^3\text{-C}_3\text{H}_5\text{Cl})_2]$ and 1,5-bis(diphenylphosphino)pentane (DPPFPent). The Suzuki–Miyaura reaction using DPPFPent–palladium catalyst is applicable to syntheses of a broad range of functionalized diarylmethanes.

During the last several years, tremendous progress has been made in palladium-catalyzed cross-couplings of organometallic compounds with unactivated aryl chlorides,¹ which had been regarded as unsuitable electrophilic coupling partners. The catalytic cross-coupling can now be conducted even with alkyl chloride.² However, the use of carboxylate functionality as a leaving group on an electrophilic substrate is unexplored except for the reactions of allylic esters.³ The acyl C–O bond of carboxylate is usually cleaved by palladium(0) complex (path a), and aryl carboxylates are often employed in the catalytic acylation of organometallic compounds (Figure 1).⁴ The oxidative addition of the C–O

formation of **A** through path b has been reported in the reaction of benzyl trifluoroacetate with palladium(0) complex.^{5,6}

Recently, we reported catalytic benzylic substitutions of benzylic carbonates, where a DPPF-ligated palladium complex was found to be the best catalyst.⁷ The catalytic reaction may involve the formation of (η^3 -benzyl)palladium through path b, and it suggests that the DPPF–palladium catalyst is efficient for the activation of the benzylic C–O bond. The




Figure 1. Oxidative Addition of Carboxylate to Palladium(0).

bond leading to (alkyl)(carboxylato)palladium **A** (path b) is desirable for the development of the cross-coupling using an ester, but the C–O bond is difficult to activate by a low-valent metal complex in general. Only one example of the

(1) Reviews: (a) Littke, A. F.; Fu, G. C. *Angew. Chem., Int. Ed.* **2002**, *41*, 4176–4211. Examples: (b) Littke, A. F.; Fu, G. C. *Angew. Chem., Int. Ed.* **1998**, *37*, 3387–3388. (c) Old, D. W.; Wolfe, J. P.; Buchwald, S. L. *J. Am. Chem. Soc.* **1998**, *120*, 9722–9723. (d) Kataoka, N.; Shelby, Q.; Stambuli, J. P.; Hartwig, J. F. *J. Org. Chem.* **2002**, *67*, 5553–5566.

(2) (a) Kirchhoff, J. H.; Dai, C.; Fu, G. C. *Angew. Chem., Int. Ed.* **2002**, *41*, 1945–1947. (b) Frisch, A. C.; Shaikh, N.; Zapf, A.; Beller, M. *Angew. Chem., Int. Ed.* **2002**, *41*, 4056–4059. (c) Zhou, J.; Fu, G. C. *J. Am. Chem. Soc.* **2003**, *125*, 12527–12530. (d) Frisch, A. C.; Rataboul, F.; Zapf, A.; Beller, M. *J. Organomet. Chem.* **2003**, *687*, 403–409.


(3) Examples: (a) Del Valle, L.; Stille, J. K.; Hegedus, L. S. *J. Org. Chem.* **1990**, *55*, 3019–3023. (b) Uozumi, Y.; Danjo, H.; Hayashi, T. *J. Org. Chem.* **1999**, *64*, 3384–3388. (c) Bouyssi, D.; Gerusz, V.; Balme, G. *Eur. J. Org. Chem.* **2002**, 2445–2448.

(4) (a) Kakino, R.; Shimizu, I.; Yamamoto, A. *Bull. Chem. Soc. Jpn.* **2001**, *72*, 371–376. (b) Tatamidani, H.; Kakiuchi, F.; Chatani, N. *Org. Lett.* **2004**, *6*, 3597–3599.

(5) Nagayama, K.; Shimizu, I.; Yamamoto, A. *Bull. Chem. Soc. Jpn.* **1999**, *72*, 799–803.

(6) Narahashi, H.; Yamamoto, A.; Shimizu, I. *Chem. Lett.* **2004**, *33*, 348–349.

Table 1. Effect of Phosphine–Palladium Complex on the Cross-Coupling of Benzyl Methyl Carbonate with Phenylboronic Acid^a



| entry | ligand | [Pd] | yield, % ^b |
|-------|------------------|--------------------------------|-----------------------|
| 1 | DPPF | $[Pd(\eta^3-C_3H_5)(cod)]BF_4$ | 3 |
| 2 | PPh_3 | $[Pd(\eta^3-C_3H_5)(cod)]BF_4$ | 2 |
| 3 | $P(C_6H_4-o-Me)$ | $[Pd(\eta^3-C_3H_5)(cod)]BF_4$ | 0 |
| 4 | $P(t-Bu)_3$ | $[Pd(\eta^3-C_3H_5)(cod)]BF_4$ | 0 |
| 5 | DPPBz | $[Pd(\eta^3-C_3H_5)(cod)]BF_4$ | 1 |
| 6 | DPPE | $[Pd(\eta^3-C_3H_5)(cod)]BF_4$ | 3 |
| 7 | DPPP | $[Pd(\eta^3-C_3H_5)(cod)]BF_4$ | 1 |
| 8 | DPPB | $[Pd(\eta^3-C_3H_5)(cod)]BF_4$ | 24 |
| 9 | DPPPent | $[Pd(\eta^3-C_3H_5)(cod)]BF_4$ | 32 |
| 10 | DPPHex | $[Pd(\eta^3-C_3H_5)(cod)]BF_4$ | 13 |
| 11 | DPPPent | $PdBr_2(cod)$ | 16 |
| 12 | DPPPent | $Pd(OAc)_2$ | 23 |
| 13 | DPPPent | $Pd(dba)_2$ | <1 |
| 14 | DPPPent | $[Pd(\eta^3-C_3H_5)Cl]_2$ | 50 (89 ^c) |

^a Reactions were conducted in DMF (1.0 mL) at 80 °C for 3 h. The ratio of **1a** (0.2 mmol):**2a**: K_2CO_3 : $[Pd]$:ligand was 20:30:60:1:1. ^b GC yield (average of two runs). ^c GC yield at 24 h.

above consideration prompted us to develop a new class of cross-couplings using benzylic carbonates as electrophiles. In this paper, we report the Suzuki–Miyaura cross-coupling of benzylic carbonates.^{8,9} The catalytic reaction offered a new straightforward approach to diverse diarylmethanes,¹⁰ which is a ubiquitous structural constituent of pharmacologically interesting compounds.¹¹

First, we attempted the cross-coupling of benzyl methyl carbonate (**1a**) with phenylboronic acid (**2a**) in the presence of $DPPF-[Pd(\eta^3-C_3H_5)(cod)]BF_4$, which was the most

(7) (a) Kuwano, R.; Kondo, Y.; Matsuyama, Y. *J. Am. Chem. Soc.* **2003**, *125*, 12104–12105. (b) Kuwano, R.; Kondo, Y. *Org. Lett.* **2004**, *6*, 3545–3547.


(8) Reviews: (a) Miyaura, N.; Suzuki, A. *Chem. Rev.* **1995**, *95*, 2457–2483. (b) Bellina, F.; Carpita, A.; Rossi, R. *Synthesis* **2004**, 2419–2440.

(9) Examples of cross-coupling of benzyl halides with organoboron compounds: (a) Chowdhury, S.; Georgiou, P. *Tetrahedron Lett.* **1999**, *40*, 7599–7603. (b) Langle, S.; Abarbri, M.; Duchêne, A. *Tetrahedron Lett.* **2003**, *44*, 9255–9258. (c) Chahen, L.; Doucet, H.; Santelli, M. *Synlett* **2003**, 1668–1672.

(10) Examples of diarylmethane synthesis through cross-coupling: (a) Negishi, E.-i.; King, A. O.; Okukado, N. *J. Org. Chem.* **1977**, *42*, 1821–1823. (b) Vanier, C.; Lorgé, F.; Wagner, A.; Mioskowski, C. *Angew. Chem., Int. Ed.* **2000**, *39*, 1679–1683. (c) Molander, G. A.; Ito, T. *Org. Lett.* **2001**, *3*, 393–396. (d) Itami, K.; Mineno, M.; Kamei, T.; Yoshida, J. *Org. Lett.* **2002**, *4*, 3635–3638.

(11) (a) Wai, J. S.; Egbertson, M. S.; Payne, L. S.; Fisher, T. E.; Embrey, M. W.; Tran, L. O.; Melamed, J. Y.; Langford, H. M.; Guare, J. P., Jr.; Zhuang, L.; Grey, V. E.; Vacca, J. P.; Holloway, M. K.; Naylor-Olsen, A. M.; Hazuda, D. J.; Felock, P. J.; Wolfe, A. L.; Stillmock, K. A.; Schleif, W. A.; Gabryelski, L. J.; Young, S. D. *J. Med. Chem.* **2000**, *43*, 4923–4926. (b) Long, Y.-Q.; Jiang, X.-H.; Dayam, R.; Sanchez, T.; Shoemaker, R.; Sei, S.; Neamati, N. *J. Med. Chem.* **2004**, *47*, 2561–2573.

Table 2. Cross-Coupling of Benzylic Carbonates (**1**) with Arylboronic Acids (**2**)^a



| entry | 1 | 2 | time, h | product (3) | yield, % ^b |
|-----------------|-----------|-----------|---------|----------------------|-----------------------|
| 1 ^c | 1a | 2a | 24 | 3a | 80 |
| 2 ^c | 1b | 2a | 5 | 3b | 88 |
| 3 | 1c | 2a | 5 | 3c | 94 |
| 4 | 1d | 2a | 12 | 3d | 94 |
| 5 | 1e | 2a | 3 | 3e | 82 |
| 6 | 1f | 2a | 24 | 3f | 84 |
| 7 | 1g | 2a | 5 | 3g | 90 |
| 8 | 1a | 2b | 24 | 3c | 90 |
| 9 | 1a | 2c | 24 | 3d | 81 |
| 10 | 1c | 2d | 24 | 3h | 99 |
| 11 | 1a | 2e | 48 | 3i | 87 |
| 12 | 1c | 2f | 72 | 3j | 85 |
| 13 | 1c | 2g | 72 | 3k | 78 |
| 14 | 1c | 2h | 3 | 3l | 92 |
| 15 | 1c | 2i | 24 | 3m | 92 |
| 16 ^c | 1a | 2j | 72 | 3n | 83 |
| 17 | 1c | 2k | 48 | 3o | 82 |


^a Reactions were conducted in DMF (1.0 mL) at 80 °C. The ratio of **1** (1.0 mmol):**2**: K_2CO_3 : $[Pd(\eta^3-C_3H_5)Cl]_2$:DPPPent was 100:110:220:0.5:1. ^b Isolated yield. ^c Reactions were conducted with 2 mol % palladium.

effective catalyst for the benzylation of malonate carbanions by benzylic carbonate.^{7a} However, the reaction failed to yield diphenylmethane (**3a**). We evaluated a broad range of phosphine ligands, palladium catalyst precursors, bases, and solvents. Selected results are shown in Table 1. The palladium catalyst generated in situ from $[Pd(\eta^3-C_3H_5)Cl]_2$ and bidentate bisphosphine 1,5-bis(diphenylphosphino)pen-

tane (DPPent) recorded the highest yield of **3a**. The choice of a phosphine ligand is essential for the catalysis. Mono-phosphine-ligated palladium complexes exhibited poor catalytic activity for the cross-coupling. Bidentate phosphine ligands possessing a small bite angle were ineffective, while increasing the ligand bite angle enhanced the reaction rate and the yield of **3a**. The yield of **3a** reached a maximum when DPPent was used. DPPHex, providing a larger P–Pd–P angle, was less effective than DPPent.¹² [Pd(η^3 -C₃H₅)Cl]₂ was superior to the other palladium catalyst precursors. DMF was the solvent of choice. K₃PO₄ and Cs₂CO₃, as well as K₂CO₃, were suitable bases for the cross-coupling of **1a** with **2a**, but K₂CO₃ is economically preferred.¹³ The catalyst loading was successfully reduced to 2 mol % palladium at a high concentration of **1a** (1.0 M) under the optimal reaction conditions, and **3a** was obtained in 80% isolated yield (Table 2, entry 1).


The optimal palladium catalyst and reaction conditions proved to be useful for cross-couplings of diverse benzylic carbonates and boronic acids, as shown in Table 2. Both electron-rich and electron-poor benzylic esters, **1c** and **1d**, respectively, were more reactive than **1a**,¹⁴ and their reactions with **2a** afforded diarylmethane **3c** and **3d** in high yields at a catalyst loading of 1 mol % palladium. On the other hand, the rate of the reaction of electron-rich boronic acid **2b** was almost equal to that of **2a**, but the electron-withdrawing group on arylboronic acid **2c** accelerated the cross-coupling.¹⁴ The ortho substituents of arylboronic acids **2d** and **2e** did not hinder the catalytic reaction. A variety of diarylmethanes bearing chloro, carboxylate, aldehyde, carbamate, alcohol, and nitro functionalities were prepared in high yields by means of the DPPent–palladium catalyst. Alkenylboronic acid **2k** was transformed into allylarene **3o** in high yield, and no undesirable migration of the double bond was observed. However, alkylboronic acids failed to react with **1a**, and the desired cross-coupling product was not obtained. Xylene- α,α' -diol dicarbonates **4** and **5** underwent two-directional cross-coupling with **2a**, as shown in Scheme 1, and yielded *p*- (**6**) and *m*-dibenzylbenzene (**7**), respectively.

Scheme 1. Syntheses of Dibenzylbenzenes



A proposed mechanism of the present reaction is shown in Scheme 2. The benzylic C–O bond of **1** is cleaved by

Scheme 2. Proposed Mechanism



DPPent-ligated palladium(0) **B**, and (η^1 -benzyl)(alkoxy)palladium intermediate **C** was formed through decarboxylation of the carbonate leaving group. The alkoxy ligand on palladium is more effective for transmetalation with organoboron **2** than acetato and halo ligands.¹⁵ No cross-coupling of benzyl acetate with **2a** occurred in the presence of the DPPent–palladium catalyst. The resulting (alkyl)(benzyl)palladium **D** provides diarylmethane **3** and regenerates palladium(0) **B** through reductive elimination.

In conclusion, we have proved that benzylic carbonates are a new class of coupling partners in Suzuki–Miyaura cross-coupling. The palladium catalyst generated from [Pd(η^3 -C₃H₅)Cl]₂ and bidentate bisphosphine DPPent is most effective for the cross-coupling using benzylic carbonates. The catalytic reaction showed high functional group compatibility, and a wide range of arylboronic acids are now commercially available. Therefore, this methodology developed by us will offer a fruitful and general synthetic method for diarylmethanes.

Acknowledgment. This work was supported by the Sumitomo Foundation and a Grant-in-Aid for Young Scientists (A) (No. 16685011) from MEXT.

Supporting Information Available: Experimental procedures and characterization data for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL050078Q

(12) We speculate that the cleavage of the benzylic C–O bond proceeded through a pathway similar to the formation of (η^3 -allyl)palladium from allylic ester. The activation energy of the process would depend on the potential energy of (η^3 -benzyl)palladium. The angle of P–Pd–P was 104.37° in the DFT-optimized structure (B3LYP/6-31G(d) & LANL2DZ for Pd) of [(H₃P)₂-Pd(η^3 -benzyl)]⁺. DPPent might provide a bite angle close to 104.37° and might form undistorted (η^3 -benzyl)palladium, which immediately isomerized to (η^1 -benzyl)palladium.

(13) When the reaction was conducted with 1.0 equiv of Cs₂CO₃ to **2a**, no formation of **3a** was observed. The acidic proton of boronic acid might cause hydrolysis of Pd–OR bond of the intermediate **C** in Scheme 2.

(14) Relative reactivities of **1a,c,d** and **2a,b,c** were estimated by competitive experiments using **2a** and **1a**, respectively. See Supporting Information.

(15) Miyaura, N.; Yamada, K.; Sugimoto, H.; Suzuki, A. *J. Am. Chem. Soc.* **1985**, *107*, 972–980.